```
In[4]:= (* Adam Beck *)
       (* Problem 1*)


       (* hosp-heart.nb data *)

       (* {M,V} M = one year mortality rate,
       percentage of patiuents that died within one year of the
        transplant operation,
       V = average annual number of transplants at that center during the same 4 years *)
       heart = {{17.9, 27}, {23.1, 4}, {40, 3}, {6.5, 35}, {14.9, 17}, {12.5, 4}, {15.7, 45},
           {9.8, 28}, {24, 6}, {5.0, 10}, {15.4, 13}, {4.8, 7}, {0, 1}, {19.1, 47}, {4.5, 6},
           {15, 56}, {12.5, 4}, {33.9, 8}, {10.7, 9}, {13, 14}, {28.3, 12}, {57.1, 2}, {6.3, 4},
           {10, 3}, {8.3, 12}, {17.5, 10}, {20, 3}, {29.3, 10}, {21.4, 7}, {27.3, 8}, {13.6, 6},
           {21.8, 30}, {36.4, 3}, {18.2, 11}, {33.3, 2}, {20, 4}, {38.5, 7}, {20.8, 18}, {12.2, 19},
           {22.2, 18}, {29, 8}, {0, 9}, {5.7, 9}, {50, 2}, {21.7, 15}, {66.7, 4}, {29.4, 17},
           {12.1, 27}, {10.7, 14}, {6.3, 4}, {16.2, 9}, {21.1, 5}, {17.4, 33}, {23.9, 17},
           {42.9, 2}, {40, 2}, {6.7, 15}, {44.4, 3}, {18.7, 34}, {14.7, 24}, {7.4, 7}, {12.6, 24},
           {9.7, 26}, {44.4, 2}, {16.7, 6}, {15.8, 14}, {83.3, 2}, {10.9, 22}, {13.3, 5},
           {11.1, 5}, {75, 2}, {19, 20}, {14, 13}, {60, 1}, {21.2, 8}, {9.7, 8}, {50, 2}, {25, 14},
           {18.6, 15}, {0.0, 1}, {35.3, 9}, {23.5, 85}, {15.6, 11}, {37.5, 2}, {14.3, 28},
           {14.3, 4}, {16.7, 6}, {20.0, 15}, {13.0, 17}, {9.6, 26}, {66.7, 3}, {30.8, 3},
           {14.0, 13}, {27.5, 10}, {37.5, 8}, {18.9, 13}, {0.0, 4}, {12.2, 44}, {57.1, 4},
           {21.4, 35}, {23.4, 16}, {10.9, 12}, {15.6, 8}, {16.7, 2}, {13.9, 9}, {18.2, 11},
           {11.5, 26}, {18.4, 13}, {16.7, 3}, {20.4, 14}, {40.0, 5}, {20.7, 56}, {19.6, 13},
           {13.5, 9}, {29.9, 36}, {8.4, 21}, {28.4, 24}, {7.7, 23}, {19.3, 29}, {0.0, 1},
           {22.2, 20}, {30.0, 5}, {7.0, 11}, {23.8, 7}, {18.8, 29}, {14.5, 16}, {17.0, 16},
           {20.0, 15}, {6.7, 15}, {11.4, 20}, {100.0, 1}, {31.4, 9}, {17.6, 26}, {19.6, 14}};

In[5]:= (* Split this M and V data into separate
        lists via Transpose[] in order to parse through *)
       heartTranspose = Transpose[heart];
       MData = heartTranspose[[1]];
       VData = heartTranspose[[2]];

In[8]:= (* Define mean, median, quantile, and variance functions *)

In[9]:= mean[x_] := Sum[x[[i]], {i, 1, Length[x]}] / Length[x];
       (* Sum elements, divide by length *)

In[10]:= median[x_] := (s = Sort[x]; s[[IntegerPart[.5 * Length[s]]]]);
       (* Sort list, take element at index 1/2*length *)

In[11]:= quantile[x_, alpha_] := (s = Sort[x]; s[[IntegerPart[alpha * Length[s]]]])
        (* Sort list, take element at index .alpha*length *)

In[12]:= variance[x_] := (m = mean[x]; Sum[(x[[i]] - m)^2, {i, 1, Length[x]}] / Length[x]);
       (* difference of every element from mean, squared, times 1/length *)

In[13]:= (* Find the mean, median, q1 and q3, and variance *)
       hospMeanM = mean[MData]

Out[13]= 21.9045
```

```
In[14]:= hospMeanV = N[mean[VData]]

Out[14]= 13.8657

In[15]:= hospMedianM = median[MData]

Out[15]= 18.2

In[16]:= hospMedianV = median[VData]

Out[16]= 10

In[17]:= hospQ1M = quantile[MData, .25]

Out[17]= 12.2

In[18]:= hospQ1V = quantile[VData, .25]

Out[18]= 4

In[72]:= hospQ3M = quantile[MData, .75]

Out[72]= 25

In[73]:= hospQ3V = quantile[VData, .75]

Out[73]= 17

In[21]:= hospVarianceM = variance[MData]

Out[21]= 268.634

In[22]:= hospVarianceV = N[variance[VData]]

Out[22]= 166.46

In[23]:= (* Histograms using two difference bin sizes *)
        (* I will use a bin size Length/2 for a very large bin,
        and Length/10 for a smaller bin *)

        (* MData large bin *)
        Histogram[MData, IntegerPart[Length[MData] / 2]]
```

In[24]:= `(* MData small bin *)`
`Histogram[MData, IntegerPart[Length[MData] / 10]]`

Out[24]=

In[25]:= `(* VData, large bin *)`
`Histogram[VData, IntegerPart[Length[VData] / 2]]`

Out[25]=

In[26]:= `(* VData, small bin *)`
`Histogram[VData, IntegerPart[Length[VData] / 10]]`

Out[26]=

In[27]:= `(* Product plots of quantile functions, moment functions, and CDFs *)`

`(* Define functions for moments and CDF *)`

`(* Sum elements raised to the kth power, divide by length *)`
`moment[x_, k_] := N[Sum[x[[i]]^k, {i, 1, Length[x]}] / Length[x]];`

In[28]:= `cdf[x_, xi_] := N[Sum[If[x[[i]] <= xi, 1, 0], {i, 1, Length[x]}]];`
`(* Count that an element is less than or equal to a given element *)`

In[29]:= `(* Plot the quantile functions *)`
`(* MData *)`
`Plot[quantile[MData, i], {i, 0, 1}]`

Out[29]=

In[30]:= `(* VData *)`
`Plot[quantile[VData, i], {i, 0, 1}]`

Out[30]=

In[31]:= `(* Plot moment functions *)`
`(* Get the first 10 moments for MData and VData in a Table *)`
`momMData = Table[moment[MData, i], {i, 1, 10}];`

In[32]:= `momVData = Table[moment[VData, i], {i, 1, 10}];`

In[33]:= `(* Plot the two moments *)`
`ListPlot[{momMData, momVData}, PlotStyle → PointSize[.03], PlotRange → {0, 9.9 * 10^17}]`
`(* Blue is MData, Orange is VData*)`

Out[33]=

In[34]:= `(* Plot CDF functions *)`
`(* MData *)`
`Plot[cdf[MData, i], {i, 0, Max[MData]}]`

Out[34]=

In[35]:= `(* VData *)`
`Plot[cdf[VData, i], {i, 0, Max[VData]}]`

Out[35]=

```
(* Although the instructions do not say to compare
 any box and whisker and QQ plots for this hospital data
 against other sets of data, I will product them anyways. QQ
 will be MData(x axis) against VDaya (y axis) *)

(* Box an whisker plots *)
(* A box and whisker plot takes a min, q1, q2 (median), q3, and max *)

(* Box and whisker, MData and VData, outliers not shown *)
BoxWhiskerChart[{MData, VData}]
```

Out[46]=

In[47]:=
```
(* Box and whisker, MData and VData, outliers shown *)
BoxWhiskerChart[{MData, VData}, "Outliers"]
```

Out[47]=

In[54]:= `(* Parametric Plots (QQ Plots), MData on x axis, VData on y axis *)`
`ParametricPlot[{quantile[MData, i], quantile[VData, i]}, {i, 0, 1}]`

Out[54]=

In[63]:= `(* If the two sets come from a population with the same distribution,`
`    the points should fall approximately`
`     along a 45 degree reference line. As we can see,`
`    the 2 batches do not appear to have come from populations with`
`     a common distribution, as they do not fit along a 45 degree line *)`

`(* resistor.nb data *)`
`(* The data represents a listing of the resistances`
` (in ohms) of 200 resistors which are all rated at 10 kiloohms. *)`
`resistors = {9.97910927, 9.833997401, 10.48797923, 9.778286587, 10.4127049, 9.729651074,`
`    10.34005333, 9.894176108, 10.07983211, 9.933230947, 9.977783398, 10.13141411,`
`    10.1266421, 9.37852757, 10.26785423, 9.907086669, 9.744503691, 9.971603949,`
`    9.693939764, 9.620137112, 12.28072506, 10.0580338, 10.33764317, 9.757096213,`
`    9.593230848, 9.713741738, 9.432574293, 9.62099431, 9.802732952, 9.971484578,`
`    10.22548428, 10.3352728, 9.989841592, 10.29860424, 9.52298034, 10.08499861,`
`    9.394148142, 9.944944954, 10.21438162, 10.36193691, 10.02987499, 9.603449021,`
`    9.742946181, 9.875414084, 10.05078967, 10.12314509, 10.15281111, 5.870566193,`
`    9.484863417, 9.973958404, 9.94911044, 9.374762262, 9.788310356, 10.06500849,`
`    9.77439594, 10.03864565, 10.32397119, 9.916142963, 9.967350072, 10.09860352,`
`    9.987682395, 10.15563395, 9.537918791, 9.945042157, 10.02686399, 9.74540807,`
`    10.26915708, 9.696347652, 10.13930795, 9.51572712, 9.367227099, 9.831637831,`
`    10.1807235, 9.88921993, 9.923452458, 9.944225885, 9.779727284, 10.26538836, 10.2298635,`
`    10.2461264, 9.694717951, 9.771545526, 9.679096242, 10.15118993, 10.25894345,`
`    9.613968464, 10.14607857, 10.3809408, 10.00425765, 10.30422606, 9.938641588,`
`    10.14989447, 9.62901378, 6.613345698, 10.48706974, 10.10426569, 10.15476425,`
`    9.839152246, 9.74229305, 9.712882265, 10.09355753, 9.655283966, 10.01073951,`
`    10.23032052, 9.896222755, 9.646005983, 10.22741355, 9.916736976, 9.853518852,`
`    9.797304974, 9.542975581, 9.582644329, 10.06420074, 10.1110437, 9.09833499,`
`    9.694181349, 10.0837185, 9.990310834, 9.680224016, 9.544769559, 10.12220661,`
`    10.35625939, 9.68922915, 9.816272486, 9.838797828, 9.787675983, 10.01512384,`
`    9.672549018, 9.166747182, 9.839861368, 10.0490497, 9.9589975, 9.707653239, 9.642065029,`
`    10.14670044, 9.704657023, 9.851454583, 9.92931813, 10.05903936, 9.749898131,`
`    10.12904658, 9.776733909, 9.956306817, 10.10913774, 9.25291271, 9.823820724,`
`    9.581313056, 9.84027462, 9.738894951, 9.923279654, 9.815685862, 9.754906605,`
`    10.19531748, 9.718578829, 9.830784043, 9.860661512, 9.665515781, 9.956836598,`
`    10.06308718, 9.401201273, 10.10992616, 9.738494773, 9.991823154, 9.877411846,`
`    10.23755441, 10.04556889, 9.978626954, 10.06519891, 9.774786454, 10.26202664,`
`    10.10298671, 9.558598995, 9.352852535, 9.611078544, 9.807194024, 9.684415081,`
`    10.17326848, 9.683191811, 10.03918111, 9.891267714, 9.707087079, 9.68933829,`
`    10.10867702, 9.770431111, 9.697278747, 10.15024178, 10.17638293, 9.676198933,`
`    9.765484028, 9.952918381, 10.15444308, 10.03372073, 9.607316647, 9.856609145,`
`    9.805244863, 9.728007162, 9.951510938, 10.03217857, 10.19504918, 10.23059564};`

In[70]:= `(* Find the mean, median, q1 and q3, and variance *)`
`resistorMean = mean[resistors]`

Out[70]= `9.87989`

In[71]:= `resistorMedian = median[resistors]`

Out[71]= `9.91674`

In[74]:= **resistorQ1 = quantile[resistors, .25]**

Out[74]= 9.71288

In[75]:= **resistorQ3 = quantile[resistors, .75]**

Out[75]= 10.1043

In[79]:= **resistorVariance = variance[resistors]**

Out[79]= 0.229448

```
(* Histograms using two difference bin sizes *)
(* I will use a bin size Length/2 for a very large bin,
and Length/10 for a smaller bin *)
```