

**Topic:** Distinct quadratic factors

**Question:** Use partial fractions to evaluate the integral.

$$\int \frac{x^3 - 8x^2 - 1}{(x^2 + x - 6)(x^2 + 1)} dx$$

**Answer choices:**

- A  $2 \ln |x + 3| - \ln |x - 2| - \tan^{-1} x + C$
- B  $2 \ln |x + 3| - \ln |x - 2| + \tan^{-1} x + C$
- C  $2 \ln |x + 3| - \ln |x - 2| + C$
- D  $2 \ln |x + 3| + \ln |x - 2| + C$



**Solution: A**

First, factor the denominator.

$$\int \frac{x^3 - 8x^2 - 1}{(x^2 + x - 6)(x^2 + 1)} dx = \int \frac{x^3 - 8x^2 - 1}{(x + 3)(x - 2)(x^2 + 1)} dx$$

Using partial fractions decomposition containing a quadratic factor, we have

$$\frac{x^3 - 8x^2 - 1}{(x + 3)(x - 2)(x^2 + 1)} = \frac{A}{x + 3} + \frac{B}{x - 2} + \frac{Cx + D}{x^2 + 1}$$

Now we'll solve for constants.

$$\left[ \frac{x^3 - 8x^2 - 1}{(x + 3)(x - 2)(x^2 + 1)} = \frac{A}{x + 3} + \frac{B}{x - 2} + \frac{Cx + D}{x^2 + 1} \right] (x + 3)(x - 2)(x^2 + 1)$$

$$x^3 - 8x^2 - 1 = A(x - 2)(x^2 + 1) + B(x + 3)(x^2 + 1) + (Cx + D)(x + 3)(x - 2)$$

$$x^3 - 8x^2 - 1 = A(x^3 - 2x^2 + x - 2) + B(x^3 + 3x^2 + x + 3) + (Cx + D)(x^2 + x - 6)$$

$$x^3 - 8x^2 - 1 = Ax^3 - 2Ax^2 + Ax - 2A + Bx^3 + 3Bx^2 + Bx + 3B$$

$$+ Cx^3 + Cx^2 - 6Cx + Dx^2 + Dx - 6D$$

$$x^3 - 8x^2 - 1 = (Ax^3 + Bx^3 + Cx^3) + (-2Ax^2 + 3Bx^2 + Cx^2 + Dx^2)$$

$$+ (Ax + Bx - 6Cx + Dx) + (-2A + 3B - 6D)$$

$$x^3 - 8x^2 - 1 = (A + B + C)x^3 + (-2A + 3B + C + D)x^2$$



$$+ (A + B - 6C + D)x + (-2A + 3B - 6D)$$

Equating coefficients on both sides, we get

$$\text{[1]} \quad A + B + C = 1$$

$$\text{[2]} \quad -2A + 3B + C + D = -8$$

$$\text{[3]} \quad A + B - 6C + D = 0$$

$$\text{[4]} \quad -2A + 3B - 6D = -1$$

Subtracting [4] from [2], we get

$$(-2A + 3B + C + D = -8) - (-2A + 3B - 6D = -1)$$

$$-2A + 2A + 3B - 3B + C + D + 6D = -8 + 1$$

$$\text{[5]} \quad C + 7D = -7$$

Subtracting [1] from [3], we get

$$(A + B - 6C + D = 0) - (A + B + C = 1)$$

$$A - A + B - B - 6C - C + D = 0 - 1$$

$$\text{[6]} \quad -7C + D = -1$$

Multiplying [6] by 7, we get

$$7(-7C + D = -1)$$

$$\text{[7]} \quad -49C + 7D = -7$$

Subtracting [7] from [5], we get



$$(C + 7D = -7) - (-49C + 7D = -7)$$

$$C + 49C + 7D - 7D = -7 + 7$$

$$50C = 0$$

$$C = 0$$

Once we've solved for one constant it gets easier to solve for the others.

$$C + 7D = -7$$

$$0 + 7D = -7$$

$$D = -1$$

Plugging the values for  $C$  and  $D$  into [2] and [3], we get

$$-2A + 3B + 0 - 1 = -8$$

$$A + B - 6(0) - 1 = 0$$

and these simplify to

$$\text{[8]} \quad -2A + 3B = -7$$

$$\text{[9]} \quad A + B = 1$$

Multiplying [9] by 2, we get

$$A + B = 1$$

$$2(A + B = 1)$$

$$\text{[10]} \quad 2A + 2B = 2$$



Now we can add [10] to [8] and get

$$(-2A + 3B = -7) + (2A + 2B = 2)$$

$$-2A + 2A + 3B + 2B = -7 + 2$$

$$5B = -5$$

$$B = -1$$

Plugging  $B = -1$  into  $A + B = 1$ , we get

$$A + B = 1$$

$$A - 1 = 1$$

$$A = 2$$

With values for all of the constants, we'll plug into the partial fractions decomposition.

$$\frac{x^3 - 8x^2 - 1}{(x + 3)(x - 2)(x^2 + 1)} = \frac{2}{x + 3} + \frac{-1}{x - 2} + \frac{0x - 1}{x^2 + 1}$$

$$\frac{x^3 - 8x^2 - 1}{(x + 3)(x - 2)(x^2 + 1)} = \frac{2}{x + 3} - \frac{1}{x - 2} - \frac{1}{x^2 + 1}$$

Then we'll put the decomposition back into the integral in place of the original function.

$$\int \frac{x^3 - 8x^2 - 1}{(x^2 + x - 6)(x^2 + 1)} dx = \int \frac{2}{x + 3} - \frac{1}{x - 2} - \frac{1}{x^2 + 1} dx$$



$$2 \int \frac{1}{x+3} dx - \int \frac{1}{x-2} dx - \int \frac{1}{x^2+1} dx$$

$$2 \ln|x+3| - \ln|x-2| - \tan^{-1}x + C$$



**Topic:** Distinct quadratic factors

**Question:** Rewrite the integral using partial fractions. Do not solve it.

$$\int \frac{6x^3 + 2x^2 - x + 12}{(2x^2 - 1)(x^2 + 1)} dx$$

**Answer choices:**

A  $\frac{4}{3} \int \frac{x}{2x^2 - 1} dx + \frac{26}{3} \int \frac{1}{2x^2 - 1} dx + \frac{7}{3} \int \frac{x}{x^2 + 1} dx - \frac{10}{3} \int \frac{1}{x^2 + 1} dx$

B  $\frac{4}{3} \int \frac{x}{2x^2 - 1} dx + \frac{46}{3} \int \frac{1}{2x^2 - 1} dx - \frac{7}{3} \int \frac{x}{x^2 + 1} dx - \frac{10}{3} \int \frac{1}{x^2 + 1} dx$

C  $\frac{4}{3} \int \frac{x}{2x^2 + 1} dx + \frac{26}{3} \int \frac{1}{2x^2 + 1} dx + \frac{7}{3} \int \frac{x}{x^2 - 1} dx + \frac{10}{3} \int \frac{1}{x^2 - 1} dx$

D  $\frac{4}{3} \int \frac{x}{2x^2 - 1} dx + \frac{46}{3} \int \frac{1}{2x^2 - 1} dx + \frac{7}{3} \int \frac{x}{x^2 + 1} dx - \frac{10}{3} \int \frac{1}{x^2 + 1} dx$



**Solution: A**

The denominator is already factored as much as it can be, which means it's a product of irreducible factors.

$$\int \frac{6x^3 + 2x^2 - x + 12}{(2x^2 - 1)(x^2 + 1)} dx$$

Since the factors are quadratic, we know the numerators are going to be  $Ax + B$ ,  $Cx + D$ ,  $Ex + F$ , etc. For the partial fractions decomposition, we get

$$\frac{6x^3 + 2x^2 - x + 12}{(2x^2 - 1)(x^2 + 1)} = \frac{Ax + B}{2x^2 - 1} + \frac{Cx + D}{x^2 + 1}$$

Now we'll solve for constants.

$$\left[ \frac{6x^3 + 2x^2 - x + 12}{(2x^2 - 1)(x^2 + 1)} = \frac{Ax + B}{2x^2 - 1} + \frac{Cx + D}{x^2 + 1} \right] (2x^2 - 1)(x^2 + 1)$$

$$6x^3 + 2x^2 - x + 12 = (Ax + B)(x^2 + 1) + (Cx + D)(2x^2 - 1)$$

$$6x^3 + 2x^2 - x + 12 = Ax^3 + Ax + Bx^2 + B + 2Cx^3 - Cx + 2Dx^2 - D$$

$$6x^3 + 2x^2 - x + 12 = Ax^3 + 2Cx^3 + Bx^2 + 2Dx^2 + Ax - Cx + B - D$$

$$6x^3 + 2x^2 - x + 12 = (A + 2C)x^3 + (B + 2D)x^2 + (A - C)x + (B - D)$$

Equating coefficients on both sides, we get

$$\text{[1]} \quad A + 2C = 6$$

$$\text{[2]} \quad A - C = -1$$





$$\text{[3]} \quad B + 2D = 2$$

$$\text{[4]} \quad B - D = 12$$

We'll solve for  $A$  and  $C$  using [1] and [2], and for  $B$  and  $D$  using [3] and [4].  
Subtracting [2] from [1], we get

$$(A + 2C = 6) - (A - C = -1)$$

$$A - A + 2C + C = 6 + 1$$

$$3C = 7$$

$$C = \frac{7}{3}$$

Plugging the value for  $C$  back into [2] to solve for  $A$ , we get

$$A - \frac{7}{3} = -1$$

$$A = \frac{4}{3}$$

Subtracting [4] from [3], we get

$$(B + 2D = 2) - (B - D = 12)$$

$$B - B + 2D + D = 2 - 12$$

$$3D = -10$$

$$D = -\frac{10}{3}$$



Plugging the value for  $D$  back into [4] to solve for  $B$ , we get

$$B - \left(-\frac{10}{3}\right) = 12$$

$$B = \frac{26}{3}$$

Plugging the values for each of the four constants back into the partial fractions decomposition, and putting the decomposition back into the integral, we get

$$\int \frac{6x^3 + 2x^2 - x + 12}{(2x^2 - 1)(x^2 + 1)} dx = \int \frac{\frac{4}{3}x + \frac{26}{3}}{2x^2 - 1} + \frac{\frac{7}{3}x - \frac{10}{3}}{x^2 + 1} dx$$

$$\int \frac{\frac{4}{3}x}{2x^2 - 1} + \frac{\frac{26}{3}}{2x^2 - 1} + \frac{\frac{7}{3}x}{x^2 + 1} + \frac{-\frac{10}{3}}{x^2 + 1} dx$$

$$\frac{4}{3} \int \frac{x}{2x^2 - 1} dx + \frac{26}{3} \int \frac{1}{2x^2 - 1} dx + \frac{7}{3} \int \frac{x}{x^2 + 1} dx - \frac{10}{3} \int \frac{1}{x^2 + 1} dx$$



**Topic:** Distinct quadratic factors

**Question:** Rewrite the integral using partial fractions. Do not solve it.

$$\int \frac{x^3 - 2x^2 + 6x - 4}{(4x^2 - 2)(2x^2 + 6)} dx$$

**Answer choices:**

A  $-\frac{13}{14} \int \frac{x}{4x^2 - 2} dx - \frac{5}{7} \int \frac{1}{4x^2 - 2} dx - \frac{3}{14} \int \frac{x}{2x^2 + 6} dx - \frac{1}{7} \int \frac{1}{2x^2 + 6} dx$

B  $\frac{13}{14} \int \frac{x}{4x^2 - 2} dx - \frac{5}{7} \int \frac{1}{4x^2 - 2} dx + \frac{3}{14} \int \frac{x}{2x^2 + 6} dx + \frac{1}{7} \int \frac{1}{2x^2 + 6} dx$

C  $\frac{13}{14} \int \frac{x}{4x^2 - 2} dx + \frac{5}{7} \int \frac{1}{4x^2 - 2} dx - \frac{3}{14} \int \frac{x}{2x^2 + 6} dx - \frac{1}{7} \int \frac{1}{2x^2 + 6} dx$

D  $\frac{13}{14} \int \frac{x}{4x^2 - 2} dx - \frac{5}{7} \int \frac{1}{4x^2 - 2} dx - \frac{3}{14} \int \frac{x}{2x^2 + 6} dx - \frac{1}{7} \int \frac{1}{2x^2 + 6} dx$



**Solution: D**

The denominator is already factored as much as it can be, which means it's a product of irreducible factors.

$$\int \frac{x^3 - 2x^2 + 6x - 4}{(4x^2 - 2)(2x^2 + 6)} dx$$

Since the factors are quadratic, we know the numerators are going to be  $Ax + B$ ,  $Cx + D$ ,  $Ex + F$ , etc. For the partial fractions decomposition, we get

$$\frac{x^3 - 2x^2 + 6x - 4}{(4x^2 - 2)(2x^2 + 6)} = \frac{Ax + B}{4x^2 - 2} + \frac{Cx + D}{2x^2 + 6}$$

Now we'll solve for constants.

$$\left[ \frac{x^3 - 2x^2 + 6x - 4}{(4x^2 - 2)(2x^2 + 6)} = \frac{Ax + B}{4x^2 - 2} + \frac{Cx + D}{2x^2 + 6} \right] (4x^2 - 2)(2x^2 + 6)$$

$$x^3 - 2x^2 + 6x - 4 = (Ax + B)(2x^2 + 6) + (Cx + D)(4x^2 - 2)$$

$$x^3 - 2x^2 + 6x - 4 = 2Ax^3 + 6Ax + 2Bx^2 + 6B + 4Cx^3 - 2Cx + 4Dx^2 - 2D$$

$$x^3 - 2x^2 + 6x - 4 = 2Ax^3 + 4Cx^3 + 2Bx^2 + 4Dx^2 + 6Ax - 2Cx + 6B - 2D$$

$$x^3 - 2x^2 + 6x - 4 = (2A + 4C)x^3 + (2B + 4D)x^2 + (6A - 2C)x + (6B - 2D)$$

Equating coefficients on both sides, we get

$$\text{[1]} \quad 2A + 4C = 1$$

$$\text{[2]} \quad 6A - 2C = 6$$



$$[3] \quad 2B + 4D = -2$$

$$[4] \quad 6B - 2D = -4$$

We'll solve for  $A$  and  $C$  using [1] and [2], and for  $B$  and  $D$  using [3] and [4]. Multiplying [1] by 3 so that both [1] and [2] contain  $6A$ , and then subtracting [2] from [1], we get

$$6A + 12C - (6A - 2C) = 3 - 6$$

$$6A + 12C - 6A + 2C = 3 - 6$$

$$14C = -3$$

$$C = -\frac{3}{14}$$

Plugging the value for  $C$  back into [2] to solve for  $A$ , we get

$$6A - 2\left(-\frac{3}{14}\right) = 6$$

$$A = \frac{13}{14}$$

Multiplying [3] by 3 so that both [3] and [4] contain  $6B$ , and then subtracting [4] from [3], we get

$$6B + 12D - (6B - 2D) = -6 - (-4)$$

$$6B + 12D - 6B + 2D = -6 - (-4)$$

$$14D = -2$$



$$D = -\frac{2}{14}$$

Plugging the value for  $D$  back into [4] to solve for  $B$ , we get

$$6B - 2\left(-\frac{1}{7}\right) = -4$$

$$B = -\frac{5}{7}$$

Plugging the values for each of the four constants back into the partial fractions decomposition, and putting the decomposition back into the integral, we get

$$\int \frac{x^3 - 2x^2 + 6x - 4}{(4x^2 - 2)(2x^2 + 6)} dx = \int \frac{\frac{13}{14}x - \frac{5}{7}}{4x^2 - 2} + \frac{-\frac{3}{14}x - \frac{1}{7}}{2x^2 + 6} dx$$

$$\int \frac{\frac{13}{14}x - \frac{5}{7}}{4x^2 - 2} - \frac{\frac{3}{14}x + \frac{1}{7}}{2x^2 + 6} dx$$

$$\int \frac{\frac{13}{14}x}{4x^2 - 2} + \frac{-\frac{5}{7}}{4x^2 - 2} - \frac{\frac{3}{14}x}{2x^2 + 6} - \frac{\frac{1}{7}}{2x^2 + 6} dx$$

$$\frac{13}{14} \int \frac{x}{4x^2 - 2} dx - \frac{5}{7} \int \frac{1}{4x^2 - 2} dx - \frac{3}{14} \int \frac{x}{2x^2 + 6} dx - \frac{1}{7} \int \frac{1}{2x^2 + 6} dx$$

