

**Topic:** Inverse hyperbolic derivatives**Question:** Find the derivative of the inverse hyperbolic function.

$$y = \cosh^{-1}(x^3)$$

**Answer choices:**

A  $y' = \frac{3x^2}{\sqrt{x^6 - 1}}$  with  $x^3 < 1$

B  $y' = \frac{3x^2}{\sqrt{x^6 + 1}}$  with  $x^3 < 1$

C  $y' = \frac{3x^2}{\sqrt{x^6 + 1}}$  with  $x^3 > 1$

D  $y' = \frac{3x^2}{\sqrt{x^6 - 1}}$  with  $x^3 > 1$



**Solution: D**

Apply the formula for the derivative of inverse hyperbolic cosine, with  $g(x) = x^3$  and  $g'(x) = 3x^2$ .

$$y' = \frac{g'(x)}{\sqrt{[g(x)]^2 - 1}} \quad \text{with } x^3 > 1$$

$$y' = \frac{3x^2}{\sqrt{(x^3)^2 - 1}} \quad \text{with } x^3 > 1$$

$$y' = \frac{3x^2}{\sqrt{x^6 - 1}} \quad \text{with } x^3 > 1$$



**Topic:** Inverse hyperbolic derivatives**Question:** Find the derivative of the inverse hyperbolic function.

$$y = \tanh^{-1}(2x^5 - 1)$$

**Answer choices:**

A  $y' = \frac{5}{2x + 2x^6}$  with  $|2x^5 - 1| < 1$

B  $y' = \frac{1}{2x - 2x^6}$  with  $|2x^5 - 1| < 1$

C  $y' = \frac{5}{x - x^6}$  with  $|2x^5 - 1| < 1$

D  $y' = \frac{5}{2x - 2x^6}$  with  $|2x^5 - 1| < 1$



**Solution: D**

Apply the formula for the derivative of inverse hyperbolic tangent, with  $g(x) = 2x^5 - 1$  and  $g'(x) = 10x^4$ .

$$y' = \frac{g'(x)}{1 - [g(x)]^2} \quad \text{with } |g(x)| < 1$$

$$y' = \frac{10x^4}{1 - (2x^5 - 1)^2} \quad \text{with } |2x^5 - 1| < 1$$

Simplify the derivative.

$$y' = \frac{10x^4}{1 - (4x^{10} - 4x^5 + 1)} \quad \text{with } |2x^5 - 1| < 1$$

$$y' = \frac{10x^4}{1 - 4x^{10} + 4x^5 - 1} \quad \text{with } |2x^5 - 1| < 1$$

$$y' = \frac{10x^4}{-4x^{10} + 4x^5} \quad \text{with } |2x^5 - 1| < 1$$

$$y' = \frac{5}{-2x^6 + 2x} \quad \text{with } |2x^5 - 1| < 1$$

$$y' = \frac{5}{2x - 2x^6} \quad \text{with } |2x^5 - 1| < 1$$



**Topic:** Inverse hyperbolic derivatives**Question:** Find the derivative of the inverse hyperbolic function.

$$f(x) = (\sinh^{-1}(2x^3))^4$$

**Answer choices:**

A  $f'(x) = \frac{24x^2(\sinh^{-1}(2x^3))^3}{\sqrt{x^2 + 1}}$

B  $f'(x) = \frac{24x^2(\sinh^{-1}(2x^3))^3}{\sqrt{4x^6 + 1}}$

C  $f'(x) = \frac{6x^2(\sinh^{-1}(2x^3))^3}{\sqrt{x^2 + 1}}$

D  $f'(x) = \frac{6x^2(\sinh^{-1}(2x^3))^3}{\sqrt{4x^6 + 1}}$



**Solution: B**

Use a substitution with  $u = \sinh^{-1}(2x^3)$  and

$$u' = \frac{6x^2}{\sqrt{(2x^3)^2 + 1}}$$

$$u' = \frac{6x^2}{\sqrt{4x^6 + 1}}$$

Then the function is

$$f(x) = u^4$$

and the derivative is

$$f'(x) = 4u^3 \cdot u'$$

$$f'(x) = 4(\sinh^{-1}(2x^3))^3 \cdot \frac{6x^2}{\sqrt{4x^6 + 1}}$$

$$f'(x) = \frac{24x^2(\sinh^{-1}(2x^3))^3}{\sqrt{4x^6 + 1}}$$

