

Arc length of a polar curve

The arc length of a polar curve is simply the length of a section of a polar parametric curve between two points a and b . We use the formula

$$L = \int_a^b \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

where L is the arc length

r is the equation of the polar curve

$\frac{dr}{d\theta}$ is the derivative of the polar curve

a and b are the endpoints of the section

Example

Find the arc length of the polar curve over the given interval.

$$r = \cos^2 \frac{\theta}{2}$$

$$0 \leq \theta \leq \frac{\pi}{2}$$

Before we can plug into the arc length formula, we need to find $dr/d\theta$.

$$\frac{dr}{d\theta} = 2 \cos \frac{\theta}{2} \left[-\sin \frac{\theta}{2} \right] \left(\frac{1}{2} \right)$$



$$\frac{dr}{d\theta} = -\cos \frac{\theta}{2} \sin \frac{\theta}{2}$$

Now we can go ahead and solve for the arc length

$$L = \int_0^{\frac{\pi}{2}} \sqrt{\left(\cos^2 \frac{\theta}{2}\right)^2 + \left(-\cos \frac{\theta}{2} \sin \frac{\theta}{2}\right)^2} d\theta$$

$$L = \int_0^{\frac{\pi}{2}} \sqrt{\cos^4 \frac{\theta}{2} + \cos^2 \frac{\theta}{2} \sin^2 \frac{\theta}{2}} d\theta$$

$$L = \int_0^{\frac{\pi}{2}} \sqrt{\cos^2 \frac{\theta}{2} \left(\cos^2 \frac{\theta}{2} + \sin^2 \frac{\theta}{2}\right)} d\theta$$

Since $\cos^2 x + \sin^2 x = 1$, we get

$$L = \int_0^{\frac{\pi}{2}} \sqrt{\left(\cos^2 \frac{\theta}{2}\right)(1)} d\theta$$

$$L = \int_0^{\frac{\pi}{2}} \cos \frac{\theta}{2} d\theta$$

$$L = 2 \sin \frac{\theta}{2} \Big|_0^{\frac{\pi}{2}}$$

$$L = 2 \sin \left(\frac{\frac{\pi}{2}}{2}\right) - 2 \sin \left(\frac{0}{2}\right)$$

$$L = 2 \sin \frac{\pi}{4} - 2 \sin 0$$



$$L = 2 \cdot \frac{\sqrt{2}}{2} - 2(0)$$

$$L = \sqrt{2}$$

Let's do another example.

Example

Find the arc length of the polar curve over the given interval.

$$r = e^{2\theta}$$

$$0 \leq \theta \leq \pi$$

Before we can plug into the arc length formula, we need to find $dr/d\theta$.

$$\frac{dr}{d\theta} = 2e^{2\theta}$$

Plugging everything into the formula, we get

$$s = \int_0^{\pi} \sqrt{(e^{2\theta})^2 + (2e^{2\theta})^2} d\theta$$

$$s = \int_0^{\pi} \sqrt{e^{4\theta} + 4e^{4\theta}} d\theta$$



$$s = \int_0^{\pi} \sqrt{5e^{4\theta}} d\theta$$

$$s = \sqrt{5} \int_0^{\pi} e^{2\theta} d\theta$$

$$s = \frac{\sqrt{5}}{2} e^{2\theta} \Big|_0^{\pi}$$

$$s = \frac{\sqrt{5}}{2} e^{2(\pi)} - \frac{\sqrt{5}}{2} e^{2(0)}$$

$$s = \frac{\sqrt{5}}{2} e^{2\pi} - \frac{\sqrt{5}}{2}$$

$$s = \frac{\sqrt{5} (e^{2\pi} - 1)}{2}$$

