



Calculus 2 Workbook Solutions

Calculus with parametric curves

krista king
MATH

TANGENT LINES OF PARAMETRIC CURVES

- 1. Find the equation of the tangent line to the parametric curve at $t = 3$.

$$x = 3t + 5$$

$$y = 7t - 2$$

Solution:

Use the formula of the tangent line as $y - y_1 = m(x - x_1)$ and transform the equation into the form $y = mx + b$.

At $t = 3$, the slope of the parametric equation is

$$m = \frac{dy}{dx} = \frac{\frac{d}{dt}(7t - 2)}{\frac{d}{dt}(3t + 5)} = \frac{7}{3}$$

At $t = 3$, the parametric equation has the values

$$x(3) = 3(3) + 5 = 14$$

$$y(3) = 7(3) - 2 = 19$$

Putting these values together gives the coordinate point $(x, y) = (14, 19)$.

Plug everything into the equation of the tangent line.

$$y - y_1 = m(x - x_1)$$



$$y - 19 = \frac{7}{3}(x - 14)$$

$$y = \frac{7}{3}x - \frac{98}{3} + \frac{57}{3}$$

$$y = \frac{7}{3}x - \frac{41}{3}$$

■ 2. Find the equation of the tangent line to the parametric curve at $t = 4$.

$$x = 3t^2 - 12$$

$$y = 2t^3 + 6$$

Solution:

Use the formula of the tangent line as $y - y_1 = m(x - x_1)$ and transform the equation into the form $y = mx + b$.

At $t = 4$, the slope of the parametric equation is

$$m = \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{\frac{d}{dt}(2t^3 + 6)}{\frac{d}{dt}(3t^2 - 12)} = \frac{6t^2}{6t} = t = 4$$

At $t = 4$, the parametric equation has the values

$$x(4) = 3(4)^2 - 12 = 36$$

$$y(4) = 2(4)^3 + 6 = 134$$



Putting these values together gives the coordinate point $(x, y) = (36, 134)$.
 Plug everything into the equation of the tangent line.

$$y - y_1 = m(x - x_1)$$

$$y - 134 = 4(x - 36)$$

$$y = 4x - 144 + 134$$

$$y = 4x - 10$$

■ 3. Find the equation of the tangent line to the parametric curve at $t = \pi/3$.

$$x = \cos^2 t$$

$$y = \sin^2 t$$

Solution:

Use the formula of the tangent line as $y - y_1 = m(x - x_1)$ and transform the equation into the form $y = mx + b$.

At $t = \pi/3$, the slope of the parametric equation is

$$m = \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{\frac{d}{dt}(\sin^2 t)}{\frac{d}{dt}(\cos^2 t)} = \frac{2 \sin t \cos t}{-2 \cos t \sin t} = -1$$

At $t = \pi/3$, the parametric equation has the values



$$x\left(\frac{\pi}{3}\right) = \cos^2\left(\frac{\pi}{3}\right) = \left(\frac{1}{2}\right)^2 = \frac{1}{4}$$

$$y\left(\frac{\pi}{3}\right) = \sin^2\left(\frac{\pi}{3}\right) = \left(\frac{\sqrt{3}}{2}\right)^2 = \frac{3}{4}$$

Putting these values together gives the coordinate point $(x, y) = (1/4, 3/4)$.

Plug everything into the equation of the tangent line.

$$y - y_1 = m(x - x_1)$$

$$y - \frac{3}{4} = -1\left(x - \frac{1}{4}\right)$$

$$y = -x + \frac{1}{4} + \frac{3}{4}$$

$$y = -x + 1$$

■ 4. Find the equation of the tangent line to the parametric curve at $t = 4$.

$$x = t^2 + t + 3$$

$$y = t^2 - 3t + 2$$

Solution:

Use the formula of the tangent line as $y - y_1 = m(x - x_1)$ and transform the equation into the form $y = mx + b$.



At $t = 4$, the slope of the parametric equation is

$$m = \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{\frac{d}{dt}(t^2 - 3t + 2)}{\frac{d}{dt}(t^2 + t + 3)} = \frac{2t - 3}{2t + 1} = \frac{2(4) - 3}{2(4) + 1} = \frac{5}{9}$$

At $t = 4$, the parametric equation has the values

$$x(4) = 4^2 + 4 + 3 = 23$$

$$y(4) = 4^2 - 3(4) + 2 = 6$$

Putting these values together gives the coordinate point $(x, y) = (23, 6)$. Plug everything into the equation of the tangent line.

$$y - y_1 = m(x - x_1)$$

$$y - 6 = \frac{5}{9}(x - 23)$$

$$y = \frac{5}{9}x - \frac{115}{9} + \frac{54}{9}$$

$$y = \frac{5}{9}x - \frac{61}{9}$$

■ 5. Find the equation of the tangent line to the parametric curve at $t = 9$.

$$x = 3\sqrt{t}$$

$$y = 5t\sqrt{t}$$



Solution:

Use the formula of the tangent line as $y - y_1 = m(x - x_1)$ and transform the equation into the form $y = mx + b$.

At $t = 9$, the slope of the parametric equation is

$$m = \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{\frac{d}{dt}(5t\sqrt{t})}{\frac{d}{dt}(3\sqrt{t})} = \frac{\frac{15}{2}\sqrt{t}}{\frac{3}{2\sqrt{t}}} = \frac{15}{2}\sqrt{t} \cdot \frac{2\sqrt{t}}{3} = 5t = 5(9) = 45$$

At $t = 9$, the parametric equation has the values

$$x(9) = 3\sqrt{9} = 3 \cdot 3 = 9$$

$$y(9) = 5 \cdot 9\sqrt{9} = 45 \cdot 3 = 135$$

Putting these values together gives the coordinate point $(x, y) = (9, 135)$. Plug everything into the equation of the tangent line.

$$y - y_1 = m(x - x_1)$$

$$y - 135 = 45(x - 9)$$

$$y = 45x - 405 + 135$$

$$y = 45x - 270$$



AREA UNDER A PARAMETRIC CURVE

- 1. Find the area under the parametric curve.

$$x(t) = 3t^2$$

$$y(t) = t + 2$$

$$0 \leq t \leq 3$$

Solution:

Find the area under the curve on $a \leq t \leq b$ using the integral formula.

$$A = \int_a^b y(t)x'(t) dt$$

$$A = \int_0^3 (t+2)(6t) dt$$

$$A = \int_0^3 6t^2 + 12t dt$$

Integrate and evaluate over the interval.

$$A = \frac{6t^3}{3} + \frac{12t^2}{2} \Big|_0^3$$



$$A = 2t^3 + 6t^2 \Big|_0^3$$

$$A = \left(2(3)^3 + 6(3)^2\right) - \left(2(0)^3 + 6(0)^2\right)$$

$$A = 54 + 54$$

$$A = 108$$

■ 2. Find the area under the parametric curve.

$$x(t) = 5t^2 - 3t + 4$$

$$y(t) = 6t - 1$$

$$0 \leq t \leq 5$$

Solution:

Find the area under the curve on $a \leq t \leq b$ using the integral formula.

$$A = \int_a^b y(t)x'(t) \, dt$$

$$A = \int_0^5 (6t - 1)(10t - 3) \, dt$$

$$A = \int_0^5 60t^2 - 28t + 3 \, dt$$



Integrate and evaluate over the interval.

$$A = \frac{60t^3}{3} - \frac{28t^2}{2} + 3t \Big|_0^5$$

$$A = 20t^3 - 14t^2 + 3t \Big|_0^5$$

$$A = \left(20(5)^3 - 14(5)^2 + 3(5)\right) - \left(20(0)^3 - 14(0)^2 + 3(0)\right)$$

$$A = 2,500 - 350 + 15$$

$$A = 2,165$$

■ 3. Find the area under the parametric curve.

$$x(t) = t + \sin t$$

$$y(t) = 4 + \cos t$$

$$0 \leq t \leq 2\pi$$

Solution:

Find the area under the curve on $a \leq t \leq b$ using the integral formula.

$$A = \int_a^b y(t)x'(t) \, dt$$



$$A = \int_0^{2\pi} (4 + \cos t)(1 + \cos t) dt$$

$$A = \int_0^{2\pi} 4 + 5 \cos t + \cos^2 t dt$$

Use the trig identity to make a substitution.

$$\cos^2 t = \frac{1}{2} + \frac{1}{2} \cos(2t)$$

Substitute.

$$A = \int_0^{2\pi} 4 + 5 \cos t + \frac{1}{2} + \frac{1}{2} \cos(2t) dt$$

$$A = \int_0^{2\pi} \frac{9}{2} + 5 \cos t + \frac{1}{2} \cos(2t) dt$$

Integrate and evaluate over the interval.

$$A = \frac{9}{2}t + 5 \sin t + \frac{1}{4} \sin(2t) \Big|_0^{2\pi}$$

$$A = \frac{9}{2}(2\pi) + 5 \sin(2\pi) + \frac{1}{4} \sin(2(2\pi)) - \left(\frac{9}{2}(0) + 5 \sin(0) + \frac{1}{4} \sin(2(0)) \right)$$

$$A = 9\pi + 5(0) + \frac{1}{4}(0)$$

$$A = 9\pi$$



- 4. Find the area under the parametric curve.

$$x(t) = t^2 + 5t - 8$$

$$y(t) = t^2 + 4t + 2$$

$$0 \leq t \leq 2$$

Solution:

Find the area under the curve on $a \leq t \leq b$ using the integral formula.

$$A = \int_a^b y(t)x'(t) dt$$

$$A = \int_0^2 (t^2 + 4t + 2)(2t + 5) dt$$

$$A = \int_0^2 2t^3 + 13t^2 + 24t + 10 dt$$

Integrate and evaluate over the interval.

$$A = \left. \frac{2t^4}{4} + \frac{13t^3}{3} + \frac{24t^2}{2} - 10t \right|_0^2$$

$$A = \left. \frac{t^4}{2} + \frac{13t^3}{3} + 12t^2 + 10t \right|_0^2$$

$$A = \left(\frac{2^4}{2} + \frac{13(2)^3}{3} + 12(2)^2 + 10(2) \right) - \left(\frac{0^4}{2} + \frac{13(0)^3}{3} + 12(0)^2 + 10(0) \right)$$



$$A = 8 + \frac{104}{3} + 48 + 20$$

$$A = \frac{104}{3} + 76$$

$$A = \frac{104}{3} + \frac{228}{3}$$

$$A = \frac{332}{3}$$



AREA UNDER ONE ARC OR LOOP

- 1. Find the area in one loop of the parametric curve.

$$x(\theta) = 2 \cos(2\theta)$$

$$y(\theta) = 4 + \sin(2\theta)$$

$$0 \leq \theta \leq \pi$$

Solution:

Plug the parametric equation and the given interval into the integral formula for the area under one arc or in one loop of the parametric curve.

$$A = \int_a^b y(\theta) \cdot x'(\theta) \, d\theta$$

$$A = \int_0^\pi (4 + \sin(2\theta))(-4 \sin(2\theta)) \, d\theta$$

$$A = \int_0^\pi -16 \sin(2\theta) - 4 \sin^2(2\theta) \, d\theta$$

Use the reduction formula

$$\sin^2(2\theta) = \frac{1}{2}(1 - \cos(4\theta)) = \frac{1}{2} - \frac{1}{2} \cos(4\theta)$$

Substitute that into the integral.



$$A = \int_0^{\pi} -16 \sin(2\theta) - 4 \left(\frac{1}{2} - \frac{1}{2} \cos(4\theta) \right) d\theta$$

$$A = \int_0^{\pi} -16 \sin(2\theta) - 2 + 2 \cos(4\theta) d\theta$$

Integrate, then evaluate over the interval.

$$A = -16 \left(-\frac{\cos(2\theta)}{2} \right) - 2\theta + 2 \left(\frac{\sin(4\theta)}{4} \right) \Big|_0^{\pi}$$

$$A = 8 \cos(2\theta) - 2\theta + \frac{1}{2} \sin(4\theta) \Big|_0^{\pi}$$

$$A = 8 \cos(2\pi) - 2\pi + \frac{1}{2} \sin(4\pi) - \left(8 \cos(2(0)) - 2(0) + \frac{1}{2} \sin(4(0)) \right)$$

$$A = 8(1) - 2\pi + \frac{1}{2}(0) - 8(1) + 0 - \frac{1}{2}(0)$$

$$A = 8 - 2\pi - 8$$

$$A = -2\pi$$

$$A = |-2\pi|$$

$$A = 2\pi$$

■ 2. Find the area in one loop of the parametric curve.

$$x(\theta) = 2 \sin \theta$$



$$y(\theta) = 5 + \cos \theta$$

$$0 \leq \theta \leq 2\pi$$

Solution:

Plug the parametric equation and the given interval into the integral formula for the area under one arc or in one loop of the parametric curve.

$$A = \int_a^b y(\theta) \cdot x'(\theta) \, d\theta$$

$$A = \int_0^{2\pi} (5 + \cos \theta)(2 \cos \theta) \, d\theta$$

$$A = \int_0^{2\pi} 10 \cos \theta + 2 \cos^2 \theta \, d\theta$$

Use the reduction formula

$$\cos^2 \theta = \frac{1}{2}(1 + \cos(2\theta)) = \frac{1}{2} + \frac{1}{2} \cos(2\theta)$$

Substitute that into the integral.

$$A = \int_0^{2\pi} 10 \cos \theta + 2 \left(\frac{1}{2} + \frac{1}{2} \cos(2\theta) \right) \, d\theta$$

$$A = \int_0^{2\pi} 10 \cos \theta + 1 + \cos(2\theta) \, d\theta$$



Integrate, then evaluate over the interval.

$$A = 10 \sin \theta + \theta + \frac{\sin(2\theta)}{2} \Big|_0^{2\pi}$$

$$A = 10 \sin(2\pi) + 2\pi + \frac{\sin(2(2\pi))}{2} - \left(10 \sin(0) + 0 + \frac{\sin(2(0))}{2} \right)$$

$$A = 10(0) + 2\pi + \frac{0}{2} - 10(0) - 0 - \frac{0}{2}$$

$$A = 2\pi$$

■ 3. Find the area in one loop of the parametric curve.

$$x(\theta) = 8 + 3 \cos \theta$$

$$y(\theta) = 9 - 2 \sin \theta$$

$$0 \leq \theta \leq 2\pi$$

Solution:

Plug the parametric equation and the given interval into the integral formula for the area under one arc or in one loop of the parametric curve.

$$A = \int_a^b y(\theta) \cdot x'(\theta) \, d\theta$$



$$A = \int_0^{2\pi} (9 - 2 \sin \theta)(-3 \sin \theta) d\theta$$

$$A = \int_0^{2\pi} -27 \sin \theta + 6 \sin^2 \theta d\theta$$

Use the reduction formula

$$\sin^2(2\theta) = \frac{1}{2}(1 - \cos(4\theta)) = \frac{1}{2} - \frac{1}{2} \cos(4\theta)$$

Substitute that into the integral.

$$A = \int_0^{2\pi} -27 \sin \theta + 6 \left(\frac{1}{2} - \frac{1}{2} \cos(2\theta) \right) d\theta$$

$$A = \int_0^{2\pi} -27 \sin \theta + 3 - 3 \cos(2\theta) d\theta$$

Integrate, then evaluate over the interval.

$$A = 27 \cos \theta + 3\theta - \frac{3}{2} \sin(2\theta) \Big|_0^{2\pi}$$

$$A = 27 \cos(2\pi) + 3(2\pi) - \frac{3}{2} \sin(2(2\pi)) - \left(27 \cos(0) + 3(0) - \frac{3}{2} \sin(2(0)) \right)$$

$$A = 27(1) + 6\pi - \frac{3}{2}(0) - 27(1) - 0 + \frac{3}{2}(0)$$

$$A = 27 + 6\pi - 27$$

$$A = 6\pi$$



- 4. Find the area in one loop of the parametric curve.

$$x(\theta) = 12 + 6 \sin \theta$$

$$y(\theta) = 12 - 6 \cos \theta$$

$$0 \leq \theta \leq 2\pi$$

Solution:

Plug the parametric equation and the given interval into the integral formula for the area under one arc or in one loop of the parametric curve.

$$A = \int_a^b y(\theta) \cdot x'(\theta) \, d\theta$$

$$A = \int_0^{2\pi} (12 - 6 \cos \theta)(6 \cos \theta) \, d\theta$$

$$A = \int_0^{2\pi} 72 \cos \theta - 36 \cos^2 \theta \, d\theta$$

Use the reduction formula

$$\cos^2 \theta = \frac{1}{2}(1 + \cos(2\theta)) = \frac{1}{2} + \frac{1}{2} \cos(2\theta)$$

Substitute that into the integral.



$$A = \int_0^{2\pi} 72 \cos \theta - 36 \left(\frac{1}{2} + \frac{1}{2} \cos(2\theta) \right) d\theta$$

$$A = \int_0^{2\pi} 72 \cos \theta - 18 - 18 \cos(2\theta) d\theta$$

Integrate, then evaluate over the interval.

$$A = 72 \sin \theta - 18\theta - 9 \sin(2\theta) \Big|_0^{2\pi}$$

$$A = 72 \sin(2\pi) - 18(2\pi) - 9 \sin(2(2\pi)) - (72 \sin(0) - 18(0) - 9 \sin(2(0)))$$

$$A = 72(0) - 36\pi - 9(0) - 72(0) + 0 + 9(0)$$

$$A = -36\pi$$

$$A = |-36\pi|$$

$$A = 36\pi$$

■ 5. Find the area in one loop of the parametric curve.

$$x(\theta) = 15 - 5 \cos \theta$$

$$y(\theta) = 5 + 15 \sin \theta$$

$$0 \leq \theta \leq 2\pi$$



Solution:

Plug the parametric equation and the given interval into the integral formula for the area under one arc or in one loop of the parametric curve.

$$A = \int_a^b y(\theta) \cdot x'(\theta) \, d\theta$$

$$A = \int_0^{2\pi} (5 + 15 \sin \theta)(5 \sin \theta) \, d\theta$$

$$A = \int_0^{2\pi} 25 \sin \theta + 75 \sin^2 \theta \, d\theta$$

Use the reduction formula

$$\sin^2(2\theta) = \frac{1}{2}(1 - \cos(4\theta)) = \frac{1}{2} - \frac{1}{2} \cos(4\theta)$$

Substitute that into the integral.

$$A = \int_0^{2\pi} 25 \sin \theta + 75 \left(\frac{1}{2} - \frac{1}{2} \cos(2\theta) \right) \, d\theta$$

$$A = \int_0^{2\pi} 25 \sin \theta + \frac{75}{2} - \frac{75}{2} \cos(2\theta) \, d\theta$$

Integrate, then evaluate over the interval.

$$A = -25 \cos \theta + \frac{75}{2} \theta - \frac{75}{4} \sin(2\theta) \Big|_0^{2\pi}$$



$$A = -25 \cos(2\pi) + \frac{75}{2}(2\pi) - \frac{75}{4} \sin(2(2\pi)) - \left(-25 \cos(0) + \frac{75}{2}(0) - \frac{75}{4} \sin(2(0)) \right)$$

$$A = -25(1) + 75\pi - \frac{75}{4}(0) + 25(1) - 0 + \frac{75}{4}(0)$$

$$A = -25 + 75\pi + 25$$

$$A = 75\pi$$



ARC LENGTH OF PARAMETRIC CURVES

- 1. Find the length of the parametric curve on the given interval.

$$x(t) = 7 - 3t$$

$$y(t) = 5 + 8t$$

$$-1 \leq t \leq 4$$

Solution:

Plug the derivatives of $x(t)$ and $y(t)$ and the given interval into the integral formula for arc length.

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$L = \int_{-1}^4 \sqrt{(-3)^2 + (8)^2} dt$$

$$L = \int_{-1}^4 \sqrt{9 + 64} dt$$

$$L = \int_{-1}^4 \sqrt{73} dt$$

Integrate, then evaluate over the interval.



$$L = \sqrt{73}t \Big|_{-1}^4$$

$$L = \sqrt{73}(4) - \sqrt{73}(-1)$$

$$L = 4\sqrt{73} + \sqrt{73}$$

$$L = 5\sqrt{73}$$

■ 2. Find the length of the parametric curve on the given interval.

$$x(t) = \cos^3 t$$

$$y(t) = \sin^3 t$$

$$0 \leq t \leq \frac{3\pi}{4}$$

Solution:

Plug the derivatives of $x(t)$ and $y(t)$ and the given interval into the integral formula for arc length.

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$L = \int_0^{\frac{3\pi}{4}} \sqrt{(3 \cos^2 t (-\sin t))^2 + (3 \sin^2 t \cos t)^2} dt$$



$$L = \int_0^{\frac{3\pi}{4}} \sqrt{9 \cos^4 t \sin^2 t + 9 \sin^4 t \cos^2 t} \, dt$$

$$L = \int_0^{\frac{3\pi}{4}} \sqrt{9 \sin^2 t \cos^2 t (\cos^2 t + \sin^2 t)} \, dt$$

$$L = \int_0^{\frac{3\pi}{4}} \sqrt{9 \sin^2 t \cos^2 t (1)} \, dt$$

$$L = \int_0^{\frac{3\pi}{4}} \sqrt{9 \sin^2 t \cos^2 t} \, dt$$

$$L = \int_0^{\frac{3\pi}{4}} 3 \sin t \cos t \, dt$$

Use u-substitution.

$$u = \sin t$$

$$du = \cos t \, dt, \text{ so } dt = \frac{du}{\cos t}$$

Substitute.

$$L = \int_{t=0}^{t=\frac{3\pi}{4}} 3u \cos t \left(\frac{du}{\cos t} \right)$$

$$L = \int_{t=0}^{t=\frac{3\pi}{4}} 3u \, du$$

Integrate, back-substitute, then evaluate over the interval.



$$L = \frac{3}{2} u^2 \Big|_{t=0}^{t=\frac{3\pi}{4}}$$

$$L = \frac{3}{2} \sin^2 t \Big|_0^{\frac{3\pi}{4}}$$

$$L = \frac{3}{2} \sin^2 \frac{3\pi}{4} - \frac{3}{2} \sin^2(0)$$

$$L = \frac{3}{2} \left(\frac{\sqrt{2}}{2} \right)^2 - \frac{3}{2} (0)^2$$

$$L = \frac{3}{2} \left(\frac{2}{4} \right)$$

$$L = \frac{3}{4}$$

■ 3. Find the length of the parametric curve on the given interval.

$$x(t) = 5t - 5 \sin t$$

$$y(t) = -5 \cos t$$

$$0 \leq t \leq 2\pi$$

Solution:



Plug the derivatives of $x(t)$ and $y(t)$ and the given interval into the integral formula for arc length.

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$L = \int_0^{2\pi} \sqrt{(5 - 5 \cos t)^2 + (5 \sin t)^2} dt$$

$$L = \int_0^{2\pi} \sqrt{25 - 50 \cos t + 25 \cos^2 t + 25 \sin^2 t} dt$$

$$L = \int_0^{2\pi} \sqrt{25 - 50 \cos t + 25(\cos^2 t + \sin^2 t)} dt$$

$$L = \int_0^{2\pi} \sqrt{25 - 50 \cos t + 25(1)} dt$$

$$L = \int_0^{2\pi} \sqrt{50 - 50 \cos t} dt$$

$$L = \int_0^{2\pi} \sqrt{50(1 - \cos t)} dt$$

$$L = \int_0^{2\pi} \sqrt{100 \cdot \frac{1}{2}(1 - \cos t)} dt$$

Use the reduction formula

$$\sin^2 \theta = \frac{1}{2}(1 - \cos(2\theta)) = \frac{1}{2} - \frac{1}{2} \cos(2\theta)$$



Substitute.

$$L = \int_0^{2\pi} \sqrt{100 \sin^2 \frac{t}{2}} dt$$

$$L = \int_0^{2\pi} 10 \sin \frac{t}{2} dt$$

Integrate, then evaluate over the interval.

$$L = -20 \cos \frac{t}{2} \Big|_0^{2\pi}$$

$$L = -20 \cos \frac{2\pi}{2} - \left(-20 \cos \frac{0}{2} \right)$$

$$L = -20 \cos \pi - (-20 \cos 0)$$

$$L = -20(-1) + 20(1)$$

$$L = 20 + 20$$

$$L = 40$$

■ 4. Find the length of the parametric curve on the given interval.

$$x(t) = \cos t$$

$$y(t) = t + \sin t$$

$$0 \leq t \leq \pi$$



Solution:

Plug the derivatives of $x(t)$ and $y(t)$ and the given interval into the integral formula for arc length.

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$L = \int_0^\pi \sqrt{(-\sin t)^2 + (1 + \cos t)^2} dt$$

$$L = \int_0^\pi \sqrt{\sin^2 t + 1 + 2 \cos t + \cos^2 t} dt$$

$$L = \int_0^\pi \sqrt{1 + 2 \cos t + (\sin^2 t + \cos^2 t)} dt$$

$$L = \int_0^\pi \sqrt{1 + 2 \cos t + (1)} dt$$

$$L = \int_0^\pi \sqrt{2 + 2 \cos t} dt$$

$$L = \int_0^\pi \sqrt{2(1 + \cos t)} dt$$

$$L = \int_0^\pi \sqrt{4 \cdot \frac{1}{2}(1 + \cos t)} dt$$

Use the reduction formula



$$\cos^2 \theta = \frac{1}{2}(1 + \cos(2\theta)) = \frac{1}{2} + \frac{1}{2} \cos(2\theta)$$

Substitute.

$$L = \int_0^{\pi} \sqrt{4 \cos^2 \frac{t}{2}} dt$$

$$L = \int_0^{\pi} 2 \cos \frac{t}{2} dt$$

Integrate, then evaluate over the interval.

$$L = 4 \sin \frac{t}{2} \Big|_0^{\pi}$$

$$L = 4 \sin \frac{\pi}{2} - 4 \sin \frac{0}{2}$$

$$L = 4(1) - 4(0)$$

$$L = 4$$



SURFACE AREA OF REVOLUTION, HORIZONTAL AXIS

- 1. Find the surface area of revolution of the parametric curve on the interval $0 \leq t \leq 3$, rotated about the x -axis.

$$x = \frac{5}{3}t$$

$$y = 4t + 6$$

Solution:

Plug the derivatives and the interval into the integral formula formula for the surface area of revolution for a parametric curve about the x -axis.

$$S = \int_a^b 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$S = \int_0^3 2\pi(4t + 6) \sqrt{\left(\frac{5}{3}\right)^2 + (4)^2} dt$$

$$S = 4\pi \int_0^3 (2t + 3) \sqrt{\frac{25}{9} + 16} dt$$

$$S = 4\pi \int_0^3 (2t + 3) \sqrt{\frac{25}{9} + \frac{144}{9}} dt$$



$$S = 4\pi \int_0^3 (2t + 3) \sqrt{\frac{169}{9}} dt$$

$$S = 4\pi \int_0^3 (2t + 3) \frac{13}{3} dt$$

$$S = \frac{52\pi}{3} \int_0^3 2t + 3 dt$$

Integrate, then evaluate over the interval.

$$S = \frac{52\pi}{3} (t^2 + 3t) \Big|_0^3$$

$$S = \frac{52\pi}{3} (3^2 + 3(3)) - \frac{52\pi}{3} (0^2 + 3(0))$$

$$S = \frac{52\pi}{3} (18) - \frac{52\pi}{3} (0)$$

$$S = 312\pi$$

■ 2. Find the surface area of revolution of the parametric curve on the interval $0 \leq t \leq 2\pi$, rotated about the x -axis.

$$x = 3 + \cos t$$

$$y = 4 + \sin t$$



Solution:

Plug the derivatives and the interval into the integral formula for the surface area of revolution for a parametric curve about the x -axis.

$$S = \int_a^b 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$S = \int_0^{2\pi} 2\pi(4 + \sin t) \sqrt{(-\sin t)^2 + (\cos t)^2} dt$$

$$S = \int_0^{2\pi} 2\pi(4 + \sin t) \sqrt{\sin^2 t + \cos^2 t} dt$$

$$S = 2\pi \int_0^{2\pi} (4 + \sin t) \sqrt{1} dt$$

$$S = 2\pi \int_0^{2\pi} 4 + \sin t dt$$

Integrate, then evaluate over the interval.

$$S = 2\pi(4t - \cos t) \Big|_0^{2\pi}$$

$$S = 2\pi(4(2\pi) - \cos(2\pi)) - 2\pi(4(0) - \cos(0))$$

$$S = 2\pi(8\pi - 1) - 2\pi(0 - 1)$$

$$S = 16\pi^2 - 2\pi + 2\pi$$

$$S = 16\pi^2$$



■ 3. Find the surface area of revolution of the parametric curve on the interval $0 \leq t \leq 2\pi$, rotated about the x -axis.

$$x = 7 - 3 \sin t$$

$$y = 6 + 3 \cos t$$

Solution:

Plug the derivatives and the interval into the integral formula formula for the surface area of revolution for a parametric curve about the x -axis.

$$S = \int_a^b 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$S = \int_0^{2\pi} 2\pi(6 + 3 \cos t) \sqrt{(-3 \cos t)^2 + (-3 \sin t)^2} dt$$

$$S = 6\pi \int_0^{2\pi} (2 + \cos t) \sqrt{9 \cos^2 t + 9 \sin^2 t} dt$$

$$S = 6\pi \int_0^{2\pi} (2 + \cos t) \sqrt{9(\cos^2 t + \sin^2 t)} dt$$

$$S = 6\pi \int_0^{2\pi} (2 + \cos t) \sqrt{9(1)} dt$$



$$S = 6\pi \int_0^{2\pi} (2 + \cos t)(3) dt$$

$$S = 18\pi \int_0^{2\pi} 2 + \cos t dt$$

Integrate, then evaluate over the interval.

$$S = 18\pi(2t + \sin t) \Big|_0^{2\pi}$$

$$S = 18\pi(2(2\pi) + \sin(2\pi)) - 18\pi(2(0) + \sin(0))$$

$$S = 18\pi(4\pi + 0) - 18\pi(0 + 0)$$

$$S = 18\pi(4\pi)$$

$$S = 72\pi^2$$

■ 4. Find the surface area of revolution of the parametric curve on the interval $0 \leq t \leq \pi$, rotated about the x -axis.

$$x = 5 - \cos(2t)$$

$$y = 3 + \sin(2t)$$

Solution:



Plug the derivatives and the interval into the integral formula for the surface area of revolution for a parametric curve about the x -axis.

$$S = \int_a^b 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$S = \int_0^\pi 2\pi(3 + \sin(2t))\sqrt{(2\sin(2t))^2 + (2\cos(2t))^2} dt$$

$$S = 2\pi \int_0^\pi (3 + \sin(2t))\sqrt{4\sin^2(2t) + 4\cos^2(2t)} dt$$

$$S = 2\pi \int_0^\pi (3 + \sin(2t))\sqrt{4(\sin^2(2t) + \cos^2(2t))} dt$$

$$S = 2\pi \int_0^\pi (3 + \sin(2t))\sqrt{4(1)} dt$$

$$S = 4\pi \int_0^\pi 3 + \sin(2t) dt$$

Integrate, then evaluate over the interval.

$$S = 4\pi \left(3t - \frac{1}{2} \cos(2t) \right) \Big|_0^\pi$$

$$S = 4\pi \left(3\pi - \frac{1}{2} \cos(2\pi) \right) - 4\pi \left(3(0) - \frac{1}{2} \cos(2(0)) \right)$$

$$S = 4\pi \left(3\pi - \frac{1}{2}(1) \right) - 4\pi \left(0 - \frac{1}{2}(1) \right)$$



$$S = 12\pi^2 - 2\pi + 2\pi$$

$$S = 12\pi^2$$



SURFACE AREA OF REVOLUTION, VERTICAL AXIS

■ 1. Find the surface area of revolution of the parametric curve on the interval $0 \leq t \leq \pi/3$, rotated about the y -axis.

$$x = 8 + \sin(6t)$$

$$y = 7 - \cos(6t)$$

Solution:

Plug the derivatives and the interval into the integral formula formula for the surface area of revolution for a parametric curve about the y -axis.

$$S = \int_a^b 2\pi x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$S = \int_0^{\pi/3} 2\pi(8 + \sin(6t))\sqrt{(6\cos(6t))^2 + (6\sin(6t))^2} dt$$

$$S = 2\pi \int_0^{\pi/3} (8 + \sin(6t))\sqrt{36\cos^2(6t) + 36\sin^2(6t)} dt$$

$$S = 2\pi \int_0^{\pi/3} (8 + \sin(6t))\sqrt{36(\cos^2(6t) + \sin^2(6t))} dt$$

$$S = 2\pi \int_0^{\pi/3} (8 + \sin(6t))\sqrt{36(1)} dt$$



$$S = 12\pi \int_0^{\frac{\pi}{3}} 8 + \sin(6t) \, dt$$

Integrate, then evaluate over the interval.

$$S = 12\pi \left(8t - \frac{1}{6} \cos(6t) \right) \Big|_0^{\frac{\pi}{3}}$$

$$S = 12\pi \left(8 \cdot \frac{\pi}{3} - \frac{1}{6} \cos \left(6 \cdot \frac{\pi}{3} \right) \right) - 12\pi \left(8(0) - \frac{1}{6} \cos(6(0)) \right)$$

$$S = 12\pi \left(\frac{8\pi}{3} - \frac{1}{6} \cos(2\pi) \right) - 12\pi \left(0 - \frac{1}{6} \cos(0) \right)$$

$$S = 12\pi \left(\frac{8\pi}{3} - \frac{1}{6}(1) \right) - 12\pi \left(0 - \frac{1}{6}(1) \right)$$

$$S = 32\pi^2 - 2\pi + 2\pi$$

$$S = 32\pi^2$$

■ 2. Find the surface area of revolution of the parametric curve on the interval $0 \leq t \leq 2\pi$, rotated about the y -axis.

$$x = 5 + 4 \sin(t)$$

$$y = 5 + 4 \cos(t)$$

Solution:



Plug the derivatives and the interval into the integral formula for the surface area of revolution for a parametric curve about the y-axis.

$$S = \int_a^b 2\pi x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$S = \int_0^{2\pi} 2\pi(5 + 4 \sin t) \sqrt{(4 \cos t)^2 + (-4 \sin t)^2} dt$$

$$S = 2\pi \int_0^{2\pi} (5 + 4 \sin t) \sqrt{16 \cos^2 t + 16 \sin^2 t} dt$$

$$S = 2\pi \int_0^{2\pi} (5 + 4 \sin t) \sqrt{16(\cos^2 t + \sin^2 t)} dt$$

$$S = 2\pi \int_0^{2\pi} (5 + 4 \sin t) \sqrt{16(1)} dt$$

$$S = 8\pi \int_0^{2\pi} 5 + 4 \sin t dt$$

Integrate, then evaluate over the interval.

$$S = 8\pi(5t - 4 \cos t) \Big|_0^{2\pi}$$

$$S = 8\pi(5(2\pi) - 4 \cos(2\pi)) - 8\pi(5(0) - 4 \cos(0))$$

$$S = 8\pi(10\pi - 4(1)) - 8\pi(0 - 4(1))$$

$$S = 80\pi^2 - 32\pi + 32\pi$$



$$S = 80\pi^2$$

■ 3. Find the surface area of revolution of the parametric curve on the interval $0 \leq t \leq 2\pi$, rotated about the y -axis.

$$x = 12 - \sin t$$

$$y = 2 + \cos t$$

Solution:

Plug the derivatives and the interval into the integral formula formula for the surface area of revolution for a parametric curve about the y -axis.

$$S = \int_a^b 2\pi x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$S = \int_0^{2\pi} 2\pi(12 - \sin t) \sqrt{(-\cos t)^2 + (-\sin t)^2} dt$$

$$S = \int_0^{2\pi} 2\pi(12 - \sin t) \sqrt{\cos^2 t + \sin^2 t} dt$$

$$S = 2\pi \int_0^{2\pi} (12 - \sin t) \sqrt{1} dt$$

$$S = 2\pi \int_0^{2\pi} 12 - \sin t dt$$



Integrate, then evaluate over the interval.

$$S = 2\pi(12t + \cos t) \Big|_0^{2\pi}$$

$$S = 2\pi(12(2\pi) + \cos(2\pi)) - 2\pi(12(0) + \cos(0))$$

$$S = 2\pi(24\pi + 1) - 2\pi(0 + 1)$$

$$S = 48\pi^2 + 2\pi - 2\pi$$

$$S = 48\pi^2$$

■ 4. Find the surface area of revolution of the parametric curve on the interval $0 \leq t \leq \pi$, rotated about the y -axis.

$$x = 4 - 3 \sin(2t)$$

$$y = 4 - 3 \cos(2t)$$

Solution:

Plug the derivatives and the interval into the integral formula formula for the surface area of revolution for a parametric curve about the y -axis.

$$S = \int_a^b 2\pi x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$



$$S = \int_0^{\pi} 2\pi(4 - 3 \sin(2t))\sqrt{(-6 \cos(2t))^2 + (6 \sin(2t))^2} dt$$

$$S = 2\pi \int_0^{\pi} (4 - 3 \sin(2t))\sqrt{36 \cos^2(2t) + 36 \sin^2(2t)} dt$$

$$S = 2\pi \int_0^{\pi} (4 - 3 \sin(2t))\sqrt{36(\cos^2(2t) + \sin^2(2t))} dt$$

$$S = 2\pi \int_0^{\pi} (4 - 3 \sin(2t))\sqrt{36(1)} dt$$

$$S = 12\pi \int_0^{\pi} 4 - 3 \sin(2t) dt$$

Integrate, then evaluate over the interval.

$$S = 12\pi \left(4t + \frac{3}{2} \cos(2t) \right) \Big|_0^{\pi}$$

$$S = 12\pi \left(4\pi + \frac{3}{2} \cos(2\pi) \right) - 12\pi \left(4(0) + \frac{3}{2} \cos(2(0)) \right)$$

$$S = 12\pi \left(4\pi + \frac{3}{2}(1) \right) - 12\pi \left(0 + \frac{3}{2}(1) \right)$$

$$S = 12\pi \left(4\pi + \frac{3}{2} \right) - 12\pi \left(\frac{3}{2} \right)$$

$$S = 48\pi^2 + 18\pi - 18\pi$$

$$S = 48\pi^2$$



■ 5. Find the surface area of revolution of the parametric curve on the interval $0 \leq t \leq 4$, rotated about the y -axis.

$$x = 6t + 5$$

$$y = 8t + 7$$

Solution:

Plug the derivatives and the interval into the integral formula formula for the surface area of revolution for a parametric curve about the y -axis.

$$S = \int_a^b 2\pi x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$S = \int_0^4 2\pi(6t + 5)\sqrt{(6)^2 + (8)^2} dt$$

$$S = 2\pi \int_0^4 (6t + 5)\sqrt{36 + 64} dt$$

$$S = 2\pi \int_0^4 (6t + 5)\sqrt{100} dt$$

$$S = 20\pi \int_0^4 6t + 5 dt$$

Integrate, then evaluate over the interval.



$$S = 20\pi(3t^2 + 5t) \Big|_0^4$$

$$S = 20\pi(3(4)^2 + 5(4)) - 20\pi(3(0)^2 + 5(0))$$

$$S = 20\pi(3(16) + 20) - 20\pi(0 + 0)$$

$$S = 20\pi(68)$$

$$S = 1,360\pi$$



VOLUME OF REVOLUTION, PARAMETRIC CURVES

- 1. Find the volume of revolution of the parametric curve, rotated about the x -axis, over the interval $1 \leq t \leq 2$.

$$x(t) = 2t^2$$

$$y(t) = 4t^2$$

Solution:

Plug the interval and the parametric equation into the integral formula for volume of revolution for a parametric equation around the x -axis.

$$V_x = \int_a^b \pi y^2 \frac{dx}{dt} dt$$

$$V_x = \int_1^2 \pi (4t^2)^2 (4t) dt$$

$$V_x = 64\pi \int_1^2 t^5 dt$$

Integrate, then evaluate over the interval.

$$V_x = 64\pi \left(\frac{1}{6} t^6 \right) \Big|_1^2$$



$$V_x = \frac{32\pi}{3} t^6 \Big|_1^2$$

$$V_x = \frac{32\pi}{3} (2)^6 - \frac{32\pi}{3} (1)^6$$

$$V_x = \frac{32\pi}{3} (64) - \frac{32\pi}{3}$$

$$V_x = \frac{32\pi}{3} (64 - 1)$$

$$V_x = \frac{32\pi}{3} (63)$$

$$V_x = 32\pi (21)$$

$$V_x = 672\pi$$

■ 2. Find the volume of revolution of the parametric curve, rotated about the y -axis, over the interval $1 \leq t \leq 3$.

$$x(t) = 3t$$

$$y(t) = 4t^2$$

Solution:

Plug the interval and the parametric equation into the integral formula for volume of revolution for a parametric equation around the y -axis.



$$V_y = \int_a^b \pi x^2 \frac{dy}{dt} dt$$

$$V_y = \int_1^3 \pi(3t)^2(8t) dt$$

$$V_y = 72\pi \int_1^3 t^3 dt$$

Integrate, then evaluate over the interval.

$$V_y = 72\pi \left(\frac{1}{4}t^4 \right) \Big|_1^3$$

$$V_y = \frac{72\pi}{4}t^4 \Big|_1^3$$

$$V_y = \frac{72\pi}{4}(3)^4 - \frac{72\pi}{4}(1)^4$$

$$V_y = \frac{72\pi}{4}(81 - 1)$$

$$V_y = \frac{72\pi}{4}(80)$$

$$V_y = 72\pi(20)$$

$$V_y = 1,440\pi$$



- 3. Find the volume of revolution of the parametric curve, rotated about the x -axis, over the interval $1 \leq t \leq 3$.

$$x(t) = 2e^{2t} - 4t$$

$$y(t) = 6e^{\frac{5t}{2}}$$

Solution:

Plug the interval and the parametric equation into the integral formula for volume of revolution for a parametric equation around the x -axis.

$$V_x = \int_a^b \pi y^2 \frac{dx}{dt} dt$$

$$V_x = \int_1^3 \pi (6e^{\frac{5t}{2}})^2 (4e^{2t} - 4) dt$$

$$V_x = 4\pi \int_1^3 (36e^{5t})(e^{2t} - 1) dt$$

$$V_x = 144\pi \int_1^3 e^{5t}e^{2t} - e^{5t} dt$$

$$V_x = 144\pi \int_1^3 e^{7t} - e^{5t} dt$$

Integrate, then evaluate over the interval.



$$V_x = 144\pi \left(\frac{1}{7}e^{7t} - \frac{1}{5}e^{5t} \right) \Big|_1^3$$

$$V_x = 144\pi \left(\frac{1}{7}e^{7(3)} - \frac{1}{5}e^{5(3)} \right) - 144\pi \left(\frac{1}{7}e^{7(1)} - \frac{1}{5}e^{5(1)} \right)$$

$$V_x = 144\pi \left(\frac{1}{7}e^{21} - \frac{1}{5}e^{15} \right) - 144\pi \left(\frac{1}{7}e^7 - \frac{1}{5}e^5 \right)$$

$$V_x = 144\pi \left(\frac{1}{7}e^{21} - \frac{1}{5}e^{15} - \frac{1}{7}e^7 + \frac{1}{5}e^5 \right)$$

$$V_x = 144\pi \left(\frac{e^{21} - e^7}{7} - \frac{e^{15} - e^5}{5} \right)$$

■ 4. Find the volume of revolution of the parametric curve, rotated about the y -axis, over the interval $0 \leq t \leq 1$.

$$x(t) = 3e^t$$

$$y(t) = e^t$$

Solution:

Plug the interval and the parametric equation into the integral formula for volume of revolution for a parametric equation around the y -axis.

$$V_y = \int_a^b \pi x^2 \frac{dy}{dt} dt$$



$$V_y = \int_0^1 \pi(3e^t)^2(e^t) dt$$

$$V_y = 9\pi \int_0^1 (e^{2t})(e^t) dt$$

$$V_y = 9\pi \int_0^1 e^{3t} dt$$

Integrate, then evaluate over the interval.

$$V_y = 9\pi \left(\frac{1}{3} e^{3t} \right) \Big|_0^1$$

$$V_y = 3\pi (e^{3t}) \Big|_0^1$$

$$V_y = 3\pi (e^{3(1)}) - 3\pi (e^{3(0)})$$

$$V_y = 3\pi(e^3) - 3\pi(1)$$

$$V_y = 3\pi(e^3 - 1)$$



