

Topic: Area under a parametric curve

Question: Find the area under the parametric curve.

$$x = t - \sin t$$

$$y = 6(1 - \cos t)$$

$$0 \leq t \leq 2\pi$$

Answer choices:

A 18

B 231π

C 18π

D $18\pi - 13$



Solution: C

We'll find the area under the curve using the integral formula

$$A = \int_{\alpha}^{\beta} y(t)x'(t) dt$$

Since the question defines an interval for our parameter, we'll be able to find a real-number answer for the area.

We've already been given $y(t)$, but we need to find $x'(t)$ before we can plug into the area formula.

$$x(t) = t - \sin t$$

$$x'(t) = 1 - \cos t$$

Plugging into our area formula, we get

$$A = \int_0^{2\pi} 6(1 - \cos t)(1 - \cos t) dt$$

$$A = 6 \int_0^{2\pi} 1 - 2 \cos t + \cos^2 t dt$$

Using the power reduction formula from trigonometry,

$$\cos^2 t = \frac{1}{2} + \frac{1}{2} \cos 2t$$

we get

$$A = 6 \int_0^{2\pi} 1 - 2 \cos t + \frac{1}{2} + \frac{1}{2} \cos 2t dt$$



$$A = 6 \int_0^{2\pi} \frac{3}{2} - 2 \cos t + \frac{1}{2} \cos 2t \, dt$$

$$A = 6 \left(\frac{3}{2}t - 2 \sin t + \frac{1}{4} \sin 2t \right) \Big|_0^{2\pi}$$

$$A = 6 \left[\left(\frac{3}{2}(2\pi) - 2 \sin(2\pi) + \frac{1}{4} \sin 2(2\pi) \right) - \left(\frac{3}{2}(0) - 2 \sin(0) + \frac{1}{4} \sin 2(0) \right) \right]$$

$$A = 6 \left[\left(3\pi - 2(0) + \frac{1}{4} \sin 4\pi \right) - \left(0 - 2(0) + \frac{1}{4}(0) \right) \right]$$

$$A = 6 \left(3\pi + \frac{1}{4}(0) \right)$$

$$A = 6(3\pi)$$

$$A = 18\pi$$



Topic: Area under a parametric curve

Question: Find the area under the parametric curve.

$$f(t) = 4t^2$$

$$g(t) = t - 4$$

$$0 \leq t \leq 4$$

Answer choices:

A 256

B $\frac{256}{3}$

C -256

D $-\frac{256}{3}$



Solution: D

We'll find the area under the curve using the integral formula

$$A = \int_{\alpha}^{\beta} y(t)x'(t) \, dt$$

Since the question defines an interval for our parameter, we'll be able to find a real-number answer for the area.

We've already been given $y(t)$ as $g(t) = t - 4$, but we need to find $x'(t)$ as $f'(t)$ before we can plug into the area formula.

$$f(t) = 4t^2$$

$$f'(t) = 8t$$

Plugging into our area formula, we get

$$A = \int_0^4 (t - 4)(8t) \, dt$$

$$A = \int_0^4 8t^2 - 32t \, dt$$

$$A = \left. \frac{8}{3}t^3 - 16t^2 \right|_0^4$$

$$A = \frac{8}{3}(4)^3 - 16(4)^2 - \left[\frac{8}{3}(0)^3 - 16(0)^2 \right]$$

$$A = \frac{512}{3} - \frac{768}{3}$$



$$A = -\frac{256}{3}$$



Topic: Area under a parametric curve**Question:** Find the area under the parametric curve.

$$f(t) = e^t$$

$$g(t) = 1 - t^2$$

$$0 \leq t \leq 4$$

Answer choices:

A $9e^4 - 1$

B $12e^4$

C $1 - 9e^4$

D $9e^4$



Solution: C

We'll find the area under the curve using the integral formula

$$A = \int_{\alpha}^{\beta} y(t)x'(t) \, dt$$

Since the question defines an interval for our parameter, we'll be able to find a real-number answer for the area.

We've already been given $y(t)$ as $g(t) = 1 - t^2$, but we need to find $x'(t)$ as $f'(t)$ before we can plug into the area formula.

$$f(t) = e^t$$

$$f'(t) = e^t$$

Plugging into our area formula, we get

$$A = \int_0^4 (1 - t^2) (e^t) \, dt$$

$$A = \int_0^4 e^t - t^2 e^t \, dt$$

$$A = \int_0^4 e^t \, dt - \int_0^4 t^2 e^t \, dt$$

For the second integral, we'll use integration by parts, setting

$$u = t^2$$

$$du = 2t \, dt$$



and

$$dv = e^t dt$$

$$v = e^t$$

Replacing the second integral using the integration by parts formula

$$\int u dv = uv - \int v du$$

we get

$$A = \int_0^4 e^t dt - \left[t^2 e^t \Big|_0^4 - \int_0^4 2te^t dt \right]$$

We'll need to use integration by parts again on this new integral. Setting

$$u = 2t$$

$$du = 2 dt$$

and

$$dv = e^t dt$$

$$v = e^t$$

Using the integration by parts formula to replace the integral again, we get

$$A = \int_0^4 e^t dt - \left[t^2 e^t \Big|_0^4 - \left[2te^t \Big|_0^4 - \int_0^4 2e^t dt \right] \right]$$



$$A = \int_0^4 e^t dt - \left[t^2 e^t \Big|_0^4 - 2te^t \Big|_0^4 + \int_0^4 2e^t dt \right]$$

$$A = \int_0^4 e^t dt - t^2 e^t \Big|_0^4 + 2te^t \Big|_0^4 - \int_0^4 2e^t dt$$

$$A = e^t \Big|_0^4 - t^2 e^t \Big|_0^4 + 2te^t \Big|_0^4 - 2e^t \Big|_0^4$$

$$A = (e^t - t^2 e^t + 2te^t - 2e^t) \Big|_0^4$$

$$A = [e^4 - 4^2 e^4 + 2(4)e^4 - 2e^4] - [e^0 - 0^2 e^0 + 2(0)e^0 - 2e^0]$$

$$A = e^4 - 16e^4 + 8e^4 - 2e^4 - (1 - 0 + 0 - 2(1))$$

$$A = -9e^4 + 1$$

$$A = 1 - 9e^4$$

