



# Calculus 1 Workbook Solutions

---

Definition of the derivative

## DEFINITION OF THE DERIVATIVE

■ 1. Use the definition of the derivative to find the derivative of  $f(x) = 2x^2 + 2x - 12$  at  $(4,28)$ .

*Solution:*

At  $(4,28)$ , the derivative will be

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{f(4+h) - f(4)}{h}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{(2(4+h)^2 + 2(4+h) - 12) - (2(4)^2 + 2(4) - 12)}{h}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{(2(16 + 8h + h^2) + 8 + 2h - 12) - (32 + 8 - 12)}{h}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{32 + 16h + 2h^2 + 8 + 2h - 12 - 32 - 8 + 12}{h}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{18h + 2h^2}{h}$$

$$f'(4) = \lim_{h \rightarrow 0} (18 + 2h)$$

Evaluate the limit to find the derivative at  $(4,28)$ .



$$f'(4) = 18 + 2(0)$$

$$f'(4) = 18$$

■ 2. Use the definition of the derivative to find the derivative of  $g(x) = 3x^3 - 4x + 7$  at  $(-2, -9)$ .

*Solution:*

At  $(-2, -9)$ , the derivative will be

$$g'(x) = \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h}$$

$$g'(-2) = \lim_{h \rightarrow 0} \frac{g(-2+h) - g(-2)}{h}$$

$$g'(-2) = \lim_{x \rightarrow 0} \frac{(3(-2+h)^3 - 4(-2+h) + 7) - (3(-2)^3 - 4(-2) + 7)}{h}$$

$$g'(-2) = \lim_{x \rightarrow 0} \frac{(3(-8 + 4h + 8h - 4h^2 - 2h^2 + h^3) + 8 - 4h + 7) - (-24 + 8 + 7)}{h}$$

$$g'(-2) = \lim_{x \rightarrow 0} \frac{-24 + 12h + 24h - 12h^2 - 6h^2 + 3h^3 + 8 - 4h + 7 + 24 - 8 - 7}{h}$$

$$g'(-2) = \lim_{x \rightarrow 0} \frac{3h^3 - 18h^2 + 32h}{h}$$

$$g'(-2) = \lim_{x \rightarrow 0} (3h^2 - 18h + 32)$$



Evaluate the limit to find the derivative at  $(-2, -9)$ .

$$g'(-2) = 3(0)^2 - 18(0) + 32$$

$$g'(-2) = 32$$

■ 3. Use the definition of the derivative to find the derivative at  $(-1, -1)$ .

$$f(x) = \frac{x}{x+2}$$

*Solution:*

At  $(-1, -1)$ , the derivative will be

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$f'(-1) = \lim_{h \rightarrow 0} \frac{f(-1+h) - f(-1)}{h}$$

$$f'(-1) = \lim_{h \rightarrow 0} \frac{\frac{-1+h}{-1+h+2} - \frac{-1}{-1+2}}{h}$$

$$f'(-1) = \lim_{h \rightarrow 0} \frac{\frac{-1+h}{h+1} - \frac{-1}{1}}{h}$$

$$f'(-1) = \lim_{h \rightarrow 0} \frac{\frac{-1+h}{h+1} + 1}{h}$$



$$f'(-1) = \lim_{h \rightarrow 0} \frac{\frac{-1 + h + h + 1}{h + 1}}{h}$$

$$f'(-1) = \lim_{h \rightarrow 0} \frac{2h}{h(h + 1)}$$

$$f'(-1) = \lim_{h \rightarrow 0} \frac{2}{h + 1}$$

Evaluate the limit to find the derivative at  $(-1, -1)$ .

$$f'(-1) = \frac{2}{0 + 1}$$

$$f'(-1) = 2$$

■ 4. Use the definition of the derivative to find the derivative of  $f(x) = \sqrt{5x - 4}$  at  $x = 4$ .

*Solution:*

At  $x = 4$ , the derivative will be

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{f(4 + h) - f(4)}{h}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{\sqrt{5(4 + h) - 4} - \sqrt{5(4) - 4}}{h}$$



$$f'(4) = \lim_{h \rightarrow 0} \frac{\sqrt{20 + 5h} - 4}{h}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{\sqrt{16 + 5h} - \sqrt{16}}{h}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{\sqrt{16 + 5h} - 4}{h}$$

Use conjugate method.

$$f'(4) = \lim_{h \rightarrow 0} \frac{\sqrt{16 + 5h} - 4}{h} \cdot \frac{\sqrt{16 + 5h} + 4}{\sqrt{16 + 5h} + 4}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{16 + 5h - 16}{h(\sqrt{16 + 5h} + 4)}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{5h}{h(\sqrt{16 + 5h} + 4)}$$

$$f'(4) = \lim_{h \rightarrow 0} \frac{5}{\sqrt{16 + 5h} + 4}$$

Evaluate the limit to find the derivative at  $x = 4$ .

$$f'(4) = \frac{5}{\sqrt{16 + 5(0)} + 4}$$

$$f'(4) = \frac{5}{4 + 4}$$



$$f'(4) = \frac{5}{8}$$

■ 5. Use the definition of the derivative to find the derivative of  $g(x) = \cos(x - 1)$  at  $x = \pi/2$ .

*Solution:*

At  $x = \pi/2$ , the derivative will be

$$g'(x) = \lim_{h \rightarrow 0} \frac{g(x + h) - g(x)}{h}$$

$$g'(x) = \lim_{h \rightarrow 0} \frac{\cos(x + h - 1) - \cos(x - 1)}{h}$$

$$g'(x) = \lim_{h \rightarrow 0} \frac{\cos(x - 1 + h) - \cos(x - 1)}{h}$$

Apply the sum-identity for cosine.

$$g'(x) = \lim_{h \rightarrow 0} \frac{\cos(x - 1)\cos h - \sin(x - 1)\sin h - \cos(x - 1)}{h}$$

$$g'(x) = \lim_{h \rightarrow 0} \frac{\cos(x - 1)(\cos h - 1) - \sin(x - 1)\sin h}{h}$$

$$g'(x) = \lim_{h \rightarrow 0} \left( \frac{\cos(x - 1)(\cos h - 1)}{h} - \frac{\sin(x - 1)\sin h}{h} \right)$$

Pull the expressions with  $x$  out in front of the limits, which only apply to  $h$ .



$$g'(x) = \cos(x-1) \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} - \sin(x-1) \lim_{h \rightarrow 0} \frac{\sin h}{h}$$

Applying formulas for the limits of these common trigonometric functions, we get

$$g'(x) = \cos(x-1)(0) - \sin(x-1)(1)$$

$$g'(x) = -\sin(x-1)$$

Evaluate the limit to find the derivative at  $x = \pi/2$ .

$$g'\left(\frac{\pi}{2}\right) = -\sin\left(\frac{\pi}{2} - 1\right)$$

If we choose, we could apply the cofunction identity for cosine,

$$\cos \theta = \sin\left(\frac{\pi}{2} - \theta\right)$$

Using that identity allows us to simplify the answer to

$$g'\left(\frac{\pi}{2}\right) = -\cos(1)$$

■ 6. Use the definition of the derivative to find the derivative of  $g(x) = |x|$  at  $x = 0$ .

*Solution:*





At  $x = 0$ , the derivative will be

$$g'(x) = \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h}$$

$$g'(0) = \lim_{h \rightarrow 0} \frac{g(0+h) - g(0)}{h}$$

$$g'(0) = \lim_{h \rightarrow 0} \frac{|h| - |0|}{h}$$

$$g'(0) = \lim_{h \rightarrow 0} \frac{|h|}{h}$$

We know that

$$|h| = \begin{cases} -h & h < 0 \\ h & h \geq 0 \end{cases}$$

$$\lim_{h \rightarrow 0^-} \frac{|h|}{h} = \lim_{h \rightarrow 0^-} \frac{-h}{h} = -1$$

$$\lim_{h \rightarrow 0^+} \frac{|h|}{h} = \lim_{h \rightarrow 0^+} \frac{h}{h} = 1$$

Since the left- and right-hand limits aren't equal, this limit doesn't exist, which means the function isn't differentiable at  $x = 0$ , and we can't find its derivative there.



