

FALL 2021 EXAMINATION II

Duration: 150 mins

There are 8 questions (total 30 marks). Answer all questions.

A permitted calculator may be used.

1. (3 marks) Write the Hamiltonian for H_2 and indicate the terms that comprise the electronic Hamiltonian.

$$\hat{H} = -\frac{1}{2M_1}\nabla_{\mathbf{R}_1}^2 - \frac{1}{2M_2}\nabla_{\mathbf{R}_2}^2 - \frac{1}{2}\nabla_{\mathbf{r}_1}^2 - \frac{1}{2}\nabla_{\mathbf{r}_2}^2 - \frac{1}{|\mathbf{R}_1-\mathbf{R}_2|} - \frac{1}{|\mathbf{r}_1-\mathbf{R}_1|} - \frac{1}{|\mathbf{r}_2-\mathbf{R}_1|} - \frac{1}{|\mathbf{r}_1-\mathbf{R}_2|} - \frac{1}{|\mathbf{r}_2-\mathbf{R}_2|} + \frac{1}{|\mathbf{r}_1-\mathbf{r}_2|}$$
$$\hat{H}_{elec} = -\frac{1}{2}\nabla_{\mathbf{r}_1}^2 - \frac{1}{2}\nabla_{\mathbf{r}_2}^2 - \frac{1}{|\mathbf{r}_1-\mathbf{R}_1|} - \frac{1}{|\mathbf{r}_2-\mathbf{R}_1|} - \frac{1}{|\mathbf{r}_1-\mathbf{R}_2|} - \frac{1}{|\mathbf{r}_2-\mathbf{R}_2|} + \frac{1}{|\mathbf{r}_1-\mathbf{r}_2|}$$

2. (3 marks) Determine the number of cartesian contracted and primitive Gaussian functions in a molecular electronic structure calculation on ethanal CH_3CHO using (i) 6-31G (ii) 6-31G* and (iii) 6-31+G* basis sets.

i) 35 contracted Gaussian functions, 82 primitive Gaussian functions

ii) Cartesian d-orbitals:

53 contracted Gaussian functions, 100 primitive Gaussian functions

iii) Cartesian d-orbitals:

65 contracted Gaussian functions, 112 primitive Gaussian functions

3. (4 marks) Write the following integrals in bra-ket notation:

i) $\int f^*(x)g(x)dx$

$$\langle f(x)|g(x)\rangle \quad (1)$$

ii) $\int f(\tau)g(\tau)d\tau$

$$\langle f^*(\tau)|g(\tau)\rangle \quad (2)$$

iii) $(\hat{H} + x^2)\psi(x) = E\psi(x)$

$$(\hat{H} + x^2)|\psi(x)\rangle = E|\psi(x)\rangle \quad (3)$$

$$\text{iv) } \psi(x) \int \psi^*(x') f(x') dx' + 3g(x)$$

$$|\psi(x)\rangle \langle \psi(x') | f(x') \rangle + 3|g(x)\rangle \quad (4)$$

4. (4 marks) Compute the Hückel molecular orbital energies of propene.

$$\begin{vmatrix} \alpha - E & \beta & 0 \\ \beta & \alpha - E & \beta \\ 0 & \beta & \alpha - E \end{vmatrix} = 0$$

$$\begin{vmatrix} -x & 1 & 0 \\ 1 & -x & 1 \\ 0 & 1 & -x \end{vmatrix} = 0$$

$$-x(x^2 - 1) + x = x(-x^2 + 2) = 0$$

$$x = 0, \pm\sqrt{2} \quad E_0 = \alpha + \sqrt{2}\beta, \quad E_1 = \alpha, \quad E_2 = \alpha - \sqrt{2}\beta$$

5. (4 marks) Confirm that the matrices $\mathbf{s}_w = \frac{\hbar}{2}\boldsymbol{\sigma}_w$, where $w = \{x, y, z\}$ and $\boldsymbol{\sigma}_w$ are the Pauli spin matrices, form a representation of angular momentum. Show that these matrices are a representation of $s = \frac{1}{2}$. The Pauli spin matrices are:

$$\boldsymbol{\sigma}_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \boldsymbol{\sigma}_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \boldsymbol{\sigma}_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

To demonstrate the matrices are a representation of angular momentum, we must show they obey the commutation relations.

$$\begin{aligned} [\mathbf{s}_x, \mathbf{s}_y] &= \frac{\hbar}{2}\boldsymbol{\sigma}_x \frac{\hbar}{2}\boldsymbol{\sigma}_y - \frac{\hbar}{2}\boldsymbol{\sigma}_y \frac{\hbar}{2}\boldsymbol{\sigma}_x = \frac{\hbar^2}{4} \left\{ \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} - \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\} \\ &= \frac{\hbar^2}{4} \left\{ \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} - \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} \right\} = \frac{\hbar^2}{4} \begin{pmatrix} 2i & 0 \\ 0 & -2i \end{pmatrix} = i\hbar \frac{\hbar}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = i\hbar \mathbf{s}_z \end{aligned}$$

Construct the matrix representation of \mathbf{s}^2 and then extract eigenvalues which should be $s(s+1)\hbar^2$ then solve for s .

$$\begin{aligned} \mathbf{s}^2 &= \mathbf{s}_x^2 + \mathbf{s}_y^2 + \mathbf{s}_z^2 = \frac{\hbar^2}{4} \left\{ \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right\} \\ &= \frac{\hbar^2}{4} \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right\} = \frac{3\hbar^2}{4} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \end{aligned}$$

As $s(s+1) = \frac{3}{4}$, so $s = 1/2$

6. (4 marks) Write the electronic energy of a restricted-open shell Slater determinant representing the i) doublet and ii) quartet wavefunction of BeH.

$${}^2E_{\text{ROHF}} = 2h_{11} + 2h_{22} + h_{33} + J_{11} + J_{22} + 4J_{12} + 2J_{13} + 2J_{23} - 2K_{12} - K_{13} - K_{23} \quad (5)$$

$${}^4E_{\text{ROHF}} = 2h_{11} + h_{22} + h_{33} + h_{44} + J_{11} + 2J_{12} + 2J_{13} + 2J_{14} + J_{23} + J_{24} + J_{34} - K_{12} - K_{13} - K_{14} - K_{23} - K_{24} - K_{34} \quad (6)$$

7. (4 marks) Consider the ground state of the hydrogen molecule with a wavefunction represented by a restricted Slater determinant, such that two electrons occupy the same spatial orbital $\psi_1(\mathbf{r})$.

i) Write out explicitly the terms contained in the Fock operator for one of the electrons.

$$\hat{F} = \hat{h} + \hat{J}_2 \quad (7)$$

ii) Compute the orbital energy of the electron described by the Fock operator in part i).

$$\langle \psi_1 | \hat{F} | \psi_1 \rangle = \langle \psi_1 | \hat{h} | \psi_1 \rangle + \langle \psi_1 | \hat{J}_2 | \psi_1 \rangle = \langle \psi_1 | \hat{h} | \psi_1 \rangle + \langle \psi_1 \psi_2 | \psi_2 \psi_1 \rangle = h_{11} + J_{12} = \varepsilon_1 \quad (8)$$

iii) Write down the ionization energy of the electron in part ii) according to Koopmans' theorem.

$$\text{IE} = -\varepsilon_1 \quad (9)$$

iv) Write the approximations involved in computing ionization energies using Koopmans' theorem.

- Orbital relaxation in response to ionization
- Electron correlation (going beyond mean-field approximation)

8. (4 marks) Given the Slater determinant wavefunction $|\Psi\rangle = N|\psi_1(\mathbf{x}_1)\psi_2(\mathbf{x}_2)\rangle$, determine the normalization constant N .

$$N^2 \langle \psi_1(\mathbf{x}_1) \psi_2(\mathbf{x}_2) | \psi_1(\mathbf{x}_1) \psi_2(\mathbf{x}_2) \rangle = 1$$

$$N^2 \langle \{ \psi_1(\mathbf{x}_1) \psi_2(\mathbf{x}_2) - \psi_1(\mathbf{x}_2) \psi_2(\mathbf{x}_1) \} | \{ \psi_1(\mathbf{x}_1) \psi_2(\mathbf{x}_2) - \psi_1(\mathbf{x}_2) \psi_2(\mathbf{x}_1) \} \rangle = 1$$

$$N^2 \{ \langle \psi_1(\mathbf{x}_1) \psi_2(\mathbf{x}_2) | \psi_1(\mathbf{x}_1) \psi_2(\mathbf{x}_2) \rangle + \langle \psi_1(\mathbf{x}_2) \psi_2(\mathbf{x}_1) | \psi_1(\mathbf{x}_2) \psi_2(\mathbf{x}_1) \rangle \\ - \langle \psi_1(\mathbf{x}_1) \psi_2(\mathbf{x}_2) | \psi_1(\mathbf{x}_2) \psi_2(\mathbf{x}_1) \rangle - \langle \psi_1(\mathbf{x}_2) \psi_2(\mathbf{x}_1) | \psi_1(\mathbf{x}_1) \psi_2(\mathbf{x}_2) \rangle \} = 1$$

$$N^2 \{ \langle \psi_1(\mathbf{x}_1) | \psi_1(\mathbf{x}_1) \rangle \langle \psi_2(\mathbf{x}_2) | \psi_2(\mathbf{x}_2) \rangle + \langle \psi_1(\mathbf{x}_2) | \psi_1(\mathbf{x}_2) \rangle \langle \psi_2(\mathbf{x}_1) | \psi_2(\mathbf{x}_1) \rangle \\ - \langle \psi_1(\mathbf{x}_1) | \psi_2(\mathbf{x}_1) \rangle \langle \psi_2(\mathbf{x}_2) | \psi_1(\mathbf{x}_2) \rangle - \langle \psi_2(\mathbf{x}_1) | \psi_1(\mathbf{x}_1) \rangle \langle \psi_1(\mathbf{x}_2) | \psi_2(\mathbf{x}_2) \rangle \} = 1$$

$$N^2 \{ 1 \times 1 + 1 \times 1 - 0 \times 0 - 0 \times 0 \} = 1$$

$$2N^2 = 1$$

$$N = \frac{1}{\sqrt{2}}$$

END OF PAPER