




normalised rate maps | x = fraction of session, y = fraction of goal-goal transition | neurons 0-47 of 1252
 per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise




normalised rate maps | x = fraction of session, y = fraction of goal-goal transition | neurons 48-95 of 1252
per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise





normalised rate maps | x = fraction of session, y = fraction of goal-goal transition | neurons 96-143 of 1252
 per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise








normalised rate maps | x = fraction of session, y = fraction of goal-goal transition
 per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise




















normalised rate maps | x = fraction of session, y = fraction of goal-goal transition | neurons 528-575 of 1252
 per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise





normalised rate maps | x = fraction of session, y = fraction of goal-goal transition | neurons 576-623 of 1252
 per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise








normalised rate maps | x = fraction of session, y = fraction of goal-goal transition | neurons 672-719 of 1252
 per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise














normalised rate maps | x = fraction of session, y = fraction of goal-goal transition | neurons 912-959 of 1252
 per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise







normalised rate maps | x = fraction of session, y = fraction of goal-goal transition | neurons 1008-1055 of 1252
per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise





normalised rate maps | x = fraction of session, y = fraction of goal-goal transition | neurons 1056-1103 of 1252
per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise





normalised rate maps | x = fraction of session, y = fraction of goal-goal transition | neurons 1152-1199 of 1252
 per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise





normalised rate maps | x = fraction of session, y = fraction of goal-goal transition | neurons 1248-1251 of 1252
per-panel peak-normalised, peak Hz in title; no reliability measure computed - many LEC maps will be Poisson noise

