

Dark stars meeting notes Jan 18

[Review of Structure Formation in Dark Sector paper](#)

[Introduction](#)

[Section 2](#)

[Section 3](#)

[Section 4](#)

[Section 4.2](#)

[Section 4.3](#)

[Objectives for next meeting](#)

Review of Structure Formation in Dark Sector paper

Introduction

- Adopting the principles of star formation to the dark sector
- Model: dark electron and dark photon (massive boson)
 - Dark photon is massive to get the right structure formation
- Jurgens idea — use dark boson instead of dark electron

Section 2

- Assume QED regime to do perturbation theory
- Dont need Lagrangian
- Mean free path is important
 - σ is the cross section of interaction (probability of interaction)
 - tells the distance a particle can travel without interaction
 - inversely related to number density
- If dark electrons scatter (self-interact) and photons escape, dark stars can cool and for stable configurations

- The idea is to compute this possibility
- Equation of state for a fermi gas — ideal gas law
 - Eq 3.4 (13) is the ideal gas law for the dark electron
 - 1st term: ideal gas law with $k_B=1$
 - 2nd term: field theory interaction term

Section 3

- Jeans length for dark stars
- Hubble expansion not important for us
- Turnaround into non-linear regime

Section 4

- Eq 4.1 (28) Jeans condition to form dark stars

Section 4.2

- IMPORTANT equations for our project
- Lambda is a cooling rate: comes from a particle physics model
 - Eq 4.13 (40) BS cooling rate is exponentially suppressed
 - We can instead use the simpler Stefan-Boltzmann law
- Eq. 4.5 (32) is an energy balance equation
- Cooling time scale ansatz
 - we can use the ones in 4.12 (39) or choose our own
 - cooling from the surface — use Stefan-Boltzmann law
- Interested in cooling of dark star into a stable configuration
 - We will need to solve the TOV equations

Section 4.3

- Figure 5 shows number density and temperature over the evolution of dark model

- Density increases, heat increases, cooling rate increases
- Eventually the energy of compression matches the rate of photon emission
- Balance of compression and cooling at the $t_{\text{cooling}}=t_{\text{free}}$ fall time scale along jeans mass lines
- This is the regime in which we can form stable configurations
- Optically thick regime: star is so dense photons cannot escape anymore
- Hierarchical fragmentation happens along the cooling line until the optically thick regime is reached
- Will want to recreate Figure 5 with our model
- Figure 7 shows the mass of compact objects from fragmentation as a function of the dark electron and dark photon masses

Objectives for next meeting

- Reread Structure Formation paper focus on Sec 4
- Review hierarchical fragmentation from Ryden
- Solve the TOV for Boson stars
 - Create TOV solver code in dimensionless units
 - Equation of state for boson stars also in dimensionless units
 - Boson stars with self-interactions on slack Eq. 16
 - relation between pressure and energy
 - energy density as a function of pressure
 -
- Dimensionless TOV equations in Compact Stars chapter 4.3 (scaling solutions)
- Eq. 4.51 and 4.52
- Chapter 4.5 applied to Boson stars

- Create your own Dark Star!