COMP0147 Discrete Mathematics for Computer Scientists Notes

Joe

March 28, 2019

Notes adapted from lecture notes by Max Kanovich and Robin Hirsch [1].

Contents

1	Fou	oundations				
	1.1	Set Th	neory	5		
		1.1.1	Set Notations	5		
		1.1.2	Properties	5		
		1.1.3	Set Equality	6		
		1.1.4	Set Operations	6		
		1.1.5	Boolean Algebra	6		
		1.1.6	Set Algebra	7		
	1.2	Funct	ions	9		
		1.2.1	Composition of Injections	10		

1 Foundations

Contents

1.1	Set Th	heory
	1.1.1	Set Notations 5
	1.1.2	Properties
	1.1.3	Set Equality
	1.1.4	Set Operations 6
	1.1.5	Boolean Algebra 6
	1.1.6	Set Algebra
1.2	Funct	ions
	1.2.1	Composition of Injections

1.1 Set Theory

1.1.1 Set Notations

- Set definition: $A = \{a, b, c\}$
- Set membership (element-of): $a \in A$
- Set builder notation: $\{x \mid x \in \mathbb{R} \land x^2 = x\}$
- Empty set: ∅

1.1.2 Properties

- No structure
- No order
- No copies

For example, a,b,c are references to actual objects in

$$\{a,b,c\} \Leftrightarrow \{c,a,b\} \Leftrightarrow \{a,b,c,b\}$$

1.1.3 Set Equality

Definition 1.1.1 (Set Equality). Set A = B iff:

- 1. $A \subseteq B \implies \forall x(x \in A \rightarrow x \in B)$
- 2. $B \subseteq A \implies \forall y(y \in B \rightarrow y \in A)$

Remark. $A = B \Leftrightarrow A \subseteq B \land B \subseteq A$

1.1.4 Set Operations

- Union: $A \cup B \equiv \{x \mid x \in A \lor x \in B\}$
- *Intersection*: $A \cap B \equiv \{x \mid x \in A \land x \in B\}$
- Relative Complement: $A \setminus B \equiv \{x \mid x \in A \land x \notin B\}$
- Absolute Complement: $A^c \equiv U \setminus A \equiv \{x \mid x \in U \land x \notin A\}$
- Symmetric Difference: $A\Delta B \equiv (A \setminus B) \cup (B \setminus A) \equiv (A \cup B) \setminus (A \cap B)$
- Cartesian Product: $A \times B \equiv \{(x, y) \mid x \in A \land y \in B\}$

1.1.5 Boolean Algebra

Definition 1.1.2 (De Morgan's Laws).

$$\neg (p \lor q) \equiv \neg p \land \neg q \tag{1.1}$$

$$\neg (p \land q) \equiv \neg p \lor \neg q \tag{1.2}$$

Definition 1.1.3 (Idempotent Laws).

$$p \lor p \equiv p \tag{1.3}$$

$$p \wedge p \equiv p \tag{1.4}$$

Definition 1.1.4 (Commutative Laws).

$$p \lor q \equiv q \lor p \tag{1.5}$$

$$p \wedge q \equiv q \wedge p \tag{1.6}$$

Definition 1.1.5 (Associative Laws).

$$p \lor (q \lor r) \equiv (p \lor q) \lor r \tag{1.7}$$

$$p \wedge (q \wedge r) \equiv (p \wedge q) \wedge r \tag{1.8}$$

Definition 1.1.6 (Distributive Laws).

$$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r) \tag{1.9}$$

$$p \lor (q \land r) \equiv (p \lor q) \land (p \lor r) \tag{1.10}$$

Definition 1.1.7 (Identity Laws).

$$p \vee F \equiv p \tag{1.11}$$

$$p \vee T \equiv T \tag{1.12}$$

$$p \wedge T \equiv p \tag{1.13}$$

$$p \wedge F \equiv F \tag{1.14}$$

Definition 1.1.8 (Absorption Laws).

$$p \lor (p \land q) \equiv p \tag{1.15}$$

$$p \land (p \lor q) \equiv p \tag{1.16}$$

Definition 1.1.9 (Implication and Negation Laws).

- *Identity*: $p \rightarrow q \equiv \neg p \lor q$
- Counter-example: $\neg(p \to q) \equiv p \land \neg q$
- Equivalences: $p \to q \to r \equiv (p \land q) \to r \equiv q \ to(p \to r)$
- Absorption:

$$p \to T \equiv T$$

$$p \to \mathcal{F} \equiv \neg p$$

$$\mathrm{T} \to p \equiv p$$

$$F \to p \equiv T$$

- Contrapositive: $p \rightarrow q \equiv \neg q \rightarrow \neg p$
- Law of Excluded Middle:

$$p \vee \neg p \equiv \mathbf{T}$$

$$p \land \neg p \equiv F$$

- Double Negation: $\neg \neg p \equiv p$
- Reduction to Absurdity: $\neg p \rightarrow F \equiv p$

1.1.6 Set Algebra

Definition 1.1.10 (De Morgan's Laws).

$$(A \cup B)^c \equiv A^c \cap B^c \tag{1.17}$$

$$(A \cap B)^c \equiv A^c \cup B^c \tag{1.18}$$

Definition 1.1.11 (Idempotent Laws).

$$A \cup A \equiv A \tag{1.19}$$

$$A \cap A \equiv A \tag{1.20}$$

Definition 1.1.12 (Commutative Laws).

$$A \cup B \equiv B \cup A \tag{1.21}$$

$$A \cap B \equiv B \cap A \tag{1.22}$$

Definition 1.1.13 (Associativity Laws).

$$A \cup (B \cup C) \equiv (A \cup B) \cup C \tag{1.23}$$

$$A \cap (B \cap C) \equiv (A \cap B) \cap C \tag{1.24}$$

Definition 1.1.14 (Distributive Laws).

$$A \cap (B \cup C) \equiv (A \cap B) \cup (B \cap C) \tag{1.25}$$

$$A \cup (B \cap C) \equiv (A \cup B) \cap (B \cup C) \tag{1.26}$$

Definition 1.1.15 (Identity Laws).

$$A \cup \emptyset \equiv A \tag{1.27}$$

$$A \cap \emptyset \equiv \emptyset \tag{1.28}$$

$$A \cap U \equiv A \tag{1.29}$$

$$A \cup U \equiv U \tag{1.30}$$

Definition 1.1.16 (Absorption Laws).

$$A \cup (A \cap B) \equiv A \tag{1.31}$$

$$A \cap (A \cup B) \equiv A \tag{1.32}$$

Definition 1.1.17 (Difference Identity Laws).

$$C \setminus (A \cup B) \equiv (C \setminus A) \cap (C \setminus B) \tag{1.33}$$

$$C \setminus (A \cap B) \equiv (C \setminus A) \cup (C \setminus B) \tag{1.34}$$

Definition 1.1.18 (Complement-Difference Identity Law).

$$C \setminus D \equiv C \cap D^c \tag{1.35}$$

Definition 1.1.19 (Double Complement Law).

$$\left(D^{c}\right)^{c} \equiv D \tag{1.36}$$

Definition 1.1.20 (Contraposition).

$$C \subseteq D \Leftrightarrow D^c \subseteq C^c \tag{1.37}$$

$$C = D \Leftrightarrow C^c = D^c \tag{1.38}$$

Definition 1.1.21 (Arbitrary Union). Given sets A_1, A_2, \dots, A_n where $I = \{1, 2, \dots, n\}$

$$A_1 \cup A_2 \cup \dots \cup A_n \equiv \bigcup_{i \in I} A_i \tag{1.39}$$

Then

$$x \in \bigcup_{i \in I} A_i \Leftrightarrow \exists i \in I \colon x \in A_i \tag{1.40}$$

Definition 1.1.22 (Arbitrary Intersection). Given sets A_1, A_2, \dots, A_n where $I = \{1, 2, \dots, n\}$

$$A_1\cap A_2\cap \cdots \cap A_n \equiv \bigcap_{i\in I} A_i \tag{1.41}$$

Then

$$x \in \bigcap_{i \in I} A_i \Leftrightarrow \forall \, i \in I \colon x \in A_i \tag{1.42}$$

1.2 Functions

Definition 1.2.1 (Function). A function f is a mapping from X to Y

$$f \colon X \mapsto Y \tag{1.43}$$

- domain(f) = X
- image(f) = f(X)

Definition 1.2.2 (Total Function). A function is *total* if

$$domain(f) = X \tag{1.44}$$

Definition 1.2.3 (Partial Function). A function is partial if

$$domain(f) \subseteq X \tag{1.45}$$

Definition 1.2.4 (Surjection). A function $f: X \mapsto Y$ is *surjective* iff

$$f(X) = Y \Leftrightarrow \forall y \in Y \colon \exists x \in X \colon f(x) = y \tag{1.46}$$

Namely each $y \in Y$ has a corresponding $x \in X$.

Definition 1.2.5 (Injection (Encodings, One-to-one)). A function $f: X \mapsto Y$ is *injective* iff

$$\forall x_1, x_2 \in X \colon x_1 \neq x_2 \to f(x_1) \neq f(x_2) \tag{1.47}$$

$$\Leftrightarrow \forall \, x_1, x_2 \in X \colon f(x_1) = f(x_2) \to x_1 = x_2 \tag{1.48}$$

Namely each distinct element $x \in X$ maps to a different element in Y.

Definition 1.2.6 (Bijection). A function $f: X \mapsto Y$ is *bijective* iff f is both *injective* and *surjective*.

$$Bijective(f) \equiv Injective(f) \land Surjective(f)$$
 (1.49)

The *inverse bijection* $f^{-1}: Y \mapsto X$ does exist.

1.2.1 Composition of Injections

Proposition 1.2.1 (Composition of Injection). Given injections $f: X \mapsto Y$ and $g: Y \mapsto Z$, then their *composition* $h: X \mapsto Z$ is given by

$$h(x) = g(f(x)) \tag{1.50}$$

Namely $h = g \circ f$ where h is a function composed from g and f with f applied first.

Proof. Given any $x_1, x_2 \in X$ where $x_1 \neq x_2$, then

$$f(x_1) \neq f(x_2) \tag{1.51}$$

as *f* is *injective*, and thus

$$h(x_1) = g(f(x_1)) \neq g(f(x_2)) = h(x_2)$$
(1.52)

h is *injective* consequently.

Bibliography

[1] Max Kanovich and Robin Hirsch. "Lecture Notes on Discrete Mathematics for Computer Scientists". URL: http://www.cs.ucl.ac.uk/1819/a4u/t2/comp0147_discrete_mathematics_for_computer_scientists/.