

1. Dragonfly Data Science, Wellington, New Zealand
2. Centre for Ocean Life, National Institute of Aquatic Resources, Technical University of Denmark, Denmark

Introduction

Size-at-maturation and maximum size are key traits, defining ecological niches and driving ecological interactions¹. Temperature has been identified in various contexts as a key driver of organism size (Bergman's rule, Temperature-size-rule...) ². Climate change may therefore have strong effects on organism size and follow on effects at population and ecosystem levels, but no clear framework to understand effects of temperature ³.

Aim: Develop a eco-physiological model⁴ to predict effects of changing temperatures on organism size over ecological and evolutionary time-scales

Model

We set up a weight (w) based model consisting of mass and oxygen balance equations: Feeding - type 2 functional response with parameters for maximum consumption rate γ , modulated by feeding activity τ , and maximum intake h , which responds to temperature via T_M^c :

$$f = \frac{\tau \gamma w^p \Theta}{\tau \gamma w^p \Theta + T_h^c h w^q}$$

Mass balance: Intake is assimilated with loss from heat increment ϕ and absorption β . Metabolic loss due to standard and feeding activity (τ) metabolic costs, modulated by temperature via T_M^c

$$P = (1 - \beta - \phi) f T_h^c h w^q - k(1 + \tau \delta) T_M^c w^n$$

Oxygen balance: Oxygen sets the limit for metabolism via the cost of metabolising one gram of feed ω relative to the potential oxygen supply $f_{O_2} w^n$:

$$P_{O_2} = f_{O_2} w^n - \omega(\beta f T_h^c h w^q + k(1 + \tau \delta) T_M^c w^n)$$

Mortality: Weight based with base mortality m_0 and activity related mortality m_τ .

$$M = (m_0 + \tau m_\tau) w^{1-p}$$

τ is optimised at any temperature via Gilliam's rule ($\text{argmax}_\tau \{P/M\}$)

Strategy trade-off: minimising mortality M or maximising energy gain P

P strategy: Maximise intake at the cost of higher standard metabolism to maintain feeding activity and larger gut. Live with higher M .

M strategy: Minimise M by minimising dangerous feeding activity, minimise resting metabolism.

Temperature impact on size




Figure 3: **On ecological time-scales**, temperature affects size via evolved maturation-reaction norms⁵. Most reaction norms found in nature have a negative slope, meaning for faster growth, maturation occurs at a larger size. Increasing temperature (left panel, purple to yellow growth curves) modifies growth, and maturation age changes according to the reaction norm (right panel).




Figure 4: **On evolutionary time-scales**, selection moves the reaction norm. Size-at-maturation reacts strongly to temperature changes in the model (left panel). Selection (S) induced changes in size are slow for M strategists, fast for P strategists.

Ecological dynamics




Figure 1: With increasing temperature, most organisms will increase activity τ to offset increased metabolic costs. Oxygen sets metabolic limits at extreme temperatures.




Figure 2: Feeding level, available energy and mortality are impacted by increased activity and the metabolic response to temperature.

Acknowledgements & References

P. Neubauer acknowledges funding from the Royal Society of New Zealand Marsden Fast Start grant 14-DFG-001. All code for the model and this poster is accessible from github at repository Adaptive-dynamics/Sizing-fx-of-temperature.

1. Andersen, K. H. & Beyer, J. E. Asymptotic size determines species abundance in the marine size spectrum. *The American Naturalist* **168**, 54–61 (2006).
2. Atkinson, D. Temperature and organism size: A biological law for ectotherms? *Advances in ecological research* **25**, 1–1 (1994).
3. Lefevre, S., McKenzie, D. J. & Nilsson, G. E. Models projecting the fate of fish populations under climate change need to be based on valid physiological mechanisms. *Glob Change Biol* (2017).
4. Holt, R. E. & Jorgensen, C. Climate warming causes life-history evolution in a model for atlantic cod (*gadus morhua*). *Conservation Physiology* **2**, (2014).
5. Marty, L., Dieckmann, U., Rochet, M.-J. & Ernande, B. Impact of environmental covariation in growth and mortality on evolving maturation reaction norms. *The American Naturalist* **177**, E98–E118 (2011).

Take home messages

- On ecological time-scales changes in adult size due to temperature alone will likely be small, and masked by changes in the environment.
- On evolutionary time-scales, size responds strongly. The relative selection response is strongest at temperature extremes.
- Size of P strategy species with high activity reacts more strongly compared to sluggish M strategy species.