

Assignment 12

Adarsh Srivastava

The link to the solution is

<https://github.com/Adarsh1310/EE5609>

Abstract—This documents solves a problem based on Row Echelon form.

1 PROBLEM

Suppose \mathbf{R} and \mathbf{R}' are 2×3 row-reduced echelon matrices and that the system $\mathbf{RX}=0$ and $\mathbf{R}'\mathbf{X}=0$ have exactly the same solutions. Prove that $\mathbf{R}=\mathbf{R}'$.

2 SOLUTION

Since \mathbf{R} and \mathbf{R}' are 2×3 row-reduced echelon matrices they can be of following three types:-

- 1) Suppose matrix \mathbf{R} has one non-zero row then $\mathbf{RX}=0$ will have two free variables. Since $\mathbf{R}'\mathbf{X}=0$ will have the exact same solution as $\mathbf{RX}=0$, $\mathbf{R}'\mathbf{X}=0$ will also have two free variables. Thus \mathbf{R}' have one non zero row. Now let's consider a matrix \mathbf{A} with the first row as the non-zero row \mathbf{R} and second row as the second row of \mathbf{R}' .

$$\mathbf{R} = \begin{pmatrix} 1 & a & b \\ 0 & 0 & 0 \end{pmatrix} \quad (2.0.1)$$

$$\mathbf{R}' = \begin{pmatrix} 1 & c & d \\ 0 & 0 & 0 \end{pmatrix} \quad (2.0.2)$$

$$\mathbf{A} = \mathbf{R} + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{R}' \quad (2.0.3)$$

$$\mathbf{A} = \begin{pmatrix} 1 & a & b \\ 1 & c & d \end{pmatrix} \quad (2.0.4)$$

Let \mathbf{X} satisfy

$$\mathbf{RX} = 0 \quad (2.0.5)$$

$$\mathbf{R}'\mathbf{X} = 0 \quad (2.0.6)$$

Now multiplying \mathbf{A} and \mathbf{X} and substituting (2.0.3).

$$= \mathbf{AX} \quad (2.0.7)$$

$$= \left(\mathbf{R} + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{R}' \right) \mathbf{X} \quad (2.0.8)$$

$$= \mathbf{RX} + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{R}'\mathbf{X} \quad (2.0.9)$$

From (2.0.5) and (2.0.6)

$$\mathbf{AX} = 0 \quad (2.0.10)$$

Now, considering the augmented matrix,

$$\left(\begin{array}{ccc|c} 1 & a & b & 0 \\ 1 & c & d & 0 \end{array} \right) \quad (2.0.11)$$

$$\xleftrightarrow{R_2=R_2-R_1} \left(\begin{array}{ccc|c} 1 & a & b & 0 \\ 0 & c-a & d-b & 0 \end{array} \right) \quad (2.0.12)$$

For (2.0.12) to be consistent,

$$c - a = 0 \quad (2.0.13)$$

$$c = a \quad (2.0.14)$$

$$d - b = 0 \quad (2.0.15)$$

$$d = b \quad (2.0.16)$$

Hence, $\mathbf{R}=\mathbf{R}'$

- 2) Let \mathbf{R} and \mathbf{R}' have all rows as non zero.

$$\mathbf{R} = \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \end{pmatrix} \quad (2.0.17)$$

$$\mathbf{R}' = \begin{pmatrix} 1 & d & e \\ 0 & 1 & f \end{pmatrix} \quad (2.0.18)$$

Now let's consider another matrix \mathbf{A} whose first two rows are from \mathbf{R} and last two rows

are from \mathbf{R}'

$$\mathbf{A} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{R} + \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{R}' \quad (2.0.19)$$

$$\mathbf{A} = \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 1 & d & e \\ 0 & 1 & f \end{pmatrix} \quad (2.0.20)$$

Let \mathbf{X} satisfy

$$\mathbf{R}\mathbf{X} = 0 \quad (2.0.21)$$

$$\mathbf{R}'\mathbf{X} = 0 \quad (2.0.22)$$

Now multiplying \mathbf{A} and \mathbf{X} and substituting (2.0.19).

$$\mathbf{A}\mathbf{X} \quad (2.0.23)$$

$$\left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{R} + \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{R}' \right) \mathbf{X} \quad (2.0.24)$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{R}\mathbf{X} + \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{R}'\mathbf{X} \quad (2.0.25)$$

From (2.0.21) and (2.0.22)

$$\mathbf{A}\mathbf{X} = 0 \quad (2.0.26)$$

Now, considering the augmented matrix,

$$\left(\begin{array}{ccc|c} 1 & a & b & 0 \\ 0 & 1 & c & 0 \\ 1 & d & e & 0 \\ 0 & 1 & f & 0 \end{array} \right) \quad (2.0.27)$$

$$\xleftrightarrow{R_3=R_3-R_1} \left(\begin{array}{ccc|c} 1 & a & b & 0 \\ 0 & 1 & c & 0 \\ 1 & d-a & e-b & 0 \\ 0 & 1 & f-c & 0 \end{array} \right) \quad (2.0.28)$$

$$\xleftrightarrow{R_4=R_4-R_2} \left(\begin{array}{ccc|c} 1 & a & b & 0 \\ 0 & 1 & c & 0 \\ 0 & d-a & e-b & 0 \\ 0 & 0 & f-c & 0 \end{array} \right) \quad (2.0.29)$$

For (2.0.29) to be consistent,

$$d - a = 0 \quad (2.0.30)$$

$$d = a \quad (2.0.31)$$

$$e - b = 0 \quad (2.0.32)$$

$$e = b \quad (2.0.33)$$

$$f - c = 0 \quad (2.0.34)$$

$$f = c \quad (2.0.35)$$

Hence, $\mathbf{R}=\mathbf{R}'$

- 3) Suppose matrix \mathbf{R} have all the rows as zero then $\mathbf{R}\mathbf{X}=0$ will be satisfied for all values of \mathbf{X} . We know that $\mathbf{R}'\mathbf{X}=0$ will have the exact same solution as $\mathbf{R}\mathbf{X}=0$ then we can say that for all values of $\mathbf{X}=0$ equation $\mathbf{R}'\mathbf{X}=0$ will be satisfied Hence, $\mathbf{R}'=\mathbf{R}=0$.