

Assignment 19

Adarsh Srivastava

The link to the solution is

<https://github.com/Adarsh1310/EE5609>

Abstract—This documents solves a problem based on Linear Transformation.

1 PROBLEM

Let T be the linear transformation from \mathbb{R}^3 into \mathbb{R}^2 defined by,

$$T \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} x_1 + x_2 \\ 2x_3 - x_1 \end{pmatrix}$$

If β is the standard ordered basis for \mathbb{R}^3 and β' is the standard ordered basis for \mathbb{R}^2 , what is the matrix of T relative to the pair β, β'

2 SOLUTION

We know that,

$$[T\alpha]_{\beta} = \mathbf{A}[\alpha]_{\beta'} \quad (2.0.1)$$

where \mathbf{A} is called the matrix of T relative to ordered basis β, β' Using the ordered basis,

$$\beta = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (2.0.2)$$

$$\beta' = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (2.0.3)$$

Let,

$$\mathbf{V}_j = [T\alpha_j]_{\beta} \quad (2.0.4)$$

$$\mathbf{V} = (\mathbf{V}_1 \quad \mathbf{V}_2 \quad \mathbf{V}_3) \quad (2.0.5)$$

Using the given transformation,

$$T \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \end{pmatrix} \quad (2.0.6)$$

$$T \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (2.0.7)$$

$$T \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 2 \end{pmatrix} \quad (2.0.8)$$

So, $T(\beta)$ can be written as,

$$\mathbf{V} = \begin{pmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \end{pmatrix} \quad (2.0.9)$$

Now, since the transformation has to be found relative to the pair β, β' we can rewrite the equation in the question as,

$$\left(\begin{array}{cc|cc} 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & -1 & 0 & 2 \end{array} \right) \quad (2.0.10)$$

but from here we can see that β' is already an identity matrix and hence no row reduction is required. So by using (2.0.1) we can say that,

$$[T\alpha]_{\beta} = \begin{pmatrix} 1 & 1 & 0 \\ -1 & 0 & 2 \end{pmatrix} [\alpha]_{\beta'} \quad (2.0.11)$$