

Assignment 16

Adarsh Srivastava

The link to the solution is

<https://github.com/Adarsh1310/EE5609>

Abstract—This documents solves a problem based on basis and dimensions.

1 PROBLEM

Let \mathbf{V} be a vector space over a subfield \mathbf{F} of complex numbers. Suppose α, β and γ are linearly independent vectors in \mathbf{V} . Prove that $(\alpha+\beta), (\beta+\gamma)$ and $(\gamma+\alpha)$ are linearly independent.

2 SOLUTION

Let α, β and γ be three $n \times 1$ dimensional vectors. We need to prove that,

$$\begin{pmatrix} \alpha + \beta & \beta + \gamma & \gamma + \alpha \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad (2.0.1)$$

will only have a trivial solution. The above equation can be written as

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}^T \begin{pmatrix} \alpha + \beta & \beta + \gamma & \gamma + \alpha \end{pmatrix}^T = 0 \quad (2.0.2)$$

$$\begin{pmatrix} x_1 & x_2 & x_3 \end{pmatrix} \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} \alpha^T \\ \beta^T \\ \gamma^T \end{pmatrix} = 0 \quad (2.0.3)$$

$$\begin{pmatrix} x_1 + x_2 & x_1 + x_3 & x_2 + x_3 \end{pmatrix} \begin{pmatrix} \alpha^T \\ \beta^T \\ \gamma^T \end{pmatrix} = 0 \quad (2.0.4)$$

Since, α, β and γ are independent Hence (2.0.4) will have a trivial solution and so, $(\alpha+\beta), (\beta+\gamma)$ and $(\gamma+\alpha)$ are linearly independent.