

# Assignment 15

Adarsh Srivastava

The link to the solution is

<https://github.com/Adarsh1310/EE5609>

**Abstract**—This documents solves a problem based on vector subspaces.

## 1 PROBLEM

Let  $\mathbf{W}$  be the set of all  $(x_1, x_2, x_3, x_4, x_5)$  in  $\mathbb{R}^5$  which satisfy

$$2x_1 - x_2 + \frac{4}{3}x_3 - x_4 = 0$$

$$x_1 + \frac{2}{3}x_3 - x_5 = 0$$

$$9x_1 - 3x_2 + 6x_3 - 3x_4 - 3x_5 = 0$$

Find a finite set of vectors which spans  $\mathbf{W}$ .

## 2 SOLUTION

The given equations are,

$$2x_1 - x_2 + \frac{4}{3}x_3 - x_4 = 0 \quad (2.0.1)$$

$$x_1 + \frac{2}{3}x_3 - x_5 = 0 \quad (2.0.2)$$

$$9x_1 - 3x_2 + 6x_3 - 3x_4 - 3x_5 = 0 \quad (2.0.3)$$

which can be written as,

$$\begin{pmatrix} 2 & -1 & \frac{4}{3} & -1 & 0 \\ 1 & 0 & \frac{2}{3} & 0 & -1 \\ 9 & -3 & 6 & -3 & -3 \end{pmatrix} \mathbf{x} = 0 \quad (2.0.4)$$

Now, the augmented matrix,

$$\left( \begin{array}{ccccc|c} 2 & -1 & \frac{4}{3} & -1 & 0 & 0 \\ 1 & 0 & \frac{2}{3} & 0 & -1 & 0 \\ 9 & -3 & 6 & -3 & -3 & 0 \end{array} \right) \quad (2.0.5)$$

$$\xleftrightarrow{R_3=R_3-3R_1-3R_2} \left( \begin{array}{ccccc|c} 2 & -1 & \frac{4}{3} & -1 & 0 & 0 \\ 1 & 0 & \frac{2}{3} & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right) \quad (2.0.6)$$

$$\xleftrightarrow{R_2=R_2-\frac{1}{2}R_1} \left( \begin{array}{ccccc|c} 2 & -1 & \frac{4}{3} & -1 & 0 & 0 \\ 0 & \frac{1}{2} & 0 & \frac{1}{2} & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right) \quad (2.0.7)$$

$$\xleftrightarrow{R_2=2R_2} \left( \begin{array}{ccccc|c} 2 & -1 & \frac{4}{3} & -1 & 0 & 0 \\ 0 & 1 & 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right) \quad (2.0.8)$$

$$\xleftrightarrow{R_1=R_1+R_2} \left( \begin{array}{ccccc|c} 2 & 0 & \frac{4}{3} & 0 & -2 & 0 \\ 0 & 1 & 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right) \quad (2.0.9)$$

So,

$$2x_1 + \frac{4}{3}x_3 - 2x_5 = 0 \quad (2.0.10)$$

$$x_2 + x_4 - 2x_5 = 0 \quad (2.0.11)$$

Solving the equations we get,

$$x_1 = -\frac{2}{3}x_3 + x_5 \quad (2.0.12)$$

$$x_2 = -x_4 + 2x_5 \quad (2.0.13)$$

which can be written as,

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} \quad (2.0.14)$$

$$= \begin{pmatrix} -\frac{2}{3}x_3 + x_5 \\ -x_4 + 2x_5 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} \quad (2.0.15)$$

$$= x_3 \begin{pmatrix} -\frac{2}{3} \\ 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 0 \\ -1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + x_5 \begin{pmatrix} 1 \\ 2 \\ 0 \\ 0 \\ 1 \end{pmatrix} \quad (2.0.16)$$

where  $x_3, x_4$  and  $x_5 \in \mathbb{R}$ . Hence, the vectors

$\begin{pmatrix} -\frac{2}{3} \\ 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ -1 \\ 0 \\ 1 \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} 1 \\ 2 \\ 0 \\ 0 \\ 1 \end{pmatrix}$  will span  $\mathbf{W}$