

# Assignment 5

Adarsh Srivastava

The link to the solution is

<https://github.com/Adarsh1310/EE5609>

**Abstract**—This documents solves a problem based on circles.

## 1 PROBLEM

Find the area of the region bounded by the circle  $\mathbf{x}^T \mathbf{x} = 4$  and  $\left\| \mathbf{x} - \begin{pmatrix} 2 \\ 0 \end{pmatrix} \right\| = 2$ .

## 2 SOLUTION

General equation of circle is  $\mathbf{x}^T \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0$   
Taking equation of the first circle to be,

$$\|\mathbf{x}\|^2 + 2\mathbf{u}_1^T \mathbf{x} + f_1 = 0 \quad (2.0.1)$$

$$\mathbf{x}^T \mathbf{x} - 4 = 0 \quad (2.0.2)$$

$$\mathbf{u}_1 = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (2.0.3)$$

$$f_1 = -4 \quad (2.0.4)$$

$$\mathbf{O}_1 = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (2.0.5)$$

Taking equation of the second circle to be,

$$\left\| \mathbf{x} - \begin{pmatrix} 2 \\ 0 \end{pmatrix} \right\|^2 = 2^2 \quad (2.0.6)$$

$$\mathbf{x}^T \mathbf{x} + 2\mathbf{u}_2^T \mathbf{x} = 0 \quad (2.0.7)$$

$$\mathbf{u}_2 = \begin{pmatrix} -2 \\ 0 \end{pmatrix} \quad (2.0.8)$$

$$f_2 = 0 \quad (2.0.9)$$

$$\mathbf{O}_2 = \begin{pmatrix} 2 \\ 0 \end{pmatrix} \quad (2.0.10)$$

Now, Subtracting equation (2.0.7) from (2.0.2) We get,

$$\mathbf{x}^T \mathbf{x} - 2\mathbf{u}_2^T \mathbf{x} + f_1 - \mathbf{x}^T \mathbf{x} = 0 \quad (2.0.11)$$

$$2\mathbf{u}_2^T \mathbf{x} = -4 \quad (2.0.12)$$

$$\begin{pmatrix} -4 & 0 \end{pmatrix} \mathbf{x} = -4 \quad (2.0.13)$$

Which can be written as:-

$$\begin{pmatrix} 1 & 0 \end{pmatrix} \mathbf{x} = 1 \quad (2.0.14)$$

$$\mathbf{x} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (2.0.15)$$

$$\mathbf{x} = \mathbf{q} + \lambda \mathbf{m} \quad (2.0.16)$$

$$\mathbf{q} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (2.0.17)$$

$$\mathbf{m} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (2.0.18)$$

Substituting (2.0.16) in (2.0.1)

$$\|\mathbf{x}\|^2 + 2\mathbf{u}_1^T \mathbf{x} + f_1 = 0 \quad (2.0.19)$$

$$\|\mathbf{q} + \lambda \mathbf{m}\|^2 + f_1 = 0 \quad (2.0.20)$$

$$(\mathbf{q} + \lambda \mathbf{m})^T (\mathbf{q} + \lambda \mathbf{m}) + f_1 = 0 \quad (2.0.21)$$

$$\mathbf{q}^T (\mathbf{q} + \lambda \mathbf{m}) + \lambda \mathbf{m}^T (\mathbf{q} + \lambda \mathbf{m}) + f_1 = 0 \quad (2.0.22)$$

$$\|\mathbf{q}\|^2 + \lambda \mathbf{q}^T \mathbf{m} + \lambda \mathbf{m}^T \mathbf{q} + \lambda^2 \|\mathbf{m}\|^2 + f_1 = 0 \quad (2.0.23)$$

$$\|\mathbf{q}\|^2 + 2\lambda \mathbf{q}^T \mathbf{m} + \lambda^2 \|\mathbf{m}\|^2 + f_1 = 0 \quad (2.0.24)$$

$$\lambda(\lambda \|\mathbf{m}\|^2 + 2\mathbf{q}^T \mathbf{m}) = -f_1 - \|\mathbf{q}\|^2 \quad (2.0.25)$$

$$\lambda^2 \|\mathbf{m}\|^2 = -f_1 - \|\mathbf{q}\|^2 \quad (2.0.26)$$

$$\lambda^2 = \frac{-f_1 - \|\mathbf{q}\|^2}{\|\mathbf{m}\|^2} \quad (2.0.27)$$

$$\lambda^2 = 3 \quad (2.0.28)$$

$$\lambda = +\sqrt{3}, -\sqrt{3} \quad (2.0.29)$$

Substituting the value of  $\lambda$  in (2.0.16)

$$\mathbf{x} = \mathbf{q} + \lambda \mathbf{m} \quad (2.0.30)$$

$$\mathbf{A} = \begin{pmatrix} 1 \\ \sqrt{3} \end{pmatrix} \quad (2.0.31)$$

$$\mathbf{B} = \begin{pmatrix} 1 \\ -\sqrt{3} \end{pmatrix} \quad (2.0.32)$$

Now finding the direction vector  $\mathbf{m}_{O_1A}$ ,  $\mathbf{m}_{O_1B}$ ,  $\mathbf{m}_{O_2A}$  and  $\mathbf{m}_{O_2B}$ .

$$\mathbf{m}_{O_1A} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} - \begin{pmatrix} 1 \\ \sqrt{3} \end{pmatrix} = \begin{pmatrix} -1 \\ -\sqrt{3} \end{pmatrix} \quad (2.0.33)$$

$$\mathbf{m}_{O_1B} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} - \begin{pmatrix} 1 \\ -\sqrt{3} \end{pmatrix} = \begin{pmatrix} -1 \\ \sqrt{3} \end{pmatrix} \quad (2.0.34)$$

$$\mathbf{m}_{O_2A} = \begin{pmatrix} 2 \\ 0 \end{pmatrix} - \begin{pmatrix} 1 \\ \sqrt{3} \end{pmatrix} = \begin{pmatrix} 1 \\ -\sqrt{3} \end{pmatrix} \quad (2.0.35)$$

$$\mathbf{m}_{O_2B} = \begin{pmatrix} 2 \\ 0 \end{pmatrix} - \begin{pmatrix} 1 \\ -\sqrt{3} \end{pmatrix} = \begin{pmatrix} 1 \\ \sqrt{3} \end{pmatrix} \quad (2.0.36)$$

Now finding the angle  $\angle O_1AB$ .

$$\mathbf{m}_{O_1A}^T \mathbf{m}_{O_1B} = \|\mathbf{m}_{O_1A}\| \|\mathbf{m}_{O_1B}\| \cos \theta_1 \quad (2.0.37)$$

$$\frac{\mathbf{m}_{O_1A}^T \mathbf{m}_{O_1B}}{\|\mathbf{m}_{O_1A}\| \|\mathbf{m}_{O_1B}\|} = \cos \theta_1 \quad (2.0.38)$$

$$\frac{-2}{4} = \cos \theta_1 \quad (2.0.39)$$

$$\frac{-1}{2} = \cos \theta_1 \quad (2.0.40)$$

$$\theta_1 = 120^\circ \quad (2.0.41)$$

Now finding the angle  $\angle O_2AB$ .

$$\mathbf{m}_{O_2A}^T \mathbf{m}_{O_2B} = \|\mathbf{m}_{O_2A}\| \|\mathbf{m}_{O_2B}\| \cos \theta_2 \quad (2.0.42)$$

$$\frac{\mathbf{m}_{O_2A}^T \mathbf{m}_{O_2B}}{\|\mathbf{m}_{O_2A}\| \|\mathbf{m}_{O_2B}\|} = \cos \theta_2 \quad (2.0.43)$$

$$\frac{-2}{4} = \cos \theta_2 \quad (2.0.44)$$

$$\frac{-1}{2} = \cos \theta_2 \quad (2.0.45)$$

$$\theta_2 = 120^\circ \quad (2.0.46)$$

Finding area of  $\mathbf{O_1AB}$  and  $\mathbf{O_2AB}$ .

$$A_{O_1AB} = \frac{\theta_1}{360} r^2 - \frac{1}{2} 2 \sqrt{3} \quad (2.0.47)$$

$$= \frac{120}{360} 4\pi - \frac{1}{2} 2 \sqrt{3} \quad (2.0.48)$$

$$A_{O_2AB} = \frac{\pi \theta_2}{360} r^2 - \frac{1}{2} 2 \sqrt{3} \quad (2.0.49)$$

$$= \frac{120}{360} 4\pi - \frac{1}{2} 2 \sqrt{3} \quad (2.0.50)$$

Area of  $\mathbf{O_1AO_2B}$

$$A_{O_1AO_2B} = \frac{120}{360} 4\pi - \frac{1}{2} 2 \sqrt{3} + \frac{120}{360} 4\pi - \frac{1}{2} 2 \sqrt{3} \quad (2.0.51)$$

$$= \frac{8\pi}{3} - 2 \sqrt{3} \quad (2.0.52)$$




Fig. 0: Figure depicting intersection points of circle