

# Assignment 1

Addagalla Satyanarayana

**Abstract**—This document explains how to find a line perpendicular to 2 lines and passing through a point.

Download the python code from

<https://github.com/AddagallaSatyanarayana/AI5006/tree/master/Assignment1>

and latex-tikz codes from

<https://github.com/AddagallaSatyanarayana/AI5006/tree/master/Assignment1/Assignment1.tex>

Let the matrix  $\mathbf{M}$  be

$$\mathbf{M} = \begin{pmatrix} \mathbf{m}_1^T \\ \mathbf{m}_2^T \end{pmatrix} \quad (3.0.3)$$

$$\mathbf{m}_1 = \begin{pmatrix} 3 \\ -16 \\ 7 \end{pmatrix}, \mathbf{m}_2 = \begin{pmatrix} 3 \\ 8 \\ -5 \end{pmatrix}, \mathbf{n} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} \quad (3.0.4)$$

$$\mathbf{M}\mathbf{n} = 0 \quad (3.0.5)$$

The matrix form is

$$\begin{pmatrix} 3 & -16 & 7 \\ 3 & 8 & -5 \end{pmatrix} \quad (3.0.6)$$

$$\begin{pmatrix} 3 & -16 & 7 \\ 3 & 8 & -5 \end{pmatrix} \xrightarrow{R_2=R_1-R_2} \begin{pmatrix} 3 & -16 & 7 \\ 0 & -24 & 12 \end{pmatrix} \quad (3.0.7)$$

$$\begin{pmatrix} 3 & -16 & 7 \\ 0 & -24 & 12 \end{pmatrix} \xrightarrow{R_2=\frac{R_2}{-24}} \begin{pmatrix} 3 & -16 & 7 \\ 0 & -2 & 1 \end{pmatrix} \quad (3.0.8)$$

$$\begin{pmatrix} 3 & -16 & 7 \\ 0 & -2 & 1 \end{pmatrix} \xrightarrow{R_1=R_1-8R_2} \begin{pmatrix} 3 & 0 & -1 \\ 0 & -2 & 1 \end{pmatrix} \quad (3.0.9)$$

We have 2 equations and 3 unknowns, we will have parametric solution

$$n_1 = \frac{k}{3} \quad (3.0.10)$$

$$n_2 = \frac{k}{2} \quad (3.0.11)$$

$$n_3 = k \quad (3.0.12)$$

$$\mathbf{n} = \frac{k}{6} \begin{pmatrix} 2 \\ 3 \\ 6 \end{pmatrix} \quad (3.0.13)$$

The equation of required line from equation (2.0.1)

$$\begin{pmatrix} 1 \\ 2 \\ -4 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 3 \\ 6 \end{pmatrix} \quad (3.0.14)$$

## 1 PROBLEM

Find the vector equation of the line passing through the point  $\begin{pmatrix} 1 \\ 2 \\ -4 \end{pmatrix}$  and perpendicular to the two lines  $\frac{x-8}{3} = \frac{y+19}{-16} = \frac{z-10}{7}$  and  $\frac{x-15}{3} = \frac{y-29}{8} = \frac{z-5}{-5}$

## 2 EXPLANATION

Equation of a line passing through the point  $\mathbf{a}$  and parallel to the line  $\mathbf{n}$  is given by:

$$\mathbf{x} = \mathbf{a} + \lambda \mathbf{n} \quad (2.0.1)$$

where  $\lambda$  is some constant. Since the line passes

through  $\begin{pmatrix} 1 \\ 2 \\ -4 \end{pmatrix}$ ;  $\mathbf{a} = \begin{pmatrix} 1 \\ 2 \\ -4 \end{pmatrix}$

$$\mathbf{x} = \begin{pmatrix} 8 \\ -19 \\ 10 \end{pmatrix} + \lambda_1 \begin{pmatrix} 3 \\ -16 \\ 7 \end{pmatrix} \quad (2.0.2)$$

$$\mathbf{x} = \begin{pmatrix} 15 \\ 29 \\ 5 \end{pmatrix} + \lambda_2 \begin{pmatrix} 3 \\ 8 \\ -5 \end{pmatrix} \quad (2.0.3)$$

## 3 SOLUTION

Let  $\mathbf{n}$  be the normal vector to both lines. If  $\mathbf{m}_1$  and  $\mathbf{m}_2$  are the direction vectors of the lines, then

$$\mathbf{m}_1^T \mathbf{n} = 0 \quad (3.0.1)$$

$$\mathbf{m}_2^T \mathbf{n} = 0 \quad (3.0.2)$$




Fig. 0: Perpendicular Line