



POLITECNICO
MILANO 1863

All new slides at a glance

New Template Showcase

10.05.2026 | Jane Smith

Outline

- **Navigation**
- **Content Layouts**
- **Math and Academic**
- **Practical Slides**



Navigation

01

Outline

- **Navigation**
- Content Layouts
- Math and Academic
- Practical Slides

TOC

On the highlighted table of contents of the previous slide...

This doesn't make sense to me if we don't have the option to deactivate the Section slide (blue one)

In order to remove it just comment the line `\AtBeginSection{\frame{\sectionpage}}` in the preamble of the .tex file and and uncomment the line `\AtBeginSection{\tocslide[currentsection]}`



Content Layouts

02

Three Columns

Column A

First approach or concept goes here.

- Advantage one
- Advantage two

Column B

Second approach or concept for comparison.

- Advantage one
- Advantage two

Column C

Third approach or concept closes the comparison.

- Advantage one
- Advantage two

Image placeholder

Image placeholder

Deep-dive: System Architecture

“Science is the poetry of reality.”

Richard Dawkins


“Science is the poetry of reality.”

平常心是道



“Science is the poetry of reality.”

Richard Dawkins




平常心是道



“Science is the poetry of reality.”

Richard Dawkins



平常心是道



“Science is the poetry of reality.”

Richard Dawkins

平常心是道



“Science is the poetry of reality.”

Key Impact Metrics

78%

Reduction in inference latency after model distillation

123.4M

Parameters in the compressed architecture

567 mln

Total training tokens across all fine-tuning stages



Math and Academic

03

Convergence Analysis

Theorem (Banach Fixed-Point)

Let (X, d) be a non-empty complete metric space and $T : X \rightarrow X$ a contraction mapping with constant $q \in [0, 1)$. Then T has a unique fixed point $x^* \in X$, and for any $x_0 \in X$ the sequence $x_{n+1} = T(x_n)$ converges to x^* .

Key Definition

Definition (Lipschitz Continuity)

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is Lipschitz continuous with constant $L \geq 0$ if

$$\|f(x) - f(y)\| \leq L \|x - y\| \quad \forall x, y \in \mathbb{R}^n.$$

Convergence Analysis

Theorem (Banach Fixed-Point)

Let (X, d) be a non-empty complete metric space and $T : X \rightarrow X$ a contraction mapping with constant $q \in [0, 1)$. Then T has a unique fixed point $x^* \in X$, and for any $x_0 \in X$ the sequence $x_{n+1} = T(x_n)$ converges to x^* .

You can also comment here on the implications of the theorem, or sketch the proof idea in a second block.

Key Definition

Definition (Lipschitz Continuity)

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is Lipschitz continuous with constant $L \geq 0$ if

$$\|f(x) - f(y)\| \leq L \|x - y\| \quad \forall x, y \in \mathbb{R}^n.$$

Theorem (Banach Fixed-Point)

Let (X, d) be a non-empty complete metric space and $T : X \rightarrow X$ a contraction mapping with constant $q \in [0, 1)$. Then T has a unique fixed point $x^* \in X$, and for any $x_0 \in X$ the sequence $x_{n+1} = T(x_n)$ converges to x^* .

Gradient Descent

A first-order iterative optimisation algorithm for finding a local minimum of a differentiable function.

Complexity:

$\mathcal{O}(n \cdot k)$ per epoch, where n is the number of parameters and k the number of samples.

Input: $f, \nabla f, \alpha, x_0, T$

Output: approximate minimiser x^*

$t \leftarrow 0$

while $t < T$ **do**

$g \leftarrow \nabla f(x_t)$


$x_{t+1} \leftarrow x_t - \alpha \cdot g$

$t \leftarrow t + 1$

end while

return x_t

Project Roadmap





Practical Slides


04



Prof. Jane Smith

Email jane.smith@polimi.it

Institution Politecnico di Milano — DEIB






Thank you for your attention




Backup / Extra Slides



An appendix

A



Extra Results

B

Supplementary: Additional Ablation

This frame is part of the backup section. It is not counted in the main slide total because `\backupmarker` was inserted beforehand and this frame uses `noframenumbering`.

Tip

Apply `[noframenumbering]` to every backup frame so the header bar counter stays frozen at the last main-talk slide number.