

## 1.5 EXERCISES

In Exercises 1–4, determine if the system has a nontrivial solution. Try to use as few row operations as possible.

1.  $2x_1 - 5x_2 + 8x_3 = 0$   
 $-2x_1 - 7x_2 + x_3 = 0$   
 $4x_1 + 2x_2 + 7x_3 = 0$
2.  $x_1 - 2x_2 + 3x_3 = 0$   
 $-2x_1 - 3x_2 - 4x_3 = 0$   
 $2x_1 - 4x_2 + 9x_3 = 0$
3.  $-3x_1 + 4x_2 - 8x_3 = 0$   
 $-2x_1 + 5x_2 + 4x_3 = 0$
4.  $5x_1 - 3x_2 + 2x_3 = 0$   
 $-3x_1 - 4x_2 + 2x_3 = 0$

In Exercises 5 and 6, follow the method of Examples 1 and 2 to write the solution set of the given homogeneous system in parametric vector form.

5.  $2x_1 + 2x_2 + 4x_3 = 0$   
 $-4x_1 - 4x_2 - 8x_3 = 0$   
 $-3x_2 - 3x_3 = 0$
6.  $x_1 + 2x_2 - 3x_3 = 0$   
 $2x_1 + x_2 - 3x_3 = 0$   
 $-1x_1 + x_2 = 0$

In Exercises 7–12, describe all solutions of  $A\mathbf{x} = \mathbf{0}$  in parametric vector form, where  $A$  is row equivalent to the given matrix.

7.  $\begin{bmatrix} 1 & 3 & -3 & 7 \\ 0 & 1 & -4 & 5 \end{bmatrix}$
8.  $\begin{bmatrix} 1 & -3 & -8 & 5 \\ 0 & 1 & 2 & -4 \end{bmatrix}$
9.  $\begin{bmatrix} 3 & -6 & 6 \\ -2 & 4 & -2 \end{bmatrix}$
10.  $\begin{bmatrix} -1 & -4 & 0 & -4 \\ 2 & -8 & 0 & 8 \end{bmatrix}$
11.  $\begin{bmatrix} 1 & -4 & -2 & 0 & 3 & -5 \\ 0 & 0 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & -4 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$
12.  $\begin{bmatrix} 1 & -2 & 3 & -6 & 5 & 0 \\ 0 & 0 & 0 & 1 & 4 & -6 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$

13. Suppose the solution set of a certain system of linear equations can be described as  $x_1 = 5 + 4x_3$ ,  $x_2 = -2 - 7x_3$ , with  $x_3$  free. Use vectors to describe this set as a line in  $\mathbb{R}^3$ .
14. Suppose the solution set of a certain system of linear equations can be described as  $x_1 = 5x_4$ ,  $x_2 = 3 - 2x_4$ ,  $x_3 = 2 + 5x_4$ , with  $x_4$  free. Use vectors to describe this set as a “line” in  $\mathbb{R}^4$ .
15. Describe and compare the solution sets of  $x_1 + 5x_2 - 3x_3 = 0$  and  $x_1 + 5x_2 - 3x_3 = -2$ .
16. Describe and compare the solution sets of  $x_1 - 2x_2 + 3x_3 = 0$  and  $x_1 - 2x_2 + 3x_3 = 4$ .
17. Follow the method of Example 3 to describe the solutions of the following system in parametric vector form. Also, give a geometric description of the solution set and compare it to that in Exercise 5.

$$\begin{aligned} 2x_1 + 2x_2 + 4x_3 &= 8 \\ -4x_1 - 4x_2 - 8x_3 &= -16 \\ -3x_2 - 3x_3 &= 12 \end{aligned}$$

18. As in Exercise 17, describe the solutions of the following system in parametric vector form, and provide a geometric comparison with the solution set in Exercise 6.


$$\begin{aligned} x_1 + 2x_2 - 3x_3 &= 5 \\ 2x_1 + x_2 - 3x_3 &= 13 \\ -x_1 + x_2 &= -8 \end{aligned}$$

In Exercises 19 and 20, find the parametric equation of the line through  $\mathbf{a}$  parallel to  $\mathbf{b}$ .

$$19. \mathbf{a} = \begin{bmatrix} -2 \\ 0 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} -5 \\ 3 \end{bmatrix} \quad 20. \mathbf{a} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} -7 \\ 6 \end{bmatrix}$$

In Exercises 21 and 22, find a parametric equation of the line  $M$  through  $\mathbf{p}$  and  $\mathbf{q}$ . [Hint:  $M$  is parallel to the vector  $\mathbf{q} - \mathbf{p}$ . See the figure below.]

$$21. \mathbf{p} = \begin{bmatrix} 3 \\ -3 \end{bmatrix}, \mathbf{q} = \begin{bmatrix} 4 \\ 1 \end{bmatrix} \quad 22. \mathbf{p} = \begin{bmatrix} -3 \\ 2 \end{bmatrix}, \mathbf{q} = \begin{bmatrix} 0 \\ -3 \end{bmatrix}$$



In Exercises 23 and 24, mark each statement True or False. Justify each answer.

23. a. A homogeneous equation is always consistent.  
 b. The equation  $A\mathbf{x} = \mathbf{0}$  gives an explicit description of its solution set.  
 c. The homogeneous equation  $A\mathbf{x} = \mathbf{0}$  has the trivial solution if and only if the equation has at least one free variable.  
 d. The equation  $\mathbf{x} = \mathbf{p} + t\mathbf{v}$  describes a line through  $\mathbf{v}$  parallel to  $\mathbf{p}$ .  
 e. The solution set of  $A\mathbf{x} = \mathbf{b}$  is the set of all vectors of the form  $\mathbf{w} = \mathbf{p} + \mathbf{v}_h$ , where  $\mathbf{v}_h$  is any solution of the equation  $A\mathbf{x} = \mathbf{0}$ .
24. a. A homogeneous system of equations can be inconsistent.  
 b. If  $\mathbf{x}$  is a nontrivial solution of  $A\mathbf{x} = \mathbf{0}$ , then every entry in  $\mathbf{x}$  is nonzero.  
 c. The effect of adding  $\mathbf{p}$  to a vector is to move the vector in a direction parallel to  $\mathbf{p}$ .  
 d. The equation  $A\mathbf{x} = \mathbf{b}$  is homogeneous if the zero vector is a solution.