

## 1.9 EXERCISES


1-13 odd, 17-27 odd

In Exercises 1–10, assume that  $T$  is a linear transformation. Find the standard matrix of  $T$ .

- $T: \mathbb{R}^2 \rightarrow \mathbb{R}^4$ ,  $T(\mathbf{e}_1) = (3, 1, 3, 1)$  and  $T(\mathbf{e}_2) = (-5, 2, 0, 0)$ , where  $\mathbf{e}_1 = (1, 0)$  and  $\mathbf{e}_2 = (0, 1)$ .
- $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$ ,  $T(\mathbf{e}_1) = (1, 3)$ ,  $T(\mathbf{e}_2) = (4, -7)$ , and  $T(\mathbf{e}_3) = (-5, 4)$ , where  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  are the columns of the  $3 \times 3$  identity matrix.
- $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  rotates points (about the origin) through  $3\pi/2$  radians (counterclockwise).
- $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  rotates points (about the origin) through  $-\pi/4$  radians (clockwise). [Hint:  $T(\mathbf{e}_1) = (1/\sqrt{2}, -1/\sqrt{2})$ .]
- $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a vertical shear transformation that maps  $\mathbf{e}_1$  into  $\mathbf{e}_1 - 2\mathbf{e}_2$  but leaves the vector  $\mathbf{e}_2$  unchanged.
- $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a horizontal shear transformation that leaves  $\mathbf{e}_1$  unchanged and maps  $\mathbf{e}_2$  into  $\mathbf{e}_2 + 3\mathbf{e}_1$ .
- $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  first rotates points through  $-3\pi/4$  radian (clockwise) and then reflects points through the horizontal  $x_1$ -axis. [Hint:  $T(\mathbf{e}_1) = (-1/\sqrt{2}, 1/\sqrt{2})$ .]
- $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  first reflects points through the horizontal  $x_1$ -axis and then reflects points through the line  $x_2 = x_1$ .
- $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  first performs a horizontal shear that transforms  $\mathbf{e}_2$  into  $\mathbf{e}_2 - 2\mathbf{e}_1$  (leaving  $\mathbf{e}_1$  unchanged) and then reflects points through the line  $x_2 = -x_1$ .
- $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  first reflects points through the vertical  $x_2$ -axis and then rotates points  $\pi/2$  radians.
- A linear transformation  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  first reflects points through the  $x_1$ -axis and then reflects points through the  $x_2$ -axis. Show that  $T$  can also be described as a linear transformation that rotates points about the origin. What is the angle of that rotation?


- Show that the transformation in Exercise 8 is merely a rotation about the origin. What is the angle of the rotation?

- Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be the linear transformation such that  $T(\mathbf{e}_1)$  and  $T(\mathbf{e}_2)$  are the vectors shown in the figure. Using the figure, sketch the vector  $T(2, 1)$ .



- Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation with standard matrix  $A = [\mathbf{a}_1 \ \mathbf{a}_2]$ , where  $\mathbf{a}_1$  and  $\mathbf{a}_2$  are shown in the figure. Using the figure, draw the image of  $\begin{bmatrix} -1 \\ 3 \end{bmatrix}$  under the

transformation  $T$ .



In Exercises 15 and 16, fill in the missing entries of the matrix, assuming that the equation holds for all values of the variables.

$$15. \begin{bmatrix} ? & ? & ? \\ ? & ? & ? \\ ? & ? & ? \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3x_1 - 2x_3 \\ 4x_1 \\ x_1 - x_2 + x_3 \end{bmatrix}$$

$$16. \begin{bmatrix} ? & ? \\ ? & ? \\ ? & ? \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 - x_2 \\ -2x_1 + x_2 \\ x_1 \end{bmatrix}$$

In Exercises 17–20, show that  $T$  is a linear transformation by finding a matrix that implements the mapping. Note that  $x_1, x_2, \dots$  are not vectors but are entries in vectors.

$$17. T(x_1, x_2, x_3, x_4) = (0, x_1 + x_2, x_2 + x_3, x_3 + x_4)$$

$$18. T(x_1, x_2) = (2x_2 - 3x_1, x_1 - 4x_2, 0, x_2)$$

$$19. T(x_1, x_2, x_3) = (x_1 - 5x_2 + 4x_3, x_2 - 6x_3)$$

$$20. T(x_1, x_2, x_3, x_4) = 2x_1 + 3x_3 - 4x_4 \quad (T: \mathbb{R}^4 \rightarrow \mathbb{R})$$

$$21. \text{ Let } T: \mathbb{R}^2 \rightarrow \mathbb{R}^2 \text{ be a linear transformation such that } T(x_1, x_2) = (x_1 + x_2, 4x_1 + 5x_2). \text{ Find } \mathbf{x} \text{ such that } T(\mathbf{x}) = (3, 8).$$

$$22. \text{ Let } T: \mathbb{R}^2 \rightarrow \mathbb{R}^3 \text{ be a linear transformation such that } T(x_1, x_2) = (x_1 - 2x_2, -x_1 + 3x_2, 3x_1 - 2x_2). \text{ Find } \mathbf{x} \text{ such that } T(\mathbf{x}) = (-1, 4, 9).$$

In Exercises 23 and 24, mark each statement True or False. Justify each answer.

- a. A linear transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  is completely determined by its effect on the columns of the  $n \times n$  identity matrix.

- b. If  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  rotates vectors about the origin through an angle  $\varphi$ , then  $T$  is a linear transformation.

- c. When two linear transformations are performed one after another, the combined effect may not always be a linear transformation.

- d. A mapping  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  is onto  $\mathbb{R}^m$  if every vector  $\mathbf{x}$  in  $\mathbb{R}^n$  maps onto some vector in  $\mathbb{R}^m$ .

- e. If  $A$  is a  $3 \times 2$  matrix, then the transformation  $\mathbf{x} \mapsto A\mathbf{x}$  cannot be one-to-one.

24. a. Not every linear transformation from  $\mathbb{R}^n$  to  $\mathbb{R}^m$  is a matrix transformation.

- b. The columns of the standard matrix for a linear transformation from  $\mathbb{R}^n$  to  $\mathbb{R}^m$  are the images of the columns of the  $n \times n$  identity matrix.



- c. The standard matrix of a linear transformation from  $\mathbb{R}^2$  to  $\mathbb{R}^2$  that reflects points through the horizontal axis, the vertical axis, or the origin has the form  $\begin{bmatrix} a & 0 \\ 0 & d \end{bmatrix}$ , where  $a$  and  $d$  are  $\pm 1$ .
- d. A mapping  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is one-to-one if each vector in  $\mathbb{R}^n$  maps onto a unique vector in  $\mathbb{R}^m$ .
- e. If  $A$  is a  $3 \times 2$  matrix, then the transformation  $\mathbf{x} \mapsto A\mathbf{x}$  cannot map  $\mathbb{R}^2$  onto  $\mathbb{R}^3$ .

In Exercises 25–28, determine if the specified linear transformation is (a) one-to-one and (b) onto. Justify each answer.

25. The transformation in Exercise 17
26. The transformation in Exercise 2
27. The transformation in Exercise 19
28. The transformation in Exercise 14

In Exercises 29 and 30, describe the possible echelon forms of the standard matrix for a linear transformation  $T$ . Use the notation of Example 1 in Section 1.2.

29.  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  is one-to-one.

30.  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  is onto.

31. Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation, with  $A$  its standard matrix. Complete the following statement to make it true: “ $T$  is one-to-one if and only if  $A$  has \_\_\_\_\_ pivot columns.” Explain why the statement is true. [Hint: Look in the exercises for Section 1.7.]

32. Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation, with  $A$  its standard matrix. Complete the following statement to make it true: “ $T$  maps  $\mathbb{R}^n$  onto  $\mathbb{R}^m$  if and only if  $A$  has \_\_\_\_\_ pivot columns.” Find some theorems that explain why the statement is true.

33. Verify the uniqueness of  $A$  in Theorem 10. Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation such that  $T(\mathbf{x}) = B\mathbf{x}$  for some

$m \times n$  matrix  $B$ . Show that if  $A$  is the standard matrix for  $T$ , then  $A = B$ . [Hint: Show that  $A$  and  $B$  have the same columns.]

34. Why is the question “Is the linear transformation  $T$  onto?” an existence question?

35. If a linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  maps  $\mathbb{R}^n$  onto  $\mathbb{R}^m$ , can you give a relation between  $m$  and  $n$ ? If  $T$  is one-to-one, what can you say about  $m$  and  $n$ ?

36. Let  $S : \mathbb{R}^p \rightarrow \mathbb{R}^n$  and  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be linear transformations. Show that the mapping  $\mathbf{x} \mapsto T(S(\mathbf{x}))$  is a linear transformation (from  $\mathbb{R}^p$  to  $\mathbb{R}^m$ ). [Hint: Compute  $T(S(c\mathbf{u} + d\mathbf{v}))$  for  $\mathbf{u}, \mathbf{v}$  in  $\mathbb{R}^p$  and scalars  $c$  and  $d$ . Justify each step of the computation, and explain why this computation gives the desired conclusion.]

[M] In Exercises 37–40, let  $T$  be the linear transformation whose standard matrix is given. In Exercises 37 and 38, decide if  $T$  is a one-to-one mapping. In Exercises 39 and 40, decide if  $T$  maps  $\mathbb{R}^5$  onto  $\mathbb{R}^5$ . Justify your answers.

37.  $\begin{bmatrix} -5 & 10 & -5 & 4 \\ 8 & 3 & -4 & 7 \\ 4 & -9 & 5 & -3 \\ -3 & -2 & 5 & 4 \end{bmatrix}$

38.  $\begin{bmatrix} 7 & 5 & 4 & -9 \\ 10 & 6 & 16 & -4 \\ 12 & 8 & 12 & 7 \\ -8 & -6 & -2 & 5 \end{bmatrix}$

39.  $\begin{bmatrix} 4 & -7 & 3 & 7 & 5 \\ 6 & -8 & 5 & 12 & -8 \\ -7 & 10 & -8 & -9 & 14 \\ 3 & -5 & 4 & 2 & -6 \\ -5 & 6 & -6 & -7 & 3 \end{bmatrix}$

40.  $\begin{bmatrix} 9 & 13 & 5 & 6 & -1 \\ 14 & 15 & -7 & -6 & 4 \\ -8 & -9 & 12 & -5 & -9 \\ -5 & -6 & -8 & 9 & 8 \\ 13 & 14 & 15 & 2 & 11 \end{bmatrix}$

### SOLUTION TO PRACTICE PROBLEMS

WEB

1. Follow what happens to  $\mathbf{e}_1$  and  $\mathbf{e}_2$ . See Figure 5. First,  $\mathbf{e}_1$  is unaffected by the shear and then is reflected into  $-\mathbf{e}_1$ . So  $T(\mathbf{e}_1) = -\mathbf{e}_1$ . Second,  $\mathbf{e}_2$  goes to  $\mathbf{e}_2 - .5\mathbf{e}_1$  by the shear transformation. Since reflection through the  $x_2$ -axis changes  $\mathbf{e}_1$  into  $-\mathbf{e}_1$  and




FIGURE 5 The composition of two transformations

## 4.2 EXERCISES

1-11 odd, 21, 23, 31

1. Determine if  $\mathbf{w} = \begin{bmatrix} 1 \\ 3 \\ -4 \end{bmatrix}$  is in  $\text{Nul } A$ , where

$$A = \begin{bmatrix} 3 & -5 & -3 \\ 6 & -2 & 0 \\ -8 & 4 & 1 \end{bmatrix}.$$

2. Determine if  $\mathbf{w} = \begin{bmatrix} 5 \\ -3 \\ 2 \end{bmatrix}$  is in  $\text{Nul } A$ , where

$$A = \begin{bmatrix} 5 & 21 & 19 \\ 13 & 23 & 2 \\ 8 & 14 & 1 \end{bmatrix}.$$

In Exercises 3–6, find an explicit description of  $\text{Nul } A$  by listing vectors that span the null space.

3.  $A = \begin{bmatrix} 1 & 3 & 5 & 0 \\ 0 & 1 & 4 & -2 \end{bmatrix}$

4.  $A = \begin{bmatrix} 1 & -6 & 4 & 0 \\ 0 & 0 & 2 & 0 \end{bmatrix}$

5.  $A = \begin{bmatrix} 1 & -2 & 0 & 4 & 0 \\ 0 & 0 & 1 & -9 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$

6.  $A = \begin{bmatrix} 1 & 5 & -4 & -3 & 1 \\ 0 & 1 & -2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$

In Exercises 7–14, either use an appropriate theorem to show that the given set,  $W$ , is a vector space, or find a specific example to the contrary.

7.  $\left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a + b + c = 2 \right\}$  8.  $\left\{ \begin{bmatrix} r \\ s \\ t \end{bmatrix} : 5r - 1 = s + 2t \right\}$

9.  $\left\{ \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} : \begin{array}{l} a - 2b = 4c \\ 2a = c + 3d \end{array} \right\}$  10.  $\left\{ \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} : \begin{array}{l} a + 3b = c \\ b + c + a = d \end{array} \right\}$

11.  $\left\{ \begin{bmatrix} b - 2d \\ 5 + d \\ b + 3d \\ d \end{bmatrix} : b, d \text{ real} \right\}$  12.  $\left\{ \begin{bmatrix} b - 5d \\ 2b \\ 2d + 1 \\ d \end{bmatrix} : b, d \text{ real} \right\}$

13.  $\left\{ \begin{bmatrix} c - 6d \\ d \\ c \end{bmatrix} : c, d \text{ real} \right\}$  14.  $\left\{ \begin{bmatrix} -a + 2b \\ a - 2b \\ 3a - 6b \end{bmatrix} : a, b \text{ real} \right\}$

In Exercises 15 and 16, find  $A$  such that the given set is  $\text{Col } A$ .

15.  $\left\{ \begin{bmatrix} 2s + 3t \\ r + s - 2t \\ 4r + s \\ 3r - s - t \end{bmatrix} : r, s, t \text{ real} \right\}$

16.  $\left\{ \begin{bmatrix} b - c \\ 2b + c + d \\ 5c - 4d \\ d \end{bmatrix} : b, c, d \text{ real} \right\}$

For the matrices in Exercises 17–20, (a) find  $k$  such that  $\text{Nul } A$  is a subspace of  $\mathbb{R}^k$ , and (b) find  $k$  such that  $\text{Col } A$  is a subspace of  $\mathbb{R}^k$ .

17.  $A = \begin{bmatrix} 2 & -6 \\ -1 & 3 \\ -4 & 12 \\ 3 & -9 \end{bmatrix}$

18.  $A = \begin{bmatrix} 7 & -2 & 0 \\ -2 & 0 & -5 \\ 0 & -5 & 7 \\ -5 & 7 & -2 \end{bmatrix}$

19.  $A = \begin{bmatrix} 4 & 5 & -2 & 6 & 0 \\ 1 & 1 & 0 & 1 & 0 \end{bmatrix}$

20.  $A = \begin{bmatrix} 1 & -3 & 9 & 0 & -5 \end{bmatrix}$

21. With  $A$  as in Exercise 17, find a nonzero vector in  $\text{Nul } A$  and a nonzero vector in  $\text{Col } A$ .

22. With  $A$  as in Exercise 3, find a nonzero vector in  $\text{Nul } A$  and a nonzero vector in  $\text{Col } A$ .

23. Let  $A = \begin{bmatrix} -6 & 12 \\ -3 & 6 \end{bmatrix}$  and  $\mathbf{w} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ . Determine if  $\mathbf{w}$  is in  $\text{Col } A$ . Is  $\mathbf{w}$  in  $\text{Nul } A$ ?

24. Let  $A = \begin{bmatrix} -8 & -2 & -9 \\ 6 & 4 & 8 \\ 4 & 0 & 4 \end{bmatrix}$  and  $\mathbf{w} = \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix}$ . Determine if  $\mathbf{w}$  is in  $\text{Col } A$ . Is  $\mathbf{w}$  in  $\text{Nul } A$ ?

In Exercises 25 and 26,  $A$  denotes an  $m \times n$  matrix. Mark each statement True or False. Justify each answer.

25. a. The null space of  $A$  is the solution set of the equation  $A\mathbf{x} = \mathbf{0}$ .

b. The null space of an  $m \times n$  matrix is in  $\mathbb{R}^m$ .

c. The column space of  $A$  is the range of the mapping  $\mathbf{x} \mapsto A\mathbf{x}$ .

d. If the equation  $A\mathbf{x} = \mathbf{b}$  is consistent, then  $\text{Col } A$  is  $\mathbb{R}^m$ .

e. The kernel of a linear transformation is a vector space.

f.  $\text{Col } A$  is the set of all vectors that can be written as  $A\mathbf{x}$  for some  $\mathbf{x}$ .

26. a. A null space is a vector space.

b. The column space of an  $m \times n$  matrix is in  $\mathbb{R}^m$ .

c.  $\text{Col } A$  is the set of all solutions of  $A\mathbf{x} = \mathbf{b}$ .

d.  $\text{Nul } A$  is the kernel of the mapping  $\mathbf{x} \mapsto A\mathbf{x}$ .

e. The range of a linear transformation is a vector space.

f. The set of all solutions of a homogeneous linear differential equation is the kernel of a linear transformation.

27. It can be shown that a solution of the system below is  $x_1 = 3$ ,  $x_2 = 2$ , and  $x_3 = -1$ . Use this fact and the theory from this section to explain why another solution is  $x_1 = 30$ ,  $x_2 = 20$ , and  $x_3 = -10$ . (Observe how the solutions are related, but make no other calculations.)

$$x_1 - 3x_2 - 3x_3 = 0$$

$$-2x_1 + 4x_2 + 2x_3 = 0$$

$$-x_1 + 5x_2 + 7x_3 = 0$$

28. Consider the following two systems of equations:

$$5x_1 + x_2 - 3x_3 = 0 \quad 5x_1 + x_2 - 3x_3 = 0$$

$$-9x_1 + 2x_2 + 5x_3 = 1 \quad -9x_1 + 2x_2 + 5x_3 = 5$$

$$4x_1 + x_2 - 6x_3 = 9 \quad 4x_1 + x_2 - 6x_3 = 45$$

It can be shown that the first system has a solution. Use this fact and the theory from this section to explain why the second system must also have a solution. (Make no row operations.)



29. Prove Theorem 3 as follows: Given an  $m \times n$  matrix  $A$ , an element in  $\text{Col } A$  has the form  $A\mathbf{x}$  for some  $\mathbf{x}$  in  $\mathbb{R}^n$ . Let  $A\mathbf{x}$  and  $A\mathbf{w}$  represent any two vectors in  $\text{Col } A$ .
- Explain why the zero vector is in  $\text{Col } A$ .
  - Show that the vector  $A\mathbf{x} + A\mathbf{w}$  is in  $\text{Col } A$ .
  - Given a scalar  $c$ , show that  $c(A\mathbf{x})$  is in  $\text{Col } A$ .
30. Let  $T : V \rightarrow W$  be a linear transformation from a vector space  $V$  into a vector space  $W$ . Prove that the range of  $T$  is a subspace of  $W$ . [Hint: Typical elements of the range have the form  $T(\mathbf{x})$  and  $T(\mathbf{w})$  for some  $\mathbf{x}, \mathbf{w}$  in  $V$ .]
31. Define  $T : \mathbb{P}_2 \rightarrow \mathbb{R}^2$  by  $T(\mathbf{p}) = \begin{bmatrix} \mathbf{p}(0) \\ \mathbf{p}(1) \end{bmatrix}$ . For instance, if  $\mathbf{p}(t) = 3 + 5t + 7t^2$ , then  $T(\mathbf{p}) = \begin{bmatrix} 3 \\ 15 \end{bmatrix}$ .
- Show that  $T$  is a linear transformation. [Hint: For arbitrary polynomials  $\mathbf{p}, \mathbf{q}$  in  $\mathbb{P}_2$ , compute  $T(\mathbf{p} + \mathbf{q})$  and  $T(c\mathbf{p})$ .]
  - Find a polynomial  $\mathbf{p}$  in  $\mathbb{P}_2$  that spans the kernel of  $T$ , and describe the range of  $T$ .
32. Define a linear transformation  $T : \mathbb{P}_2 \rightarrow \mathbb{R}^2$  by  $T(\mathbf{p}) = \begin{bmatrix} \mathbf{p}(0) \\ \mathbf{p}(0) \end{bmatrix}$ . Find polynomials  $\mathbf{p}_1$  and  $\mathbf{p}_2$  in  $\mathbb{P}_2$  that span the kernel of  $T$ , and describe the range of  $T$ .
33. Let  $M_{2 \times 2}$  be the vector space of all  $2 \times 2$  matrices, and define  $T : M_{2 \times 2} \rightarrow M_{2 \times 2}$  by  $T(A) = A + A^T$ , where  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ .
- Show that  $T$  is a linear transformation.
  - Let  $B$  be any element of  $M_{2 \times 2}$  such that  $B^T = B$ . Find an  $A$  in  $M_{2 \times 2}$  such that  $T(A) = B$ .
  - Show that the range of  $T$  is the set of  $B$  in  $M_{2 \times 2}$  with the property that  $B^T = B$ .
  - Describe the kernel of  $T$ .
34. (Calculus required) Define  $T : C[0, 1] \rightarrow C[0, 1]$  as follows: For  $\mathbf{f}$  in  $C[0, 1]$ , let  $T(\mathbf{f})$  be the antiderivative  $\mathbf{F}$  of  $\mathbf{f}$  such that  $\mathbf{F}(0) = 0$ . Show that  $T$  is a linear transformation, and describe the kernel of  $T$ . (See the notation in Exercise 20 of Section 4.1.)
35. Let  $V$  and  $W$  be vector spaces, and let  $T : V \rightarrow W$  be a linear transformation. Given a subspace  $U$  of  $V$ , let  $T(U)$  denote the set of all images of the form  $T(\mathbf{x})$ , where  $\mathbf{x}$  is in  $U$ . Show that  $T(U)$  is a subspace of  $W$ .
36. Given  $T : V \rightarrow W$  as in Exercise 35, and given a subspace  $Z$  of  $W$ , let  $U$  be the set of all  $\mathbf{x}$  in  $V$  such that  $T(\mathbf{x})$  is in  $Z$ . Show that  $U$  is a subspace of  $V$ .
37. [M] Determine whether  $\mathbf{w}$  is in the column space of  $A$ , the null space of  $A$ , or both, where
- $$\mathbf{w} = \begin{bmatrix} 1 \\ 1 \\ -1 \\ -3 \end{bmatrix}, \quad A = \begin{bmatrix} 7 & 6 & -4 & 1 \\ -5 & -1 & 0 & -2 \\ 9 & -11 & 7 & -3 \\ 19 & -9 & 7 & 1 \end{bmatrix}$$
38. [M] Determine whether  $\mathbf{w}$  is in the column space of  $A$ , the null space of  $A$ , or both, where
- $$\mathbf{w} = \begin{bmatrix} 1 \\ 2 \\ 1 \\ 0 \end{bmatrix}, \quad A = \begin{bmatrix} -8 & 5 & -2 & 0 \\ -5 & 2 & 1 & -2 \\ 10 & -8 & 6 & -3 \\ 3 & -2 & 1 & 0 \end{bmatrix}$$
39. [M] Let  $\mathbf{a}_1, \dots, \mathbf{a}_5$  denote the columns of the matrix  $A$ , where
- $$A = \begin{bmatrix} 5 & 1 & 2 & 2 & 0 \\ 3 & 3 & 2 & -1 & -12 \\ 8 & 4 & 4 & -5 & 12 \\ 2 & 1 & 1 & 0 & -2 \end{bmatrix}, \quad B = [\mathbf{a}_1 \quad \mathbf{a}_2 \quad \mathbf{a}_4]$$
- Explain why  $\mathbf{a}_3$  and  $\mathbf{a}_5$  are in the column space of  $B$ .
  - Find a set of vectors that spans  $\text{Nul } A$ .
  - Let  $T : \mathbb{R}^5 \rightarrow \mathbb{R}^4$  be defined by  $T(\mathbf{x}) = A\mathbf{x}$ . Explain why  $T$  is neither one-to-one nor onto.
40. [M] Let  $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$  and  $K = \text{Span}\{\mathbf{v}_3, \mathbf{v}_4\}$ , where
- $$\mathbf{v}_1 = \begin{bmatrix} 5 \\ 3 \\ 8 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 1 \\ 3 \\ 4 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix}, \mathbf{v}_4 = \begin{bmatrix} 0 \\ -12 \\ -28 \end{bmatrix}.$$
- Then  $H$  and  $K$  are subspaces of  $\mathbb{R}^3$ . In fact,  $H$  and  $K$  are planes in  $\mathbb{R}^3$  through the origin, and they intersect in a line through  $\mathbf{0}$ . Find a nonzero vector  $\mathbf{w}$  that generates that line. [Hint:  $\mathbf{w}$  can be written as  $c_1\mathbf{v}_1 + c_2\mathbf{v}_2$  and also as  $c_3\mathbf{v}_3 + c_4\mathbf{v}_4$ . To build  $\mathbf{w}$ , solve the equation  $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 = c_3\mathbf{v}_3 + c_4\mathbf{v}_4$  for the unknown  $c_j$ 's.]