

Introductory Mathematics II (MTS 101)

Trigonometry

Department of Mathematical Sciences
Federal University of Technology, Akure

1 Introduction

1.1 Definition

The word ‘trigonometry’ suggests ‘tri’-three, ‘gono’-angle, ‘metry’-measurement. As such, trigonometry is basically about triangles, most especially right-angled triangles.

1.2 Revision

VOTE KONNECT FOR 006

- Circular Measures
- Triangles
- Pythagoras’ Theorem

2 Trigonometric Ratios

From the right-angled triangle shown in Figure 1, we have the following trigonometric ratios:

$$\begin{aligned}\bullet \sin \theta &= \frac{a}{c} \\ \bullet \cos \theta &= \frac{b}{c} \\ \bullet \tan \theta &= \frac{a}{b}\end{aligned}$$

$$\begin{aligned}\bullet \csc \theta &= \frac{1}{\sin \theta} = \frac{c}{a} \\ \bullet \sec \theta &= \frac{1}{\cos \theta} = \frac{c}{b} \\ \bullet \cot \theta &= \frac{1}{\tan \theta} = \frac{b}{a}\end{aligned}$$




Figure 1: A Right-Angled Triangle

3 Trigonometric Identities

3.1 Basic Identities

$$\frac{\sin \theta}{\cos \theta} = \frac{\frac{a}{c}}{\frac{b}{c}} = \frac{a}{c} \times \frac{c}{b} = \frac{a}{b} = \tan \theta \quad (3.1)$$

$$\frac{\cos \theta}{\sin \theta} = \frac{\frac{b}{c}}{\frac{a}{c}} = \frac{b}{c} \times \frac{c}{a} = \frac{b}{a} = \cot \theta \quad (3.2)$$

With reference to Figure 1, using Pythagoras' Theorem,

$$a^2 + b^2 = c^2 \quad (3.3)$$

Dividing (3.3) through by c^2 , we have

$$\begin{aligned} \frac{a^2}{c^2} + \frac{b^2}{c^2} &= \frac{c^2}{c^2} \\ \frac{a^2}{c^2} + \frac{b^2}{c^2} &= 1 \\ \sin^2 \theta + \cos^2 \theta &= 1 \end{aligned} \quad (3.4)$$

Dividing (3.3) through by b^2 , we have

$$\begin{aligned} \frac{a^2}{b^2} + \frac{b^2}{b^2} &= \frac{c^2}{b^2} \\ \frac{a^2}{b^2} + 1 &= \frac{c^2}{b^2} \\ \tan^2 \theta + 1 &= \sec^2 \theta \end{aligned} \quad (3.5)$$

Dividing (3.3) through by a^2 , we have

$$\begin{aligned} \frac{a^2}{a^2} + \frac{b^2}{a^2} &= \frac{c^2}{a^2} \\ 1 + \frac{b^2}{a^2} &= \frac{c^2}{a^2} \\ 1 + \cot^2 \theta &= \csc^2 \theta \end{aligned} \quad (3.6)$$

3.2 Examples

3.2.1 Example 1

To show that $\sin^2 \theta + (1 + \cos \theta)^2 = 2(1 + \cos \theta)$,

$$\begin{aligned} \sin^2 \theta + (1 + \cos \theta)^2 &= \sin^2 \theta + 1 + 2 \cos \theta + \cos^2 \theta \\ &= \sin^2 \theta + \cos^2 \theta + 1 + 2 \cos \theta \\ &= 1 + 1 + 2 \cos \theta \quad (\text{Recall that } \sin^2 \theta + \cos^2 \theta = 1) \\ &= 2 + 2 \cos \theta \\ &= 2(1 + \cos \theta) \end{aligned}$$

$$\therefore \sin^2 \theta + (1 + \cos \theta)^2 = 2(1 + \cos \theta)$$

3.2.2 Example 2

To show that $\frac{1+\sin \theta}{1+\cos \theta} \cdot \frac{1+\sec \theta}{1+\csc \theta} = \tan \theta$,

$$\begin{aligned}
 \frac{1+\sin \theta}{1+\cos \theta} \cdot \frac{1+\sec \theta}{1+\csc \theta} &= \frac{1+\sin \theta}{1+\cos \theta} \cdot \frac{1+\frac{1}{\cos \theta}}{1+\frac{1}{\sin \theta}} \\
 &= \frac{1+\sin \theta}{1+\cos \theta} \cdot \frac{\frac{\cos \theta+1}{\cos \theta}}{\frac{\sin \theta+1}{\sin \theta}} \\
 &= \frac{1+\sin \theta}{1+\cos \theta} \cdot \frac{\cos \theta+1}{\cos \theta} \div \frac{\sin \theta+1}{\sin \theta} \\
 &= \frac{1+\sin \theta}{1+\cos \theta} \cdot \frac{\cos \theta+1}{\cos \theta} \cdot \frac{\sin \theta}{\sin \theta+1} \\
 &= \frac{\sin \theta}{\cos \theta} \\
 &= \tan \theta
 \end{aligned}$$

$$\therefore \frac{1+\sin \theta}{1+\cos \theta} \cdot \frac{1+\sec \theta}{1+\csc \theta} = \tan \theta$$

3.2.3 Example 3

To show that $\frac{1+\cos \theta}{1-\cos \theta} \cdot \frac{\sec \theta-1}{\sec \theta+1} = 1$,

$$\begin{aligned}
 \frac{1+\cos \theta}{1-\cos \theta} \cdot \frac{\sec \theta-1}{\sec \theta+1} &= \frac{1+\cos \theta}{1-\cos \theta} \cdot \frac{\frac{1}{\cos \theta}-1}{\frac{1}{\cos \theta}+1} \\
 &= \frac{1+\cos \theta}{1-\cos \theta} \cdot \frac{\frac{1-\cos \theta}{\cos \theta}}{\frac{1+\cos \theta}{\cos \theta}} \\
 &= \frac{1+\cos \theta}{1-\cos \theta} \cdot \frac{1-\cos \theta}{\cos \theta} \div \frac{1+\cos \theta}{\cos \theta} \\
 &= \frac{1+\cos \theta}{1-\cos \theta} \cdot \frac{1-\cos \theta}{\cos \theta} \cdot \frac{\cos \theta}{1+\cos \theta} \\
 &= \frac{\cos \theta}{\cos \theta} \\
 &= 1
 \end{aligned}$$

$$\therefore \frac{1+\cos \theta}{1-\cos \theta} \cdot \frac{\sec \theta-1}{\sec \theta+1} = 1$$

3.2.4 Example 4

To show that $(x^j)^2 + (y^j)^2 = x^2 + y^2$ if $x^j = x \cos \theta + y \sin \theta$ and $y^j = x \sin \theta - y \cos \theta$,

$$\begin{aligned}
 (x^j)^2 + (y^j)^2 &= (x \cos \theta + y \sin \theta)^2 + (x \sin \theta - y \cos \theta)^2 \\
 &= x^2 \cos^2 \theta + 2xy \sin \theta \cos \theta + y^2 \sin^2 \theta + x^2 \sin^2 \theta - 2xy \sin \theta \cos \theta + y^2 \cos^2 \theta \\
 &= x^2 \sin^2 \theta + x^2 \cos^2 \theta + y^2 \sin^2 \theta + y^2 \cos^2 \theta \\
 &= x^2(\sin^2 \theta + \cos^2 \theta) + y^2(\sin^2 \theta + \cos^2 \theta) \\
 &= x^2(1) + y^2(1) \quad (\text{Recall that } \sin^2 \theta + \cos^2 \theta = 1) \\
 &= x^2 + y^2
 \end{aligned}$$

$$\therefore (x^j)^2 + (y^j)^2 = x^2 + y^2 \text{ if } x^j = x \cos \theta + y \sin \theta \text{ and } y^j = x \sin \theta - y \cos \theta$$

3.2.5 Example 5

To show that $x^2 + y^2 + z^2 = c^2$ if $x = c \sin \theta \cos \varphi$, $y = c \sin \theta \sin \varphi$ and $z = c \cos \theta$,

$$\begin{aligned}
 x^2 + y^2 + z^2 &= (c \sin \theta \cos \varphi)^2 + (c \sin \theta \sin \varphi)^2 + (c \cos \theta)^2 \\
 &= c^2 \sin^2 \theta \cos^2 \varphi + c^2 \sin^2 \theta \sin^2 \varphi + c^2 \cos^2 \theta \\
 &= c^2 \sin^2 \theta (\cos^2 \varphi + \sin^2 \varphi) + c^2 \cos^2 \theta \\
 &= c^2 \sin^2 \theta (1) + c^2 \cos^2 \theta \\
 &= c^2 \sin^2 \theta + c^2 \cos^2 \theta \\
 &= c^2 (\sin^2 \theta + \cos^2 \theta) \\
 &= c^2 (1) \\
 &= c^2
 \end{aligned}$$

$\therefore x^2 + y^2 + z^2 = c^2$ if $x = c \sin \theta \cos \varphi$, $y = c \sin \theta \sin \varphi$ and $z = c \cos \theta$

4 The Addition Formulae

4.1 Basic Formulae

$$\sin(A + B) = \sin A \cos B + \cos A \sin B \quad (4.1)$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B \quad (4.2)$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B \quad (4.3)$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B \quad (4.4)$$

$$\begin{aligned}
 \tan(A + B) &= \frac{\sin(A + B)}{\cos(A + B)} \\
 &= \frac{\sin A \cos B + \cos A \sin B}{\cos A \cos B - \sin A \sin B} \\
 &= \frac{\frac{\sin A \cos B}{\cos A \cos B} + \frac{\cos A \sin B}{\cos A \cos B}}{\frac{\cos A \cos B}{\cos A \cos B} - \frac{\sin A \sin B}{\cos A \cos B}} \\
 &= \frac{\tan A + \tan B}{1 - \tan A \tan B}
 \end{aligned} \quad (4.5)$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B} \quad (4.6)$$

4.2 Examples

4.2.1 Example 1

To calculate $\sin 15^\circ$ given that $\sin 45^\circ = \cos 45^\circ = \frac{\sqrt{2}}{2}$, $\sin 30^\circ = \frac{1}{2}$ and $\cos 30^\circ = \frac{\sqrt{3}}{2}$,

$$\begin{aligned}
 \sin 15^\circ &= \sin(45^\circ - 30^\circ) \\
 &= \sin 45^\circ \cos 30^\circ - \cos 45^\circ \sin 30^\circ \\
 &= \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{3}}{2} - \frac{\sqrt{2}}{2} \cdot \frac{1}{2} \\
 &= \frac{\sqrt{2}(\sqrt{3} - 1)}{4}
 \end{aligned}$$

$$\begin{aligned}\sin 15^\circ &= \frac{\sqrt{3}}{2\sqrt{2}} - \frac{1}{2\sqrt{2}} \\ &= \frac{\sqrt{3}-1}{2\sqrt{2}} \\ &= \frac{\sqrt{3}-1}{\sqrt{6}-\sqrt{2}} \cdot \frac{\sqrt{6}+\sqrt{2}}{\sqrt{6}+\sqrt{2}} \\ &= \frac{(\sqrt{3}-1)(\sqrt{6}+\sqrt{2})}{4}\end{aligned}$$

4.2.2 Example 2

To show that $\tan(45^\circ + A) = \frac{1 + \tan A}{1 - \tan A}$,

Recall that $\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$, so

$$\begin{aligned}\tan(45^\circ + A) &= \frac{\tan 45^\circ + \tan A}{1 - \tan 45^\circ \tan A} \\ &= \frac{1 + \tan A}{1 - (1)\tan A} \quad (\text{Recall that } \tan 45^\circ = 1) \\ \therefore \tan(45^\circ + A) &= \frac{1 + \tan A}{1 - \tan A}\end{aligned}$$

4.2.3 Example 3

To show that $(\cos \theta + \cos \varphi)^2 + (\sin \theta + \sin \varphi)^2 = 2 + 2 \cos(\theta - \varphi)$,

$$\begin{aligned}(\cos \theta + \cos \varphi)^2 + (\sin \theta + \sin \varphi)^2 &= \cos^2 \theta + 2 \cos \theta \cos \varphi + \cos^2 \varphi + \sin^2 \theta + 2 \sin \theta \sin \varphi + \sin^2 \varphi \\&= (\cos^2 \theta + \sin^2 \theta) + (\cos^2 \varphi + \sin^2 \varphi) + (2 \cos \theta \cos \varphi + 2 \sin \theta \sin \varphi) \\&= 1 + 1 + 2(\cos \theta \cos \varphi + \sin \theta \sin \varphi) \\&= 2 + 2 \cos(\theta - \varphi) \quad [\text{Recall that } \cos(A - B) = \cos A \cos B + \sin A \sin B]\end{aligned}$$

4.2.4 Example 4

To show that $\tan(A + B + C) = \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan A \tan C - \tan B \tan C}$,

$$\begin{aligned}
 \tan(A + B + C) &= \frac{\sin(A + B + C)}{\cos(A + B + C)} \\
 &= \frac{\sin[(A + B) + C]}{\cos[(A + B) + C]} \\
 &= \frac{\sin(A + B) \cos C + \cos(A + B) \sin C}{\cos(A + B) \cos C - \sin(A + B) \sin C} \\
 &= \frac{(\sin A \cos B + \cos A \sin B) \cos C + (\cos A \cos B - \sin A \sin B) \sin C}{(\cos A \cos B - \sin A \sin B) \cos C - (\sin A \cos B + \cos A \sin B) \sin C} \\
 &= \frac{\sin A \cos B \cos C + \cos A \sin B \cos C + \cos A \cos B \sin C - \sin A \sin B \sin C}{\cos A \cos B \cos C - \sin A \sin B \cos C - \sin A \cos B \sin C - \cos A \sin B \sin C} \\
 &= \frac{\frac{\sin A \cos B \cos C}{\cos A \cos B \cos C} + \frac{\cos A \sin B \cos C}{\cos A \cos B \cos C} + \frac{\cos A \cos B \sin C}{\sin A \cos B \sin C} - \frac{\sin A \sin B \sin C}{\cos A \sin B \sin C}}{\frac{\cos A \cos B \cos C}{\cos A \cos B \cos C} - \frac{\sin A \sin B \cos C}{\cos A \cos B \cos C} - \frac{\sin A \cos B \sin C}{\cos A \cos B \cos C} - \frac{\cos A \sin B \sin C}{\cos A \cos B \cos C}} \\
 &= \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan A \tan C - \tan B \tan C}
 \end{aligned}$$

$$\therefore \tan(A + B + C) = \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan A \tan C - \tan B \tan C}$$

4.2.5 Example 5

To show that $P + Q + R = 45^\circ$ if $\tan P = \frac{1}{2}$, $\tan Q = \frac{1}{5}$ and $\tan R = \frac{1}{8}$,

$$\begin{aligned} \tan(P + Q + R) &= \frac{\tan P + \tan Q + \tan R - \tan P \tan Q \tan R}{1 - \tan P \tan Q - \tan P \tan R - \tan Q \tan R} \\ &= \frac{\frac{1}{2} + \frac{1}{5} + \frac{1}{8} - \frac{1}{2} \cdot \frac{1}{5} \cdot \frac{1}{8}}{1 - \frac{1}{2} \cdot \frac{1}{5} - \frac{1}{2} \cdot \frac{1}{8} - \frac{1}{5} \cdot \frac{1}{8}} \\ &= \frac{\frac{40+16+10-1}{80}}{\frac{80-8-5-2}{80}} \\ &= \frac{\frac{65}{80}}{\frac{65}{80}} \\ &= \frac{65}{80} \div \frac{65}{80} \\ &= \frac{65}{80} \cdot \frac{80}{65} \\ &= 1 \\ P + Q + R &= \tan^{-1} 1 \\ &= 45^\circ \end{aligned}$$

$$\therefore P + Q + R = 45^\circ \text{ if } \tan P = \frac{1}{2}, \tan Q = \frac{1}{5} \text{ and } \tan R = \frac{1}{8}$$

VOTE KONNECT
FOR ALTRUISTIC
INFORMATION
WITH CLEAR
DISTINCT

5 Multiple Angles

5.1 Double Angles

$$\begin{aligned} \sin 2A &= \sin(A + A) \\ &= \sin A \cos A + \cos A \sin A \\ &= 2 \sin A \cos A \end{aligned}$$

$$\begin{aligned} \cos 2A &= \cos(A + A) \\ &= \cos A \cos A - \sin A \sin A \\ &= \cos^2 A - \sin^2 A \\ &= \cos^2 A - (1 - \cos^2 A) \\ &= \cos^2 A - 1 + \cos^2 A \\ &= 2 \cos^2 A - 1 \end{aligned}$$

$$\begin{aligned} \tan 2A &= \tan(A + A) \\ &= \frac{\tan A + \tan A}{1 - \tan A \tan A} \\ &= \frac{2 \tan A}{1 - \tan^2 A} \end{aligned}$$

5.2 Triple Angles

$$\begin{aligned}
 \sin 3A &= \sin(A + 2A) \\
 &= \sin A \cos 2A + \cos A \sin 2A \\
 &= \sin A \cos(A + A) + \cos A \sin(A + A) \\
 &= \sin A(\cos A \cos A - \sin A \sin A) + \cos A(\sin A \cos A + \cos A \sin A) \\
 &= \sin A(\cos^2 A - \sin^2 A) + \cos A(2 \sin A \cos A) \\
 &= \sin A \cos^2 A - \sin^3 A + 2 \sin A \cos^2 A \\
 &= 3 \sin A \cos^2 A - \sin^3 A \\
 &= 3 \sin A(1 - \sin^2 A) - \sin^3 A \\
 &= 3 \sin A - 3 \sin^3 A - \sin^3 A \\
 &= 3 \sin A - 4 \sin^3 A
 \end{aligned}$$

$$\begin{aligned}
 \cos 3A &= \cos(A + 2A) \\
 &= \cos A \cos 2A - \sin A \sin 2A \\
 &= \cos A \cos(A + A) - \sin A \sin(A + A) \\
 &= \cos A(\cos A \cos A - \sin A \sin A) - \sin A(\sin A \cos A + \cos A \sin A) \\
 &= \cos A(\cos^2 A - \sin^2 A) - \sin A(2 \sin A \cos A) \\
 &= \cos^3 A - \sin^2 A \cos A - 2 \sin^2 A \cos A \\
 &= \cos^3 A - 3 \sin^2 A \cos A \\
 &= \cos^3 A - 3(1 - \cos^2 A) \cos A \\
 &= \cos^3 A - 3 \cos A + 3 \cos^3 A \\
 &= 4 \cos^3 A - 3 \cos A
 \end{aligned}$$

$$\begin{aligned}
 \tan 3A &= \tan(A + 2A) \\
 &= \frac{\tan A + \tan 2A}{1 - \tan A \tan 2A} \\
 &= \frac{\tan A + \tan(A + A)}{1 - \tan A \tan(A + A)} \\
 &= \frac{\tan A + \left(\frac{\tan A + \tan A}{1 - \tan A \tan A}\right)}{1 - \tan A \left(\frac{\tan A + \tan A}{1 - \tan A \tan A}\right)} \\
 &= \frac{\tan A + \frac{2 \tan A}{1 - \tan^2 A}}{1 - \tan A \left(\frac{2 \tan A}{1 - \tan^2 A}\right)} \\
 &= \frac{\tan A(1 - \tan^2 A) + 2 \tan A}{(1 - \tan^2 A) - 2 \tan^2 A} \\
 &= \frac{\tan A - \tan^3 A + 2 \tan A}{1 - \tan^2 A - 2 \tan^2 A} \\
 &= \frac{3 \tan A - \tan^3 A}{1 - 3 \tan^2 A}
 \end{aligned}$$

(Multiplying through by $1 - \tan A$)

VOTE KONNECT
FOR ALTRUISTIC
INFORMATION
WITH CLEAR
DISTINCT

2

6 Submultiple Angles

$$\begin{aligned}\sin A &= \sin \left(\frac{A}{2} + \frac{A}{2} \right) \\ &= \sin \frac{A}{2} \cos \frac{A}{2} + \cos \frac{A}{2} \sin \frac{A}{2} \\ &= 2 \sin \frac{A}{2} \cos \frac{A}{2}\end{aligned}$$

$$\begin{aligned}\cos A &= \cos \left(\frac{A}{2} + \frac{A}{2} \right) \\ &= \cos \frac{A}{2} \cos \frac{A}{2} - \sin \frac{A}{2} \sin \frac{A}{2} \\ &= \cos^2 \frac{A}{2} - \sin^2 \frac{A}{2} \\ &= \cos^2 \frac{A}{2} - (1 - \cos^2 \frac{A}{2}) \\ &= \cos^2 \frac{A}{2} - 1 + \cos^2 \frac{A}{2} \\ &= 2 \cos^2 \frac{A}{2} - 1\end{aligned}$$

$$\begin{aligned}\tan A &= \tan \left(\frac{A}{2} + \frac{A}{2} \right) \\ &= \frac{\tan \frac{A}{2} + \tan \frac{A}{2}}{1 - \tan \frac{A}{2} \tan \frac{A}{2}} \\ &= \frac{2 \tan \frac{A}{2}}{1 - \tan^2 \frac{A}{2}}\end{aligned}$$

7 Product and Sum Formulae

7.1 Derivation of the Formulae

Recall the following:

$$\sin(A + B) = \sin A \cos B + \cos A \sin B \quad (7.1)$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B \quad (7.2)$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B \quad (7.3)$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B \quad (7.4)$$

(7.1) + (7.2) gives

$$\sin(A + B) + \sin(A - B) = 2 \sin A \cos B \quad (7.5)$$

(7.1) - (7.2) gives

$$\sin(A + B) - \sin(A - B) = 2 \cos A \sin B \quad (7.6)$$

(7.3) + (7.4) gives

$$\cos(A + B) + \cos(A - B) = 2 \cos A \cos B \quad (7.7)$$

(7.3) - (7.4) gives

$$\cos(A + B) - \cos(A - B) = -2 \sin A \sin B \quad (7.8)$$

$$\cos(A - B) - \cos(A + B) = 2 \sin A \sin B \quad (7.9)$$

Given that

$$A + B = C \quad (7.10)$$

$$A - B = D \quad (7.11)$$

We have

$$A = \frac{C + D}{2} \quad (7.12)$$

$$B = \frac{C - D}{2} \quad (7.13)$$

Substituting (7.10)-(7.13) into (7.5)-(7.9), we have

$$\sin C + \sin D = 2 \sin \frac{C + D}{2} \cos \frac{C - D}{2} \quad (7.14)$$

$$\sin C - \sin D = 2 \cos \frac{C + D}{2} \sin \frac{C - D}{2} \quad (7.15)$$

$$\cos C + \cos D = 2 \cos \frac{C + D}{2} \cos \frac{C - D}{2} \quad (7.16)$$

$$\cos D - \cos C = 2 \sin \frac{C + D}{2} \sin \frac{C - D}{2} \quad (7.17)$$

7.2 Examples

7.2.1 Example 1

To express $\sin 5x + \sin x$ as a product of trigonometric ratios,

Comparing

with

$$\sin C + \sin D = 2 \sin \frac{C + D}{2} \cos \frac{C - D}{2}$$

It is clear that $C = 5x$ and $D = x$, so

$$\begin{aligned} \sin 5x + \sin x &= 2 \sin \frac{5x + x}{2} \cos \frac{5x - x}{2} \\ &= 2 \sin \frac{6x}{2} \cos \frac{4x}{2} \\ &= 2 \sin 3x \cos 2x \end{aligned}$$

7.2.2 Example 2

To express $2 \sin 3x \cos 2x$ as a sum of trigonometric ratios,

Comparing

with
$$2 \sin \frac{C+D}{2} \cos \frac{C-D}{2} = \sin C + \sin D$$

We see that,

$$\begin{aligned} \frac{C+D}{2} &= 3x \\ C+D &= 6x \end{aligned} \quad (7.18)$$

Also,

$$\begin{aligned} \frac{C-D}{2} &= 2x \\ C-D &= 4x \end{aligned} \quad (7.19)$$

(7.18)+(7.19) gives

$$\begin{aligned} 2C &= 10x \\ C &= 5x \end{aligned}$$

(7.18)-(7.19) gives

$$\begin{aligned} 2D &= 2x \\ D &= x \end{aligned}$$

$$\therefore 2 \sin 3x \cos 2x = \sin 5x + \sin x$$

7.2.3 Example 3

To show that $\sin 2A \cos 4A + \sin 3A \cos 9A = \frac{1}{2} (\sin 12A - \sin 2A)$,

From
$$\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2}$$

We have
$$\sin \frac{C+D}{2} \cos \frac{C-D}{2} = \frac{1}{2} (\sin C + \sin D)$$

For $\sin 2A \cos 4A$,

$$\begin{aligned} \frac{C+D}{2} &= 2A \\ C+D &= 4A \end{aligned} \quad (7.20)$$

$$\begin{aligned} \frac{C-D}{2} &= 4A \\ C-D &= 8A \end{aligned} \quad (7.21)$$

(7.20)+(7.21) gives

$$2C = 12A$$

$$C = 6A$$

(7.20)-(7.21) gives

$$2D = -4A$$

$$D = -2A$$

$$\therefore \sin 2A \cos 4A = \frac{1}{2} (\sin 6A + \sin(-2A)) = \frac{1}{2} (\sin 6A - \sin 2A)$$

For $\sin 3A \cos 9A$,

$$\frac{C+D}{2} = 3A$$

$$C + D = 6A$$

(7.22)

$$\frac{C-D}{2} = 9A$$

$$C - D = 18A$$

(7.23)

(7.22)+(7.23) gives

$$2C = 24A$$

$$C = 12A$$

(7.22)-(7.23) gives

$$2D = -12A$$

$$D = -6A$$

$$\therefore \sin 3A \cos 9A = \frac{1}{2} (\sin 12A + \sin(-6A)) = \frac{1}{2} (\sin 12A - \sin 6A)$$

Now,

$$\begin{aligned} \sin 2A \cos 4A + \sin 3A \cos 9A &= \frac{1}{2} (\sin 6A - \sin 2A) + \frac{1}{2} (\sin 12A - \sin 6A) \\ &= \frac{1}{2} [(\sin 6A - \sin 2A) + (\sin 12A - \sin 6A)] \\ &= \frac{1}{2} (\sin 6A - \sin 2A + \sin 12A - \sin 6A) \\ &= \frac{1}{2} (\sin 12A - \sin 2A) \end{aligned}$$

$$\therefore \sin 2A \cos 4A + \sin 3A \cos 9A = \frac{1}{2} (\sin 12A - \sin 2A)$$

7.2.4 Example 4

To show that $\sin 7x + \sin x - 2 \sin 2x \cos 3x = 4 \cos^2 3x \sin x$,

Since $\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2}$, we have

$$\begin{aligned} \sin 7x + \sin x - 2 \sin 2x \cos 3x &= 2 \sin \frac{7x+x}{2} \cos \frac{7x-x}{2} - 2 \sin 2x \cos 3x \\ &= 2 \sin \frac{8x}{2} \cos \frac{6x}{2} - 2 \sin 2x \cos 3x \\ &= 2 \sin 4x \cos 3x - 2 \sin 2x \cos 3x \\ &= 2 \cos 3x (\sin 4x - \sin 2x) \end{aligned}$$

Since $\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}$, we have

$$\begin{aligned} \sin 7x + \sin x - 2 \sin 2x \cos 3x &= 2 \cos 3x \cdot 2 \cos \frac{4x+2x}{2} \sin \frac{4x-2x}{2} \\ &= 2 \cos 3x \cdot 2 \cos \frac{6x}{2} \sin \frac{2x}{2} \\ &= 2 \cos 3x (2 \cos 3x \sin x) \\ &= 4 \cos^2 3x \sin x \end{aligned}$$

$$\therefore \sin 7x + \sin x - 2 \sin 2x \cos 3x = 4 \cos^2 3x \sin x$$

7.2.5 Example 5

If $\sin \theta + \sin \varphi = a$ and $\cos \theta + \cos \varphi = b$, to show that $\cos^2 \frac{1}{2}(\theta - \varphi) = \frac{1}{4}(a^2 + b^2)$,

Since $\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2}$,

$$2 \sin \frac{\theta + \varphi}{2} \cos \frac{\theta - \varphi}{2} = a \quad (7.24)$$

Since $\cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2}$,

$$2 \cos \frac{\theta + \varphi}{2} \cos \frac{\theta - \varphi}{2} = b \quad (7.25)$$

Squaring (7.24) and (7.25),

$$4 \sin^2 \frac{\theta + \varphi}{2} \cos^2 \frac{\theta - \varphi}{2} = a^2 \quad (7.26)$$

$$4 \cos^2 \frac{\theta + \varphi}{2} \cos^2 \frac{\theta - \varphi}{2} = b^2 \quad (7.27)$$

Adding (7.26) and (7.27),

$$\begin{aligned}
4 \sin^2 \frac{\theta + \varphi}{2} \cos^2 \frac{\theta - \varphi}{2} + 4 \cos^2 \frac{\theta + \varphi}{2} \sin^2 \frac{\theta - \varphi}{2} &= a^2 + b^2 \\
4 \cos^2 \frac{\theta - \varphi}{2} \sin^2 \frac{\theta + \varphi}{2} + 4 \sin^2 \frac{\theta - \varphi}{2} \cos^2 \frac{\theta + \varphi}{2} &= a^2 + b^2 \\
4 \cos^2 \frac{\theta - \varphi}{2} &= a^2 + b^2 \quad (1) \\
\cos^2 \frac{\theta - \varphi}{2} &= \frac{1}{4}(a^2 + b^2) \\
\cos^2 \frac{1}{2}(\theta - \varphi) &= \frac{1}{4}(a^2 + b^2)
\end{aligned}$$

$\therefore \cos^2 \frac{1}{2}(\theta - \varphi) = \frac{1}{4}(a^2 + b^2)$ given that $\sin \theta + \sin \varphi = a$ and $\cos \theta + \cos \varphi = b$

8 A Revisit to Sub-Multiple Angles

8.1 A Transformation of the Sub-Multiple Angles

Recall that

$$\tan x = \frac{2 \tan \frac{x}{2}}{1 - \tan^2 \frac{x}{2}}$$

If $\tan \frac{x}{2} = t$, then

$$\tan x = \frac{2t}{1 - t^2} \quad (8.1)$$

For $\sin x$, we proceed as follows

$$\begin{aligned}
\sin x &= 2 \sin \frac{x}{2} \cos \frac{x}{2} \\
&= \frac{2 \sin \frac{x}{2} \cos \frac{x}{2}}{1} \\
&= \frac{2 \sin \frac{x}{2} \cos \frac{x}{2}}{\cos^2 \frac{x}{2} + \sin^2 \frac{x}{2}} \\
&= \frac{\frac{2 \sin \frac{x}{2} \cos \frac{x}{2}}{\cos^2 \frac{x}{2}}}{\frac{\cos^2 \frac{x}{2}}{\cos^2 \frac{x}{2}} + \frac{\sin^2 \frac{x}{2}}{\cos^2 \frac{x}{2}}} \\
&= \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} \\
\therefore \sin x &= \frac{2t}{1 + t^2} \quad (8.2)
\end{aligned}$$

For $\cos x$, we have

$$\begin{aligned}
 \cos x &= \frac{\cos^2 \frac{x}{2} - \sin^2 \frac{x}{2}}{\cos^2 \frac{x}{2} + \sin^2 \frac{x}{2}} \\
 &= \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} \\
 &= \frac{\frac{\cos^2 \frac{x}{2}}{\cos^2 \frac{x}{2}} - \frac{\sin^2 \frac{x}{2}}{\cos^2 \frac{x}{2}}}{\frac{\cos^2 \frac{x}{2}}{\cos^2 \frac{x}{2}} + \frac{\sin^2 \frac{x}{2}}{\cos^2 \frac{x}{2}}} \\
 &= \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} \\
 \therefore \cos x &= \frac{1 - t^2}{1 + t^2} \quad (8.3)
 \end{aligned}$$

8.2 Examples

8.2.1 Example 1

To calculate $\tan \frac{1}{2} \theta$, if $\tan \theta = \frac{24}{7}$ and θ is acute,

$$\begin{aligned}
 \tan \theta &= \frac{2 \tan \frac{1}{2} \theta}{1 - \tan^2 \frac{1}{2} \theta} \\
 \tan \theta &= \frac{2t}{1 - t^2} \quad (\text{Since } \tan \frac{1}{2} \theta = t) \\
 \frac{24}{7} &= \frac{2t}{1 - t^2} \\
 7(2t) &= 24(1 - t^2) \\
 14t &= 24 - 24t^2 \\
 24t^2 + 14t - 24 &= 0 \\
 12t^2 + 7t - 12 &= 0 \\
 12t^2 + 16t - 9t - 12 &= 0 \\
 (12t^2 + 16t) - (9t + 12) &= 0 \\
 4t(3t + 4) - 3(3t + 4) &= 0 \\
 (3t + 4)(4t - 3) &= 0 \\
 t &= -\frac{4}{3} \text{ or } \frac{3}{4}
 \end{aligned}$$

8.2.2 Example 2

Given that $\sec \theta - \tan \theta = x$. Show that $t = \frac{1-x}{1+x}$ where $t = \tan(\frac{\theta}{2})$.
(This is left as an exercise for students).

9 Trigonometric Equations

9.1 Overview

Trigonometric equations are equations which involve the six trigonometric ratios.

In solving trigonometric equations, we get the values of the angles in the four quadrants from $0^\circ - 360^\circ$ that satisfy the equations.

We recall the following about the four quadrants:




Figure 2: The Four Quadrants

In the first quadrant,

$$\sin \theta = \sin \theta$$

$$\cos \theta = \cos \theta$$

$$\tan \theta = \tan \theta$$

In the second quadrant,

$$\sin \theta = \sin(180 - \theta)^\circ$$

$$\cos \theta = - \cos(180 - \theta)^\circ$$

$$\tan \theta = - \tan(180 - \theta)^\circ$$

In the third quadrant,

$$\sin \theta = - \sin(\theta - 180)^\circ$$

$$\cos \theta = - \cos(\theta - 180)^\circ$$

$$\tan \theta = \tan(\theta - 180)^\circ$$

In the fourth quadrant,

$$\sin \theta = - \sin(360 - \theta)^\circ$$

$$\cos \theta = \cos(360 - \theta)^\circ$$

$$\tan \theta = - \tan(360 - \theta)^\circ$$

9.2 Examples

9.2.1 Example 1

To find the angles less than 360° which satisfy $6 \sin \theta = \tan \theta$,

$$\begin{aligned}
6 \sin \theta &= \frac{\tan \theta}{\sin \theta} \\
6 \sin \theta &= \frac{1}{\cos \theta} \\
6 \sin \theta \cos \theta &= 1 \\
6 \cos \theta &= \frac{1}{\sin \theta} \\
\cos \theta &= \frac{1}{6} \\
\theta &= \cos^{-1} \frac{1}{6} \\
\theta &= \cos^{-1}(0.1667) \\
\theta &= 80.4^\circ, 279.6^\circ
\end{aligned}$$

9.2.2 Example 2

To find the angles less than 360° which satisfy $3 \cos^2 \theta + 5 \sin^2 \theta = 4$,

$$\begin{aligned}
3 \cos^2 \theta + 5 \sin^2 \theta &= 4 \\
3 \cos^2 \theta + 5(1 - \cos^2 \theta) &= 4 \\
3 \cos^2 \theta + 5 - 5 \cos^2 \theta &= 4 \\
3 \cos^2 \theta - 5 \cos^2 \theta &= 4 - 5 \\
-2 \cos^2 \theta &= -1 \\
2 \cos^2 \theta &= 1 \\
\cos^2 \theta &= \frac{1}{2} \\
\cos \theta &= \pm \sqrt{\frac{1}{2}} \\
\cos \theta &= \pm \frac{1}{\sqrt{2}} \\
\cos \theta &= \frac{1}{\sqrt{2}} \text{ or } \cos \theta = -\frac{1}{\sqrt{2}} \\
\theta &= \cos^{-1} \frac{1}{\sqrt{2}} \text{ or } \theta = \cos^{-1} -\frac{1}{\sqrt{2}} \\
\theta &= 45^\circ, 315^\circ \text{ or } \theta = 135^\circ, 225^\circ \\
\therefore \theta &= 45^\circ, 315^\circ, 135^\circ, 225^\circ
\end{aligned}$$

9.2.3 Example 3

To solve $5 \tan^2 \theta - \sec^2 \theta = 11$ for $0^\circ \leq \theta \leq 360^\circ$,

$$\begin{aligned}
5 \tan^2 \theta - \sec^2 \theta &= 11 \\
5 \tan^2 \theta - (1 + \tan^2 \theta) &= 11 \\
5 \tan^2 \theta - 1 - \tan^2 \theta &= 11 \\
5 \tan^2 \theta - \tan^2 \theta &= 11 + 1 \\
4 \tan^2 \theta &= 12
\end{aligned}$$

$$\begin{aligned}\tan^2 \theta &= \frac{3}{\sqrt{3}} \\ \tan \theta &= \pm \sqrt{3} \\ \theta &= \tan^{-1} \sqrt{3} \text{ or } \theta = \tan^{-1}(-\sqrt{3}) \\ \theta &= 60^\circ, 240^\circ \text{ or } \theta = 120^\circ, 300^\circ \\ \theta &= 60^\circ, 240^\circ, 120^\circ, 300^\circ\end{aligned}$$

9.2.4 Example 4

To solve $6 \cos^2 \theta + \sin \theta - 5 = 0$ for $0^\circ \leq \theta \leq 360^\circ$,

$$\begin{aligned}6 \cos^2 \theta + \sin \theta - 5 &= 0 \\ 6(1 - \sin^2 \theta) + \sin \theta - 5 &= 0 \\ 6 - 6 \sin^2 \theta + \sin \theta - 5 &= 0 \\ -6 \sin^2 \theta + \sin \theta + 1 &= 0 \\ 6 \sin^2 \theta - \sin \theta - 1 &= 0 \\ (\text{Let } \sin \theta = x) \quad 6x^2 - x - 1 &= 0 \\ 6x^2 - 3x + 2x - 1 &= 0 \\ (6x^2 - 3x) + (2x - 1) &= 0 \\ 3x(2x - 1) + 1(2x - 1) &= 0 \\ 2x - 1 = 0 \text{ or } 3x + 1 &= 0 \\ x = \frac{1}{2} \text{ or } x = -\frac{1}{3} \\ (\text{Recall that } x = \sin \theta) \quad \sin \theta = \frac{1}{2} \text{ or } \sin \theta = -\frac{1}{3} \\ \theta = \sin^{-1} \frac{1}{2} \text{ or } \theta = \sin^{-1} -\frac{1}{3} \\ \theta = 30^\circ, 150^\circ \text{ or } \theta = 340.5^\circ, 119.5^\circ\end{aligned}$$

9.3 Exercise

Solve the following for $0^\circ \leq \theta \leq 360^\circ$:

- $4 \cos 2\theta - 3 = 0$
- $3 \sin 2\theta = 18$
- $4 \cos \theta = 3 \tan \theta$
- $2 \tan^2 \theta - 3 \tan \theta + 1 = 0$
- $\sin \theta + \sin^2 \theta = 0$

Reference

B.D. Bunday and H. Mulholland, “Pure Mathematics for Advanced Level”. Heinemann Educational Books Ltd, Halley Court Jordan Hill, Oxford OX2 8EJ, P.M.B. 5205, Ibadan. P.O. Box 45314, Nairobi, 2004.

Compiled by:

- **Afolabi, Ayodeji Sunday** (asafolabi@futa.edu.ng)
- **Dansu, Emmanuel Jesuyon** (ejdansu@futa.edu.ng)

VOTE KONNECT FOR ALTRUISTIC
INFORMATION WITH CLEAR DISTINCT

KONNECT'18

KONNECT'18