

PARTICULARS TO APPEAR ON THE OUTER PACKAGE {CARDBOARD BOX}

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Zycortal 25 mg/ml prolonged-release suspension for injection

2. STATEMENT OF ACTIVE SUBSTANCES

Each ml contains:
Desoxycortone pivalate 25 mg

3. PACKAGE SIZE

4 ml

4. TARGET SPECIES

Dogs.

5. INDICATIONS

6. ROUTES OF ADMINISTRATION

Subcutaneous use.

7. WITHDRAWAL PERIODS

8. EXPIRY DATE

Exp. {mm/yyyy}
Once opened use within 4 months.

9. SPECIAL STORAGE PRECAUTIONS

Do not store above 30 °C.
Do not freeze.

10. THE WORDS “READ THE PACKAGE LEAFLET BEFORE USE”

Read the package leaflet before use.

11. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

12. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

13. NAME OF THE MARKETING AUTHORISATION HOLDER

Dechra Regulatory B.V.

14. MARKETING AUTHORISATION NUMBERS

Vm 50406/5053

15. BATCH NUMBER

Lot {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING
UNITS {VIAL LABEL}**

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Zycortal

2. QUANTITATIVE PARTICULARS OF THE ACTIVE SUBSTANCES

Desoxycortone pivalate 25 mg/ml

3. BATCH NUMBER

Lot {number}

4. EXPIRY DATE

Exp. {mm/yyyy}

Once opened use within 4 months.

PARTICULARS TO APPEAR ON THE PACKAGE LEAFLET:

PACKAGE LEAFLET

1. Name of the veterinary medicinal product

Zycortal 25 mg/ml Prolonged-release Suspension for Injection for Dogs

2. Composition

Each ml contains:

Active substances:

Desoxycortone pivalate 25 mg

Excipients:

Chlorocresol 1 mg

Opaque white suspension.

3. Target species

Dogs.

4. Indications for use

For use as replacement therapy for mineralocorticoid deficiency in dogs with primary hypoadrenocorticism (Addison's disease).

5. Contraindications

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

6. Special warnings

Special warnings:

Before starting treatment with the veterinary medicinal product, it is important that Addison's disease has been definitively diagnosed. Any dog presenting with severe hypovolaemia, dehydration, pre-renal azotaemia and inadequate tissue perfusion (also known as "Addisonian crisis") must be rehydrated with intravenous fluid (saline) therapy before starting treatment with the veterinary medicinal product.

Special precautions for safe use in the target species:

Use with caution in dogs with congestive heart disease, severe renal disease, primary hepatic failure or oedema.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Avoid contact with the eyes and skin. In case of accidental spillage onto the skin or eyes, wash the affected area with water. If irritation occurs, seek medical advice immediately and show the package leaflet or the label to the physician.

This veterinary medicinal product may cause pain and swelling at the injection site if accidentally self-administered.

This veterinary medicinal product may cause adverse events on male reproductive organs and, as a result, fertility.

This veterinary medicinal product may cause adverse developmental effects on unborn children and neonates.

The veterinary medicinal product should not be administered by pregnant or breast-feeding women.

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during breeding, pregnancy or lactation. Use only according to the benefit-risk assessment by the responsible veterinarian.

Interaction with other medicinal products and other forms of interaction:

Use caution when administering Zycortal concurrently with medicinal products which affect either serum sodium or potassium concentrations, or cellular transportation of sodium or potassium, for example: trimethoprim, amphotericin B, or digoxin or insulin.

Overdose:

When given to dogs at three to five times the recommended dose, injection site reactions, characterised by erythema and oedema occurred.

As expected from the pharmacodynamic effects, escalating doses of desoxycortone are associated with a dose-related trend for increased serum sodium, and decreased blood urea nitrogen, serum potassium and urine specific gravity. Polyuria and polydipsia may be observed.

Hypertension has been observed in dogs receiving 20 mg/kg of desoxycortone pivalate.

There is no specific antidote. In case of signs of overdose, the dog should be treated symptomatically, and subsequent doses reduced.

Major incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

7. Adverse events

Dogs:

Very common (>1 animal / 10 animals treated):	Polydipsia (excessive drinking) Polyuria (excessive urination)
Common (1 to 10 animals / 100 animals treated):	Inappropriate urination Lethargy, Decreased appetite, Anorexia, Decreased activity, Depression, Polyphagia (excessive eating), Tiredness Alopecia (hair loss) Panting Vomiting, Diarrhoea Shaking Urinary tract infection
Uncommon (1 to 10 animals / 1 000 animals treated):	Injection site pain
Rare (1 to 10 animals / 10 000 animals treated):	Pancreas disorder ^a

^a The concurrent administration of glucocorticoids may contribute to these signs.

Reporting adverse events is important. It allows continuous safety monitoring of a product. If you notice any side effects, even those not already listed in this package leaflet, or you think that the medicine has not worked, please contact, in the first instance, your veterinarian. You can also report any adverse events to the marketing authorisation holder or its local representative using the contact details at the end of this leaflet, or via your national reporting system at:

Website: <https://www.gov.uk/report-veterinary-medicine-problem/animal-reacts-medicine>

e-mail: adverse.events@vmd.gov.uk

8. Dosage for each species, routes and method of administration

Subcutaneous use.

Zycortal replaces the mineralocorticoid hormones only. Dogs with combined glucocorticoid and mineralocorticoid deficiency should also receive a glucocorticoid such as prednisolone taking into account the current scientific knowledge.

Zycortal is intended for long term administration at intervals and doses dependent upon individual response. Tailor the dose of Zycortal and the concurrently administered glucocorticoid replacement therapy to the individual dog based on clinical response and normalization of Na^+ and K^+ serum concentrations.

Initial dose of Zycortal:

The initial dose is 2.2 mg/kg body weight, administered by subcutaneous injection.

Interim monitoring visit:

Re-evaluate the dog and measure the serum sodium/potassium ratio (Na^+/K^+ ratio) approximately 10 days after the first dose (which is the time to maximum concentration (T_{max}) of desoxycortone). If the dog's clinical signs have worsened or not resolved, adjust the dose of glucocorticoid and/or investigate other causes of the clinical signs.

Second dose of Zycortal:

At approximately 25 days after the first dose, re-evaluate the dog and measure the Na^+/K^+ ratio.

- If the dog is both clinically normal and has a normal Na^+/K^+ ratio (i.e. 27 to 32) on Day 25, adjust the dose based on the Day 10 Na^+/K^+ ratio using the guidelines in Table 1, below.
- If the dog is clinically normal and has a Na^+/K^+ ratio > 32 on Day 25, either adjust the dose based on the Day 10 Na^+/K^+ ratio according to Table 1 or delay the dose (see Prolonging the dosing interval).
- If the dog is either not clinically normal or if the Na^+/K^+ ratio is abnormal on Day 25, adjust the dose of glucocorticoid or Zycortal (see Subsequent doses and long term management).

Table 1: Day 25: Administering the Second Dose of Zycortal

If the Day 10 Na^+/K^+ ratio is:		25 days after the first dose, administer Zycortal, as follows:
≥ 34	Do not administer Dose 2 on Day 10.	Decrease dose to: 2.0 mg/kg body weight
32 to < 34		Decrease dose to: 2.1 mg/kg body weight
27 to < 32		Continue 2.2 mg/kg body weight
≥ 24 to < 27		Increase dose to: 2.3 mg/kg body weight
< 24		Increase dose to: 2.4 mg/kg body weight

Prolonging the dosing interval:

If the dog is clinically normal and the Day 25 Na^+/K^+ ratio is > 32 , it is possible to prolong the dosing interval instead of adjusting the dose as described in Table 1. Evaluate the electrolytes every 5–9 days until the Na^+/K^+ ratio is < 32 , and then administer 2.2 mg/kg of Zycortal.

Subsequent doses and long term management:

Once the optimal dose and dosing interval have been determined, maintain the same regimen. If the dog develops abnormal clinical signs or Na^+ or K^+ serum concentrations, use the following guidelines for subsequent doses:

- Clinical signs of polyuria/polydipsia: Decrease the glucocorticoid dose first. If the polyuria/polydipsia persists and the Na^+/K^+ ratio is >32 , then decrease the dose of Zycortal without changing the dosing interval.
- Clinical signs of depression, lethargy, vomiting, diarrhoea or weakness: Increase the glucocorticoid dose.
- Hyperkalaemia, hyponatremia or Na^+/K^+ ratio < 27 : Decrease the Zycortal dosing interval by 2–3 days or increase the dose.
- Hypokalaemia, hypernatremia or Na^+/K^+ ratio > 32 : Decrease the Zycortal dose.

Prior to a stressful situation, consider temporarily increasing the dose of glucocorticoid.

In the clinical trial, the mean final dose of Zycortal was 1.9 mg/kg (range 1.2–2.5 mg/kg) and the mean final dosing interval was 38.7 ± 12.7 days (range 20–99 days) with the majority of dogs having a dosing interval between 20 and 46 days.

9. Advice on correct administration

Prior to use, gently shake the vial to resuspend the product.

Use an appropriately graduated syringe to accurately administer the required dose volume. This is particularly important when injecting small volumes.

To ensure a correct dosage, body weight should be determined as accurately as possible.

The use of suitably calibrated measuring equipment is recommended.

10. Withdrawal periods

Not applicable.

11. Special storage precautions

Keep out of the sight and reach of children.

Do not store above 30 °C.

Do not freeze.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and the vial after Exp. The expiry date refers to the last day of that month.

Shelf life after first opening the immediate packaging: 4 months.

12. Special precautions for disposal

Medicines should not be disposed of via wastewater.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any applicable national collection systems. These measures should help to protect the environment.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required.

13. Classification of veterinary medicinal products

Veterinary medicinal product subject to prescription.

14. MARKETING AUTHORISATION NUMBERS AND PACK SIZES

Vm 50406/5053

Type I glass vial (4 ml) with a coated chlorobutyl rubber stopper and aluminium seal with a plastic flip-off cap.

One vial of 4 ml in a cardboard box.

15. PID LINK (Do not print heading)

[The following statement must be included where reference to the European Union Product Database is included on the product information. This statement is relevant to both UK(GB) and UK(NI) products:]

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

16. Contact details

Marketing authorisation holder:

Dechra Regulatory B.V.
Handelsweg 25
5531 AE Bladel
The Netherlands

Manufacturer responsible for batch release:

Eurovet Animal Health B.V.
Handelsweg 25
5531 AE Bladel
The Netherlands

Local representatives and contact details to report suspected adverse events:

Dechra Veterinary Products Limited
Sansaw Business Park
Hadnall
Shrewsbury
Shropshire
SY4 4AS
United Kingdom
Tel: +44 (0) 1939 211200

17. Other information

POM-V

Gavin Hall

Approved: 10 December 2025