

What is a (Mathematical) Convolution

- The definition

- Continuous (<https://mathworld.wolfram.com/Convolution.html>)

$$[f * g](t) = \int_0^t f(\tau)g(t - \tau)d\tau$$

- Discrete

$$[x * w](n) = \sum_{m=-\infty}^{\infty} x[n - m]w[m]$$

- What to do with finite sequences
 - pad with zeros and suffer edge effects
 - circular convolution (assumed periodicity)
- 2D and more (can be generalised to images/matrices, tensors, groups ++)

“

- Another Mathsy View

- Convolution is a linear transform by a Toeplitz matrix

“

- The Signal Processing View

- Think of x as a signal, and w as a *kernel*
- A convolution is a Linear Time-Invariant (LTI) filter
 - Keywords: Low-pass, High-pass, Band-pass, ARMA, IIR, FIR, ...
- Implemented as taps, block diagrams, ...

“

- EEEngineers view

- Convolution is a LTI *system*, e.g., represents dynamics

“

- The Frequency View
 - $x * w = F^{-1}(F(x)F(w))$
 - Fourier transform is just a set of convolutions (1 for each frequency band)

“

- The Statisticians View
 - A convolution is just a correlation coefficient between x and lagged versions of \bar{w} (time reversed w)
 - In time series, a convolution is used to create an ARMA system
 - A convolution is used in kernel smoothing/density approaches

“

- The probabilists View
 - A convolution is how you compute the distribution function of the sum of two independent RVs

“

- The physicist View
 - Doppler blurring
 - Out-of-focus optics

“

- Machine Learning
 - CNNs
 - blurring, smoothing, edge detection, ...

These views are all the same.

