

Optimal Time-Stepping with Non-Iterative Defect Correction

for the Moment Hierarchy of Stochastic Reaction–Diffusion Equations

A living paper: derivation, verification ledger, and an honest map from closed moment systems to Cole–Hopf and KPZ observables

Aditya Tiwari

Living document, version 0.3 — September 16, 2026

Repository: OTS-NIDC (this paper is generated from it)

1 Review for Single Variable PDEs

Throughout, $\delta_x^2 u_j = (u_{j+1} - 2u_j + u_{j-1})/\Delta x^2$ is the centered second difference and refinement means $\Delta x \rightarrow 0$ with Δt slaved to Δx .

1.1 The scalar family and the optimal time step

Consider the deterministic family

$$u_t = D u_{xx} + \alpha u, \quad D > 0, \alpha \in \mathbb{R}, \quad (1)$$

whose stochastic motivation (the moment equations of (??), with $\alpha = \frac{k(k-1)}{2}\sigma^2$) is deferred to §??. The pure diffusion case $\alpha = 0$ is exactly Chu’s setting (Chu, 2011).

Apply forward Euler in time and the three-point stencil in space. Taylor expansion of both sides, followed by repeated differentiation of (1) to convert time derivatives of u into space derivatives (the operators $D\partial_{xx}$ and α commute, giving the binomial identity $u_{t^k} = (D\partial_{xx} + \alpha)^k u$), yields the local truncation error organized by derivative type:

$$\tau = \underbrace{\left(\frac{D^2\Delta t}{2} - \frac{D\Delta x^2}{12}\right)u_{xxxx}}_{\text{Group 1}} + \underbrace{\alpha D \Delta t u_{xx}}_{\text{Group 2}} + \underbrace{\frac{\alpha^2\Delta t}{2}u}_{\text{Group 3}} + O(\Delta t^2, \Delta x^4). \quad (2)$$

Only Group 1 couples Δt to Δx . Setting it to zero:

Theorem 1 (Optimal time step (Chu, 2011)). *For (1) discretized by forward Euler / centered differences, the unique step size annihilating the leading (u_{xxxx}) truncation term is*

$$\boxed{\Delta t_{\text{opt}} = \frac{\Delta x^2}{6D}} \quad (3)$$

and no single step annihilates both the u_{xxxx} and u_{xxxxx} truncation terms, since the level-2 condition is $\Delta t = \Delta x^2/(\sqrt{60}D) \neq \Delta t_{\text{opt}}$.

(Proof: one-line cancellation in (2); uniqueness by the level-2 computation. Full algebra in `sign-change.tex` §1.)

Proposition 2 (α -independence). *Δt_{opt} is independent of the reaction parameter α : the cancellation operates on the diffusive group alone. For the stochastic hierarchy this makes Δt_{opt} independent of the noise amplitude σ and of the moment order k .*

Following the derivation inspecting the truncation error, at $\Delta t = \Delta t_{\text{opt}}$ the Group-1 term vanishes and the surviving residual is

$$\tau|_{\Delta t_{\text{opt}}} = \frac{\alpha \Delta x^2}{6} u_{xx} + \frac{\alpha^2 \Delta x^2}{12D} u + O(\Delta x^4). \quad (4)$$

Then the truncation error τ as assembled in (2) is the *forward* residual: $\text{exact} - \text{scheme} = \Delta t \tau + \dots$. Cancelling it therefore requires *adding* $\Delta t \tau$.

Lemma 3 (Closed form correction term). *The corrected factor at any given time step is,*

$$C_j^{(1)} = +\frac{\alpha \Delta x^4}{36D} \delta_x^2 u_j^n + \frac{\alpha^2 \Delta x^4}{72D^2} u_j^n. \quad (5)$$

Theorem 4 (First-level accuracy). *The corrected scheme*

$$u_j^{n+1} = u_j^n + \Delta t_{\text{opt}} \left[D \delta_x^2 u_j^n + \alpha u_j^n \right] + C_j^{(1)} \quad (6)$$

has step error $O(\Delta x^6)$ and global order 4: the correction $C^{(1)}$ is itself $O(\Delta x^4)$, so its evaluation with the second-order operator δ_x^2 contributes only $O(\Delta x^6)$ to the step — the three-point stencil is mathematically sufficient at $p = 1$.

Status: [V]. Measured orders and errors, with the harness described in Appendix ??:

Table 1: Scalar $p = 1$ convergence, $u_t = Du_{xx} + \alpha u$ with $D = 0.1$, $\alpha \sigma^2 = 0.25$, Neumann BCs, $T = 0.05$; L^2 error against the exact cosine-series solution. Data: `code/implementation/results.md` (generated by `convergence_test.py`, 2026-07). All methods use the same grid and step size Δt_{opt} at each level.

Method	$N = 32$	$N = 64$	$N = 128$	$N = 256$
OTS-NIDC ($p = 1$)	8.54e-09	5.26e-10	3.27e-11	2.04e-12
Forward Euler	1.23e-05	3.02e-06	7.52e-07	1.88e-07
4th-order FD + FE	1.52e-05	3.79e-06	9.45e-07	2.36e-07
Reference expm	2.77e-05	6.81e-06	1.70e-06	4.23e-07
<i>Measured order (log-log slope, mean over levels):</i>				
OTS-NIDC	4.01	(FE / 4th-order FD / expm: 2.00–2.01)		

The 4th-order-FD row carries the paper’s cleanest negative control: high-order spatial accuracy alone buys nothing when the step size is slaved to Δx^2 — the $O(\Delta t)$ time error caps the whole scheme at second order. The boost must come from the time-step structure; that is the method’s thesis, verified.

The p -level tower and the stencil bound

move this into an appendix, perhaps?

The general correction is generated by powers of the operator, with the sign of Remark ??:

$$C_j^{(\leq p)} = + \sum_{\ell=1}^p \frac{\Delta t_{\text{opt}}^\ell}{(\ell+1)!} (D\mathcal{D}_2 + \alpha)^\ell u_j^n, \quad (7)$$

where $\mathcal{D}_2$ is a spatial operator whose accuracy must be raised as ℓ grows. Spatial discretization error breaks the binomial symmetry of the time expansion, giving the stencil-scaling bound:

Proposition 5 (Stencil bound; sketch). *Global order $2p+2$ requires corrections evaluated on at least a $(2p+3)$ -point stencil: $p=1$ needs the 3-point main stencil plus a 5-point evaluation inside $C^{(1)}$ when full fourth-order spatial accuracy is demanded of the correction itself; $p=2$ needs a 7-point stencil to represent the uncanceled u_{xxxxx} residual.*

Status: [O] for $p \geq 2$ numerics (no harness yet; the bound's derivation is a consistency argument in OTS-NIDC.md §6). The $p=1$ row is [V] via Table 1.

1.2 Stability of Single Variable Systems

1.2.1 Setting and criterion

Set $u_j^n = G^n e^{ikj\Delta x}$. The three-point Laplacian has symbol

$$\delta_x^2 u_j^n \rightarrow -\frac{4}{\Delta x^2} s u_j^n, \quad s = \sin^2(k\Delta x/2) \in [0, 1].$$

Let $\mu = D\Delta t/\Delta x^2$, $\eta = \alpha\Delta x^2/D$ (α already contains σ^2 for the moment equations), and $a = \alpha\Delta t = \mu\eta$. The exact one-step factor is $E(s) = e^{-4\mu s + a}$, so $|E| \leq e^a$: for $\alpha > 0$ the equation grows, and $|G| \leq 1$ is the wrong requirement. We use the von Neumann condition with growth allowance,

$$|G(s)| \leq 1 + C \Delta t \quad \text{uniformly in } s \in [0, 1], \quad (8)$$

along refinement paths $\Delta t = \mu\Delta x^2/D$ with μ fixed (Lax–Richtmyer). The sharper, natural comparison is $|G(s)| \leq e^a$.

1.2.2 Amplification factors

We frame OTS-NIDC as a perturbation of the simple Finite-Euler method, so we begin by deriving the amplification factor G_{FE} and then transform it into G_{NIDC} as follows in,

$$G_{\text{FE}} = 1 - 4\mu s + \mu\eta, \quad (9)$$

$$G_{\text{NIDC}} = 1 - 4\mu s + \mu\eta - \frac{\eta}{9}s + \frac{\eta^2}{72}, \quad (10)$$

the latter from $C^{(1)}$ in (5): $\frac{\alpha\Delta x^4}{36D} \cdot (-\frac{4s}{\Delta x^2}) = -\frac{\eta}{9}s$ and $\frac{\alpha^2\Delta x^4}{72D^2} = \frac{\eta^2}{72}$. Both are affine in s with negative slope, so the extremes over $s \in [0, 1]$ are at $s=0$ (physical mode) and $s=1$ (grid mode).

phrase all
of this as
lemmas
please god

1.2.3 Stability at the optimal step

At $\Delta t = \Delta t_{\text{opt}}$, $\mu = 1/6$, $\eta = 6a$, and

$$G_{\text{NIDC}}(s) = \left(1 - \frac{2}{3}s\right)(1 + a) + \frac{a^2}{2}. \quad (11)$$

Compare $E(s) = e^{-2s/3}e^a$: the correction supplies exactly the $a^2/2$ term of e^a at $s = 0$, and $-\frac{2}{3}as$ is the cross term. Since $1 - \frac{2}{3}s \in [\frac{1}{3}, 1]$,

$$\frac{1}{3}(1 + a) + \frac{a^2}{2} \leq G_{\text{NIDC}}(s) \leq 1 + a + \frac{a^2}{2} \leq e^a \quad (a \geq 0).$$

Hence $G_{\text{NIDC}} > 0$ for all s (no oscillatory modes) and $|G_{\text{NIDC}}| \leq e^{\alpha\Delta t}$: (8) holds for every $\eta \geq 0$, with no restriction on η . The offset $+\eta^2/72$ is therefore not destabilizing; it is what makes $G(0)$ match e^a to $O(a^2)$. For plain FE at the same μ , $G_{\text{FE}}(s) = (1 - \frac{2}{3}s) + a$, also inside $[\frac{1}{3} + a, 1 + a]$.

1.2.4 Grid-scale boundary for general μ

Requiring the lower side of (8) at $s = 1$, i.e. $G(1) \geq -(1 + \mu\eta)$:

$$\mu_{\text{FE,max}} = \frac{2}{4 - 2\eta} = \frac{1}{2 - \eta}, \quad (12)$$

$$\mu_{\text{NIDC,max}} = \frac{2 - \eta/9 + \eta^2/72}{4 - 2\eta}, \quad (13)$$

with ratio

$$\frac{\mu_{\text{NIDC,max}}}{\mu_{\text{FE,max}}} = 1 - \frac{\eta}{18} + \frac{\eta^2}{144}. \quad (14)$$

(The stricter test $G(1) \geq -1$ gives denominators $4 - \eta$ and the same ratio.) At $\eta = 0.30$: $\mu_{\text{FE,max}} = 0.5882$, $\mu_{\text{NIDC,max}} = 0.5788$, ratio 0.98396 (-1.6%). The shrink is $O(\Delta x^2)$, and the effective NIDC CFL is $\Delta t \lesssim (1 - \eta/18) \Delta x^2 / (2D)$.

1.2.5 Scope of the correction ($\Delta t \neq \Delta t_{\text{opt}}$)

$C^{(1)}$ is derived at $\Delta t = \Delta t_{\text{opt}}$ and is independent of Δt . At $s = 0$,

$$G_{\text{NIDC}}(0) - e^a \approx \frac{\eta^2}{72}(1 - 36\mu^2),$$

which is ≤ 0 iff $\mu \geq 1/6$. For $\mu < 1/6$ the excess is Δt -independent, so at fixed Δx with $\Delta t \rightarrow 0$ the accumulated growth $(1 + \text{const})^{T/\Delta t}$ diverges. Stability statements therefore refer to refinement paths μ fixed, where the excess is $O(\Delta t^2)$ and (8) holds; the $\Delta t/\Delta t_{\text{opt}} \in \{0.5, \dots, 2\}$ table is a sweep over such paths, not a fixed- Δx study. The method is designed for $\mu = 1/6$.

1.2.6 Pair equations

Replace $D \rightarrow D_i + D_j$, $\alpha \rightarrow \alpha_{ij}$; (11) holds verbatim with $\mu_{ij} = 1/6$ at $\Delta t_{\text{opt}}^{(ij)}$. For the coupled discrete system there is still no independent proof (open item ??).

Theorem 6 (Von Neumann stability). *With $s = \sin^2(k\Delta x/2)$, $\mu = D\Delta t/\Delta x^2$, $\eta = \alpha\Delta x^2/D$, the amplification factors are $G_{\text{FE}} = 1 - 4\mu s + \mu\eta$ and*

$$G_{\text{NIDC}} = 1 - 4\mu s + \mu\eta - \frac{\eta}{9}s + \frac{\eta^2}{72}.$$

For the moment hierarchy $\eta \geq 0$ and the grid-scale stability bound tightens from $\mu \leq 2/(4 - \eta)$ to $\mu \leq (2 - \eta/9 + \eta^2/72)/(4 - \eta)$: a relative shrink of $1 - \eta/18 + O(\eta^2)$, i.e. $O(\Delta x^2)$, about -1.6% at the coarse $\eta = 0.3$ test point. The optimal step $\mu = 1/6$ sits a factor of 3 inside the CFL ceiling for both schemes.

Status: [V](analytic derivation above + numerical check in `notebooks/ai-src/stability/`; the earlier “expands the CFL” conclusion was an artifact of the pre-correction sign and has been fixed in the same release). The correction is stability-neutral to first order; it never threatens the optimal step. Note the honest asymmetry: this is a *scalar* analysis; the coupled discrete schemes of §2 have no independent stability proof (open item ??).

1.3 Efficiency profile

The method’s cost argument is architectural, not just arithmetic: versus multi-stage Runge–Kutta, OTS-NIDC is single-pass (one read, one write of the state vector; no stage storage, no stage barriers); versus compact schemes, it is strictly local (no tridiagonal solves), so it maps onto GPUs and bandwidth-bound hardware without global communication. At $N = 256$ the verified scalar runs show $\sim 10^3$ – $10^4 \times$ lower error than forward Euler at identical grid and step [V]— order wins over constants at refinement.

2 The Step to Coupled PDEs

We now couple species to allow for multi-linear systems, a direct expansion from the canon work we expand on!

2.1 Reaction-Diffusion Family and OTS

For N species with diagonal diffusion, linear reaction coupling, and (here) zero noise — the deterministic skeleton of the moment systems to come —

$$\partial_t u^{(i)} = D_i \partial_{xx} u^{(i)} + \sum_j R_{ij} u^{(j)}. \quad (15)$$

Lemma 7 (OTS-NIDC for fixed indices). *For equation (15), the optimal step for each $u^{(i)}$ individually, as derived in the single variable case, is exactly $\Delta t_{\text{opt}}^{(i)} = \frac{\Delta x^2}{6D_i}$, with first-level corrections $C^{(1),(i)}$ of the same algebraic form as (5), derived term-by-term in *OTS-NIDC.md* §8–9.*

With this, we get the desired theorem as follows.

Theorem 8 (First-level accuracy for fixed indices). *The scheme*

$$u_k^{(i),n+1} = u_k^{(i),n} + \Delta t_{\text{opt}} \left[D_i \delta_x^2 u_k^{(i),n} + \sum_j R_{ij} u_k^{(j),n} \right] + C^{(1),(i)} \quad (16)$$

has similar step error and global order as given by Theorem 4.

2.1.1 Competing Time Steps and Subcycling

Notice that the scheme presented by Theorem 8 relies on having $u_k^{(j),n}$ from having the value of separate PDE $u^{(j)}$ at time-step n . However, since each equation itself relies on their own time-step, we lose the guarantee of having the required value for any given equation readily available if $D_i = D, \forall i$ such that $\Delta t_{\text{opt}}^{(i)} = \Delta t_{\text{opt}}$ for the whole system.

The two failure modes of a naive single step frame the problem.

Failure 1: shared step without correction. If species 1 (fast, D_1) is forced onto species 2's optimum $\Delta t^* = \tau_2$, its Group-1 cancellation breaks. The one-step identity

$$e(\Delta t^*) = \Delta t^* D_1 \tilde{u}_{xxxx} \left[\frac{\Delta t^* D_1}{2} - \frac{\Delta x^2}{12} \right] + O((\Delta t^*)^3) \quad (17)$$

is exact for every Δt^* , and writing $\Delta t^* = \tau_1 + \epsilon$ turns the bracket into $\epsilon D_1/2$ *identically* (no Taylor truncation in ϵ). Hence:

Theorem 9 (Exact mismatch correction). *For any shared step $\Delta t^* = \tau_1 + \epsilon > 0$ that is stable for species 1, subtracting (17) (with $\tilde{u}_{xxxx}$ from a 5-point stencil) restores species 1's full accuracy. The correction is algebraic, not asymptotic, in ϵ — including ϵ of the same size as τ_1 itself.*

Status: [V](worked example $D_1 = 3, D_2 = 1$ at $r = 3$, `examples/test_regime1_D1.3_D2.1.py`; correction there is grid-independent — the Δx^4 cancels, leaving pure stencil weights).

Failure 2: naive splitting. Lie–Trotter splitting with the coupling frozen over a macro step H incurs a sustained $O(H)$ forcing error; accumulated over T/H cycles this is a global $O(H) = O(\Delta x^2)$ error — worse than the uncorrected base scheme. Freezing is not viable; coupling data must be interpolated (order $m \geq p$ by the Gear–Wells bookkeeping (see Thm. ??'s setting), cubic Hermite sufficing through $p = 2$ (Gear and Wells, 1984; Fok and Rosales, 2010)).

References

- Kevin T. Chu. Boosting the accuracy of finite difference schemes via optimal time step selection and non-iterative defect correction. *Applied Mathematics and Computation*, 218(7):3596–3614, 2011. doi: 10.1016/j.amc.2011.08.108. arXiv:0811.2847.
- Pak-Wing Fok and Robert R. Rosales. Multirate integration of axisymmetric step-flow equations. *arXiv preprint*, 2010. arXiv:0810.2517.
- C. William Gear and David R. Wells. Multirate linear multistep methods. *BIT Numerical Mathematics*, 24:484–502, 1984.