Optimal Time-Stepping with Non-Iterative Defect Correction for the Moment Hierarchy of Stochastic Reaction–Diffusion Equations

A living paper: derivation, verification ledger, and an honest map from closed moment systems to Cole–Hopf and KPZ observables

Aditya Tiwari

Living document, version 0.3 — September 16, 2026

Repository: `OTS-NIDC` (this paper is generated from it)

1 Review for Single Variable PDEs

Throughout, $\delta_x^2 u_j = (u_{j+1} - 2u_j + u_{j-1})/\Delta x^2$ is the centered second difference and refinement means $\Delta x \to 0$ with Δt slaved to Δx.

1.1 The scalar family and the optimal time step

Consider the deterministic family

$$u_t = D\,u_{xx} + \alpha\,u, \qquad D > 0,\ \alpha \in \mathbb{R}, \tag{1}$$

whose stochastic motivation (the moment equations of (**??**), with $\alpha = \frac{k(k-1)}{2}\sigma^2$) is deferred to §3. The pure diffusion case $\alpha = 0$ is exactly Chu's setting (Chu, 2011).

Apply forward Euler in time and the three-point stencil in space. Taylor expansion of both sides, followed by repeated differentiation of (1) to convert time derivatives of u into space derivatives (the operators $D\partial_{xx}$ and α commute, giving the binomial identity $u_{t^k} = (D\partial_{xx} + \alpha)^k u$), yields the local truncation error organized by derivative type:

$$\tau \;=\; \underbrace{\Big(\tfrac{D^2\Delta t}{2} - \tfrac{D\Delta x^2}{12}\Big)u_{xxxx}}_{\text{Group 1}} \;+\; \underbrace{\alpha D\,\Delta t\,u_{xx}}_{\text{Group 2}} \;+\; \underbrace{\tfrac{\alpha^2\Delta t}{2}\,u}_{\text{Group 3}} \;+\; O(\Delta t^2, \Delta x^4). \tag{2}$$

Only Group 1 couples Δt to Δx. Setting it to zero:

Theorem 1 (Optimal time step (Chu, 2011))**.** *For (1) discretized by forward Euler / centered differences, the unique step size annihilating the leading (u_{xxxx}) truncation term is*

$$\boxed{\Delta t_{\text{opt}} = \frac{\Delta x^2}{6D}} \tag{3}$$

and no single step annihilates both the u_{xxxx} and u_{xxxxxx} truncation terms, since the level-2 condition is $\Delta t = \Delta x^2/(\sqrt{60}\,D) \neq \Delta t_{\text{opt}}$.

(Proof: one-line cancellation in (2); uniqueness by the level-2 computation. Full algebra in `sign-change.tex` §1.)

Proposition 2 (α-independence)**.** *Δt_{opt} is independent of the reaction parameter α: the cancellation operates on the diffusive group alone. For the stochastic hierarchy this makes Δt_{opt} independent of the noise amplitude σ and of the moment order k.*

Following the derivation inspecting the truncation error, at $\Delta t = \Delta t_{\text{opt}}$ the Group-1 term vanishes and the surviving residual is

$$\tau\big|_{\Delta t_{\text{opt}}} = \frac{\alpha \Delta x^2}{6} u_{xx} + \frac{\alpha^2 \Delta x^2}{12 D} u + O(\Delta x^4). \tag{4}$$

Then the truncation error τ as assembled in (2) is the *forward* residual: exact − scheme = $\Delta t\, \tau + \cdots$. Cancelling it therefore requires *adding* $\Delta t\, \tau$.

Lemma 3 (Closed form correction term)**.** *The corrected factor at any given time step is,*

$$C_j^{(1)} = +\frac{\alpha\, \Delta x^4}{36 D}\, \delta_x^2 u_j^n + \frac{\alpha^2\, \Delta x^4}{72 D^2}\, u_j^n. \tag{5}$$

Theorem 4 (First-level accuracy)**.** *The corrected scheme*

$$u_j^{n+1} = u_j^n + \Delta t_{\text{opt}} \left[D\, \delta_x^2 u_j^n + \alpha\, u_j^n \right] + C_j^{(1)} \tag{6}$$

has step error $O(\Delta x^6)$ and global order 4: the correction $C^{(1)}$ is itself $O(\Delta x^4)$, so its evaluation with the second-order operator δ_x^2 contributes only $O(\Delta x^6)$ to the step — the three-point stencil is mathematically sufficient at $p = 1$.

Status: [V]. Measured orders and errors, with the harness described in Appendix **??**:

Table 1: Scalar $p = 1$ convergence, $u_t = D u_{xx} + \alpha u$ with $D = 0.1$, $\alpha\sigma^2 = 0.25$, Neumann BCs, $T = 0.05$; L^2 error against the exact cosine-series solution. Data: `code/implementation/results.md` (generated by `convergence_test.py`, 2026-07). All methods use the same grid and step size Δt_{opt} at each level.

Method	$N = 32$	$N = 64$	$N = 128$	$N = 256$
OTS-NIDC ($p = 1$)	8.54e-09	5.26e-10	3.27e-11	2.04e-12
Forward Euler	1.23e-05	3.02e-06	7.52e-07	1.88e-07
4th-order FD + FE	1.52e-05	3.79e-06	9.45e-07	2.36e-07
Reference expm	2.77e-05	6.81e-06	1.70e-06	4.23e-07
Measured order (log–log slope, mean over levels):				
OTS-NIDC	4.01	(FE / 4th-order FD / expm: 2.00–2.01)		

The 4th-order-FD row carries the paper's cleanest negative control: high-order spatial accuracy alone buys nothing when the step size is slaved to Δx^2 — the $O(\Delta t)$ time error caps the whole scheme at second order. The boost must come from the time-step structure; that is the method's thesis, verified.

The p-level tower and the stencil bound

move this into an appendix, perhaps?

The general correction is generated by powers of the operator, with the sign of Remark **??**:

$$C_j^{(\le p)} \;=\; +\sum_{\ell=1}^{p} \frac{\Delta t_{\mathrm{opt}}^{\ell}}{(\ell+1)!}\big(D\mathcal{D}_2+\alpha\big)^{\ell} u_j^n, \tag{7}$$

where $\mathcal{D}_2$ is a spatial operator whose accuracy must be raised as ℓ grows. Spatial discretization error breaks the binomial symmetry of the time expansion, giving the stencil-scaling bound:

Proposition 5 (Stencil bound; sketch)**.** *Global order $2p+2$ requires corrections evaluated on at least a $(2p+3)$-point stencil: $p=1$ needs the 3-point main stencil plus a 5-point evaluation inside $C^{(1)}$ when full fourth-order spatial accuracy is demanded of the correction itself; $p=2$ needs a 7-point stencil to represent the uncancelled u_{xxxxxx} residual.*

Status: [O]for $p \ge 2$ numerics (no harness yet; the bound's derivation is a consistency argument in `OTS-NIDC.md` §6). The $p=1$ row is [V]via Table 1.

1.2 Stability of Single Variable Systems

phrase all of this as lemmas please god

1.2.1 Setting and criterion

Set $u_j^n = G^n e^{ikj\Delta x}$. The three-point Laplacian has symbol

$$\delta_x^2 u_j^n \;\to\; -\frac{4}{\Delta x^2}\, s\, u_j^n, \qquad s=\sin^2(k\Delta x/2)\in[0,1].$$

Let $\mu = D\Delta t/\Delta x^2$, $\eta = \alpha\Delta x^2/D$ (α already contains σ^2 for the moment equations), and $a = \alpha\Delta t = \mu\eta$. The exact one-step factor is $E(s)=e^{-4\mu s+a}$, so $|E|\le e^a$: for $\alpha>0$ the equation grows, and $|G|\le 1$ is the wrong requirement. We use the von Neumann condition with growth allowance,

$$|G(s)| \le 1 + C\,\Delta t \quad \text{uniformly in } s\in[0,1], \tag{8}$$

along refinement paths $\Delta t = \mu\Delta x^2/D$ with μ fixed (Lax–Richtmyer). The sharper, natural comparison is $|G(s)|\le e^a$.

1.2.2 Amplification factors

We frame OTS-NIDC as a perturbation of the simple Finite-Euler method, so we begin by deriving the amplification factor G_{FE} and then transform it into G_{NIDC} as follows in,

$$G_{\mathrm{FE}} = 1 - 4\mu s + \mu\eta, \tag{9}$$

$$G_{\mathrm{NIDC}} = 1 - 4\mu s + \mu\eta - \tfrac{\eta}{9}s + \tfrac{\eta^2}{72}, \tag{10}$$

the latter from $C^{(1)}$ in (5): $\frac{\alpha\Delta x^4}{36D}\cdot(-\frac{4s}{\Delta x^2}) = -\frac{\eta}{9}s$ and $\frac{\alpha^2\Delta x^4}{72D^2} = \frac{\eta^2}{72}$. Both are affine in s with negative slope, so the extremes over $s\in[0,1]$ are at $s=0$ (physical mode) and $s=1$ (grid mode).

1.2.3 Stability at the optimal step

At $\Delta t = \Delta t_{\text{opt}}$, $\mu = 1/6$, $\eta = 6a$, and

$$G_{\text{NIDC}}(s) = \left(1 - \tfrac{2}{3}s\right)(1 + a) + \tfrac{a^2}{2}. \tag{11}$$

Compare $E(s) = e^{-2s/3}e^a$: the correction supplies exactly the $a^2/2$ term of e^a at $s = 0$, and $-\frac{2}{3}as$ is the cross term. Since $1 - \frac{2}{3}s \in [\frac{1}{3}, 1]$,

$$\tfrac{1}{3}(1 + a) + \tfrac{a^2}{2} \;\le\; G_{\text{NIDC}}(s) \;\le\; 1 + a + \tfrac{a^2}{2} \;\le\; e^a \qquad (a \ge 0).$$

Hence $G_{\text{NIDC}} > 0$ for all s (no oscillatory modes) and $|G_{\text{NIDC}}| \le e^{\alpha \Delta t}$: (8) holds for every $\eta \ge 0$, with no restriction on η. The offset $+\eta^2/72$ is therefore not destabilizing; it is what makes $G(0)$ match e^a to $O(a^2)$. For plain FE at the same μ, $G_{\text{FE}}(s) = (1 - \frac{2}{3}s) + a$, also inside $[\,\frac{1}{3} + a,\, 1 + a\,]$.

1.2.4 Grid-scale boundary for general μ

Requiring the lower side of (8) at $s = 1$, i.e. $G(1) \ge -(1 + \mu\eta)$:

$$\mu_{\text{FE,max}} = \frac{2}{4 - 2\eta} = \frac{1}{2 - \eta}, \tag{12}$$

$$\mu_{\text{NIDC,max}} = \frac{2 - \eta/9 + \eta^2/72}{4 - 2\eta}, \tag{13}$$

with ratio

$$\frac{\mu_{\text{NIDC,max}}}{\mu_{\text{FE,max}}} = 1 - \frac{\eta}{18} + \frac{\eta^2}{144}. \tag{14}$$

(The stricter test $G(1) \ge -1$ gives denominators $4 - \eta$ and the same ratio.) At $\eta = 0.30$: $\mu_{\text{FE,max}} = 0.5882$, $\mu_{\text{NIDC,max}} = 0.5788$, ratio 0.98396 ($-1.6\%$). The shrink is $O(\Delta x^2)$, and the effective NIDC CFL is $\Delta t \lesssim (1 - \eta/18)\,\Delta x^2/(2D)$.

1.2.5 Scope of the correction ($\Delta t \neq \Delta t_{\text{opt}}$)

$C^{(1)}$ is derived at $\Delta t = \Delta t_{\text{opt}}$ and is independent of Δt. At $s = 0$,

$$G_{\text{NIDC}}(0) - e^a \;\approx\; \frac{\eta^2}{72}\left(1 - 36\mu^2\right),$$

which is ≤ 0 iff $\mu \ge 1/6$. For $\mu < 1/6$ the excess is Δt-independent, so at fixed Δx with $\Delta t \to 0$ the accumulated growth $(1 + \text{const})^{T/\Delta t}$ diverges. Stability statements therefore refer to refinement paths μ fixed, where the excess is $O(\Delta t^2)$ and (8) holds; the $\Delta t/\Delta t_{\text{opt}} \in \{0.5, \ldots, 2\}$ table is a sweep over such paths, not a fixed-Δx study. The method is designed for $\mu = 1/6$.

1.2.6 Pair equations

Replace $D \to D_i + D_j$, $\alpha \to \alpha_{ij}$; (11) holds verbatim with $\mu_{ij} = 1/6$ at $\Delta t_{\text{opt}}^{(ij)}$. For the coupled discrete system there is still no independent proof (open item **??**).

Theorem 6 (Von Neumann stability)**.** *With $s = \sin^2(k\Delta x/2)$, $\mu = D\Delta t/\Delta x^2$, $\eta = \alpha\Delta x^2/D$, the amplification factors are $G_{\text{FE}} = 1 - 4\mu s + \mu\eta$ and*

$$G_{\text{NIDC}} = 1 - 4\mu s + \mu\eta - \tfrac{\eta}{9}s + \tfrac{\eta^2}{72}.$$

For the moment hierarchy $\eta \geq 0$ and the grid-scale stability bound tightens from $\mu \leq 2/(4-\eta)$ to $\mu \leq (2-\eta/9+\eta^2/72)/(4-\eta)$: a relative shrink of $1-\eta/18+O(\eta^2)$, i.e. $O(\Delta x^2)$, about -1.6% at the coarse $\eta = 0.3$ test point. The optimal step $\mu = 1/6$ sits a factor of 3 inside the CFL ceiling for both schemes.

Status: [V](analytic derivation above + numerical check in `notebooks/ai-src/stability/`; the earlier "expands the CFL" conclusion was an artifact of the pre-correction sign and has been fixed in the same release). The correction is stability-neutral to first order; it never threatens the optimal step. Note the honest asymmetry: this is a *scalar* analysis; the coupled discrete schemes of §2 have no independent stability proof (open item **??**).

1.3 Efficiency profile

The method's cost argument is architectural, not just arithmetic: versus multi-stage Runge–Kutta, OTS-NIDC is single-pass (one read, one write of the state vector; no stage storage, no stage barriers); versus compact schemes, it is strictly local (no tridiagonal solves), so it maps onto GPUs and bandwidth-bound hardware without global communication. At $N = 256$ the verified scalar runs show $\sim 10^3$–$10^4\times$ lower error than forward Euler at identical grid and step [V]— order wins over constants at refinement.

2 The Step to Coupled PDEs

We now couple species to allow for multi-linear systems, a direct expansion from the canon work we expand on!

2.1 Reaction-Diffusion Family and OTS

For N species with diagonal diffusion, linear reaction coupling, and (here) zero noise — the deterministic skeleton of the moment systems to come —

$$\partial_t u^{(i)} = D_i\, \partial_{xx} u^{(i)} + \sum_j R_{ij} u^{(j)}. \tag{15}$$

Lemma 7 (OTS-NIDC for fixed indices)**.** *For equation (15), the optimal step for each $u^{(i)}$ individually, as derived in the single variable case, is exactly $\Delta t^{(i)}_{\text{opt}} = \frac{\Delta x^2}{6D_i}$, with first-level corrections $C^{(1),(i)}$ of the same algebraic form as (5), derived term-by-term in `OTS-NIDC.md` §8–9.*

With this, we get the desired theorem as follows.

Theorem 8 (First-level accuracy for fixed indices)**.** *The scheme*

$$u_k^{(i),n+1} = u_k^{(i),n} + \Delta t_{\text{opt}} \left[D_i \delta_x^2 u_k^{(i),n} + \sum_j R_{ij} u_k^{(j),n} \right] + C^{(1),(i)} \tag{16}$$

has similar step error and global order as given by Theorem 4.

2.1.1 Competing Time Steps and Subcycling

Notice that the scheme presented by Theorem 8 relies on having $u_k^{(j),n}$ from having the value of separate PDE $u^{(j)}$ at time-step n. However, since each equation itself relies on their own time-step, we lose the guarantee of having the required value for any given equation readily available if $D_i = D, \forall i$ such that $\Delta t_{\text{opt}}^{(}i) = \Delta t_{\text{opt}}$ for the whole system.

The two failure modes of a naive single step frame the problem.

Failure 1: shared step without correction. If species 1 (fast, D_1) is forced onto species 2's optimum $\Delta t^* = \tau_2$, its Group-1 cancellation breaks. The one-step identity

$$e(\Delta t^*) \;=\; \Delta t^* D_1\, \tilde{u}_{xxxx}\left[\tfrac{\Delta t^* D_1}{2} - \tfrac{\Delta x^2}{12}\right] + O((\Delta t^*)^3) \tag{17}$$

is exact for every Δt^*, and writing $\Delta t^* = \tau_1 + \epsilon$ turns the bracket into $\epsilon D_1/2$ *identically* (no Taylor truncation in ϵ). Hence:

Theorem 9 (Exact mismatch correction)**.** *For any shared step $\Delta t^* = \tau_1 + \epsilon > 0$ that is stable for species 1, subtracting (17) (with $\tilde{u}_{xxxx}$ from a 5-point stencil) restores species 1's full accuracy. The correction is algebraic, not asymptotic, in ϵ — including ϵ of the same size as τ_1 itself.*

Status: [V](worked example $D_1 = 3, D_2 = 1$ at $r = 3$, `examples/test_regime1_D1_3_D2_1.py`; correction there is grid-independent — the Δx^4 cancels, leaving pure stencil weights).

We remain, however, that this stencil is not ideal for each step, and it leads to questions of stability. In fact,

Failure 2: naive splitting. Lie–Trotter splitting with the coupling frozen over a macro step H incurs a sustained $O(H)$ forcing error; accumulated over T/H cycles this is a global $O(H) = O(\Delta x^2)$ error — worse than the uncorrected base scheme. Freezing is not viable; coupling data data must be interpolated.

Therefore, we introduce a subcycle protocol that fares to match timesteps where information can nicely be interpolated. First we find that,

Theorem 10 (Regime boundary)**.** *Forcing the fast species onto the slow species' optimal step $\Delta t^* = \tau_2 = \Delta x^2/6D_2$ is stable for the fast species iff*

$$r := D_1/D_2 \;\le\; 3.$$

For $r > 3$ the shared step is unconditionally unstable and subcycling is mandatory.

Proof sketch. FTCS stability needs $\Delta t^* \le \Delta x^2/2D_1$; substituting $\Delta t^* = \Delta x^2/6D_2$ gives $r \le 3$. The threshold is grid-independent because both sides scale as Δx^2. □

Status: [S]— derived from plain FTCS; the corrected scheme's stability region (Prop. 6 is scalar-only) has not been re-derived, so the boundary is supported, not proven, for the corrected coupled scheme (open item **??**).

Then, via this regime boundary, we can strictly define the two regimes:

Regime 1: Single unoptimal step with exact mismatch correction. For any two equations with distinct $\Delta t_{\text{opt}}^{(1)} < \Delta t_{\text{opt}}^{(2)}$ and $r = D_2/D_1 \le 3$, we evolve both equations by $\Delta t_{\text{opt}}^* = \Delta t_{\text{opt}}^{(2)}$ such that $u^(2)$ is optimally solved by nature, and the discretization of $u^{(1)}$ is evolved with $\Delta t_{\text{opt}}^{(1)} + \epsilon$ such that the explicit mismatch correction term recovers order 4 accuracy.

Regime 2: Sub-cycling with Hermite Polynomial interpolation. For any two equations with distinct $\Delta t^{(1)}_{\text{opt}} < \Delta t^{(2)}_{\text{opt}}$ and $r = D_2/D_1 > 3$, evolve the fast equation $u^{(1)}$ via $n = \lfloor r \rfloor$ sub-cycled steps of $\Delta t^{(1)}_{\text{opt}}$ and a single remainder step $\rho = \Delta t^{(1)}_{\text{opt}} - n\Delta t^{(2)}_{\text{opt}}$ with explicit mismatch correction given by ϵ in $\rho = \Delta t^{(2)}_{\text{opt}} - \epsilon$. The slow equation $u^{(2)}$ can evolve with it's own $\Delta t^{(2)}_{\text{opt}}$, but requires cubic Hermite interpolation to overcome the $O(H)$ forcing error through $p = 2$ (Gear and Wells, 1984; Fok and Rosales, 2010)).

With these regimes, we complete the derivation of OTS-NIDC for coupled PDEs.

2.2 Stability

LOOOL, i totally believe in myself

3 The method III: stochastic reaction–diffusion via the moment hierarchy

Everything so far is deterministic, but our greatest contribution is primarily in the stochastic domain. The stochastic payload enters through a single structural fact: *for the right noise class, the SPDE's moments obey closed deterministic reaction–diffusion equations* — the exact family (1) – (15) of §1–2.

3.1 Moment systems for the linear N-species model

hOOOOOLI idek

Setting. Let $D \subset \mathbb{R}$ be the periodic interval $\mathbb{R}/L\mathbb{Z}$ or a bounded interval with Neumann conditions, and let Δ be the corresponding Laplacian on $H = L^2(D)$. Let $(e_m)_{m\geq 0}$ be its real orthonormal eigenbasis, $-\Delta e_m = \lambda_m e_m$, $\lambda_m \geq 0$.

The noise has spatial covariance $q : D \times D \to \mathbb{R}$, which is symmetric, continuous and positive semidefinite. By Mercer's theorem, $q = \sum_p \mu_p\, f_p \otimes f_p$ with (f_p) an orthonormal basis of H, $\mu_p \geq 0$, and $\sum_p \mu_p < \infty$. Put $\bar{q} = \sup |q| < \infty$. Let

$$W_k(t) = \sum_p \sqrt{\mu_p}\, \beta^p_k(t)\, f_p,$$

where the β^p_k are independent standard Brownian motions, so the channels k are independent with common spatial covariance q. Consider

$$du_i = \Big(D_i \Delta u_i + \sum_j R_{ij} u_j\Big) dt + \sum_k \sigma_{ik}\, u_k\, dW_k, \qquad u_i(0) = u^0_i, \tag{18}$$

with $D_i \geq 0$, R, σ constant matrices, and u^0 independent of W with $\mathbb{E}\|u^0\|^4_H < \infty$.

Hypothesis (H). (18) has a unique mild solution $u \in C([0,T]; H)$ a.s. with $\sup_{t\leq T} \mathbb{E}\|u(t)\|^4_H < \infty$.

(H) holds under the assumptions above. The map $v \mapsto \sum_k \sigma_{ik}\, v\, dW_k$ is Lipschitz $H \to \mathrm{HS}(Q^{1/2}H, H)$, because $\sum_p \mu_p \|v f_p\|^2 = \int q(x,x) v(x)^2 dx \leq \bar{q}\|v\|^2$. Existence, uniqueness and moment bounds then follow from the standard theory (Da Prato–Zabczyk, *Stochastic Equations in Infinite Dimensions*, Thm. 7.2 and its p-th moment version).

Moment objects. Define the Bochner expectations

$$m_i(t) = \mathbb{E}[u_i(t)] \in L^2(D), \qquad V_{ij}(t) = \mathbb{E}\big[u_i(t) \otimes u_j(t)\big] \in L^2(D \times D),$$

so that formally $V_{ij}(t, x, y) = \mathbb{E}[u_i(t, x)\, u_j(t, y)]$. These are well defined because $\|u_i \otimes u_j\| = \|u_i\|\|u_j\|$ is integrable. The pointwise ("same-point") moment is the restriction $V_{ij}(t, x, x)$, which is *not* an object of this proposition.

Proposition 11 (Closed moment systems)**.** *Under (H):*

(a) ***Mean.*** *$m = (m_i)$ is the unique mild solution of*

$$\partial_t m_i = D_i \Delta m_i + \sum_j R_{ij} m_j. \tag{19}$$

(b) ***Two-point function.*** *$V = (V_{ij})$ is the unique mild solution in $C([0, T]; L^2(D \times D)^{N \times N})$ of*

$$\begin{aligned} \partial_t V_{ij}(x, y) = \big(D_i \Delta_x + D_j \Delta_y\big) V_{ij}(x, y) + \sum_k \big(R_{ik} V_{kj}(x, y) + R_{jk} V_{ik}(x, y)\big) \\ + \; q(x, y) \sum_k \sigma_{ik} \sigma_{jk}\, V_{kk}(x, y), \end{aligned} \tag{20}$$

with $V_{ij}(0) = \mathbb{E}[u_i^0 \otimes u_j^0]$. Here Δ_x, Δ_y carry the boundary condition of Δ in each variable separately; in the Neumann case this is $\partial_{n_x} V_{ij} = \partial_{n_y} V_{ij} = 0$, in the weak sense.

In particular, (20) is a closed, linear, deterministic system for (V_{ij}): the noise source involves only the two-point functions $V_{kk}(x, y)$ and the covariance $q(x, y)$.

Status: ***[P]****, proved below modulo the cited well-posedness (H). The weak-sense Neumann condition on V is also a consequence of the proof (it settles the item previously marked "used but not proven").*

this is claude, i believe goes without saying

Proof. **Step 1 (mode equations).** Let $u_i^m(t) = \langle u_i(t), e_m \rangle$. Testing the mild formulation against e_m, and using that Δ is self-adjoint with $\Delta e_m = -\lambda_m e_m$, gives

$$u_i^m(t) = u_i^m(0) + \int_0^t \Big(- D_i \lambda_m u_i^m + \sum_j R_{ij} u_j^m \Big) ds + M_i^m(t),$$

$$M_i^m(t) = \sum_k \sigma_{ik} \sum_p \sqrt{\mu_p} \int_0^t \langle e_m, u_k f_p \rangle \, d\beta_k^p.$$

The β_k^p are independent across (k, p), so

$$d\langle M_i^m, M_j^n \rangle_t = \sum_k \sigma_{ik} \sigma_{jk} \sum_p \mu_p \langle e_m, u_k f_p \rangle \langle e_n, u_k f_p \rangle \, dt.$$

By Mercer's theorem the series $\sum_p \mu_p f_p(x) f_p(y)$ converges uniformly to $q(x, y)$, and $u_k \in L^2 \subset L^1$, so we may interchange sum and integral:

$$d\langle M_i^m, M_j^n \rangle_t = \sum_k \sigma_{ik} \sigma_{jk} \iint e_m(x) e_n(y)\, q(x, y)\, u_k(t, x) u_k(t, y)\, dx\, dy\, dt. \tag{21}$$

Also $\sum_p \mu_p \langle e_m, u f_p \rangle^2 = \iint e_m(x) e_m(y)\, q(x, y)\, u(x) u(y) \le \bar{q} \|u\|^2$, so $d\langle M_i^m \rangle \le C \bar{q} \|u\|^2 dt$.

Step 2 (Itô product and expectation). By the Itô product rule for continuous semimartingales,

$$d\big(u_i^m u_j^n\big) = u_i^m\, du_j^n + u_j^n\, du_i^m + d\langle M_i^m, M_j^n\rangle.$$

The stochastic integrals $\int u_i^m\, dM_j^n$ are true martingales: since $|u_i^m| \le \|u_i\|$ and $d\langle M_j^n\rangle \le C\bar{q}\|u_j\|^2 dt$,

$$\mathbb{E}\int_0^T (u_i^m)^2\, d\langle M_j^n\rangle \le C\bar{q}\int_0^T \mathbb{E}\big[\|u_i\|^2\|u_j\|^2\big]dt < \infty$$

by the fourth-moment bound in (H). Taking expectations, writing $V_{ij}^{mn} = \mathbb{E}[u_i^m u_j^n] = \langle V_{ij}, e_m \otimes e_n\rangle$, and applying Fubini in (21) (the integrand is dominated by $\bar{q}\,|e_m(x)e_n(y)|\,\mathbb{E}[|u_k(x)||u_k(y)|]$, which is integrable), we get

$$\frac{d}{dt}V_{ij}^{mn} = -(D_i\lambda_m + D_j\lambda_n)V_{ij}^{mn} + \sum_k \big(R_{ik}V_{kj}^{mn} + R_{jk}V_{ik}^{mn}\big) + \sum_k \sigma_{ik}\sigma_{jk}\,\big\langle q\,V_{kk},\, e_m \otimes e_n\big\rangle. \quad (22)$$

The right side depends only on the family $\{V_{kl}\}$, not on any higher moment. This is the closure.

Step 3 (identification and uniqueness). Let $X = L^2(D \times D)^{N\times N}$. Let B act on the (i,j) component as $D_i\Delta_x + D_j\Delta_y$ with the stated boundary conditions. It is self-adjoint and nonpositive, diagonal in the basis $e_m \otimes e_n$ with eigenvalue $-(D_i\lambda_m + D_j\lambda_n)$, hence generates a contraction C_0-semigroup on X; the boundary conditions on V are those of the domain of B. Let $\mathcal{L}$ collect the remaining terms of (20). It is bounded on X:

$$\|(\mathcal{L}v)_{ij}\| \le \sum_k \big(|R_{ik}|\|v_{kj}\| + |R_{jk}|\|v_{ik}\|\big) + \bar{q}\sum_k |\sigma_{ik}\sigma_{jk}|\,\|v_{kk}\|,$$

because multiplication by the bounded function q is bounded on $L^2(D \times D)$. So $B + \mathcal{L}$ generates a C_0-semigroup on X.

The map $t \mapsto V(t) \in X$ is continuous, by dominated convergence from continuity of u in H and the fourth-moment bound. By (22), V is a weak solution of $V' = (B + \mathcal{L})V$ tested against the basis $e_m \otimes e_n$, which is dense in $D(B)$ in graph norm. A continuous weak solution coincides with the mild solution (Ball, 1977), and the mild solution is unique. This proves (b).

Part (a) is identical with the noise term absent: take expectations in Step 1, using $\mathbb{E}M_i^m = 0$, to get $\frac{d}{dt}m_i^m = -D_i\lambda_m m_i^m + \sum_j R_{ij}m_j^m$. □

Remark 12 (Why the pointwise equation does not close)**.** The former Prop. 12 asserted $\partial_t V_{ij}(x,x) = (D_i + D_j)\partial_{xx}V_{ij}(x,x) + \cdots$ for the same-point function. Set $v_{ij}(x) = V_{ij}(x,x)$. The chain rule gives

$$\partial_{xx}\, v_{ij} = \big(\Delta_x + \Delta_y + 2\partial_x\partial_y\big)V_{ij}\Big|_{y=x}.$$

For $D_i = D_j = D$ the diffusion term in (20), restricted to the diagonal, is therefore

$$D\,\partial_{xx}v_{ij} \;-\; 2D\,\mathbb{E}\big[\partial_x u_i\,\partial_x u_j\big](x),$$

which contains the gradient moment $\mathbb{E}[\partial_x u_i\,\partial_x u_j]$. That moment is not a function of v_{ij}; its own evolution involves higher derivatives of V, so the pointwise hierarchy does not close. For $D_i \neq D_j$ the diagonal diffusion term cannot even be written as a multiple of $\partial_{xx}v_{ij}$ plus a gradient term.

Deterministic counterexample ($\sigma = 0$, $N = 1$, $R = 0$). Take $u_t = Du_{xx}$ and $u_0 = 2 + \cos x$, so $V(0,x,y) = u_0(x)u_0(y)$. At $x = \pi/2$ we have $u_0 = 2$, $u_{0,x} = -1$, $u_{0,xx} = 0$. Then

$$\partial_t V(x,x)\big|_{t=0} = 2D\,u_0u_{0,xx} = 0, \qquad D\,\partial_{xx}\big[V(x,x)\big]\big|_{t=0} = 2D\big(u_{0,x}^2 + u_0u_{0,xx}\big) = 2D.$$

The same-point equation fails even for heat flow.

Stationary check. If $V_{ii}(x,y) = c(x-y)$ is homogeneous, then v_{ii} is constant, so the pointwise equation predicts no diffusive damping of the variance. The two-point equation instead gives a diffusive contribution $2D\,c''(0) = -2D\,\mathbb{E}[(\partial_x u_i)^2] < 0$ on the diagonal, as it should.

Remark 13 (Cases where the diagonal does close)**.** If $D_i = 0$ for all i (no spatial coupling), then (20) contains no spatial derivatives, so the diagonal restriction (with $q(x,x)$ in the source) closes and reduces to the per-cell finite-dimensional Itô computation. The failure is entirely due to the diffusion term, which couples a point to its neighbours. The mean system (19) closes for any noise, because the equation is linear in u.

Remark 14 (Consequences for OTS–NIDC (status [O]))**.** (20) is a 2-dimensional linear reaction–diffusion system in (x,y) with a potential-like coupling $q(x,y)$. This has three consequences.

- The scalar analysis of §1 applies to (20) only after extension to two space dimensions. The squared Laplacian $(D_i\Delta_x + D_j\Delta_y)^2$ contains a mixed term $2D_iD_j\,\partial_{xxyy}$, which the one-dimensional cancellation at Δt_{opt} does not remove; an additional cross-derivative correction is required.
- $q(x,y)$ is a variable coefficient, so $D_i\Delta$ and the reaction term do not commute; the binomial identity behind Eq. (2) fails and Group 2 and Group 3 pick up derivatives of q.
- The cost scales as N_x^2 for the two-point function, so the efficiency comparison against Euler–Maruyama must be restated.

For white noise ($q = \delta(x-y)$) the source is $\delta(x-y)\sum_k \sigma_{ik}\sigma_{jk}V_{kk}(x,x)$; the equation is plausible in the distributional sense in one dimension, but this proof does not cover it (it needs $\bar{q} < \infty$), and V then has a gradient jump on the diagonal, which affects the Taylor-based order claims. This is open.

Remark 15 (Notation)**.** The former text wrote M_i in (19); use m_i for means and reserve M_k for nothing. The old Prop. 11 is subsumed by (a) for $k = 1$; it is retracted for $k \geq 2$.

decide which proof is actually worth to keep

Alternative proof of Proposition 11. Write $\mathcal{D}$ for the spatial domain and $H = L^2(\mathcal{D})$. All expectations and integrals below are justified by hypothesis (H), the boundedness of q, and the Cauchy–Schwarz inequality. We derive the equations in weak form, so no pointwise differentiability of the random field is assumed.

Step 1: The mean equation. Fix a species i and a smooth test function φ satisfying the periodic or homogeneous Neumann boundary condition. Testing (18) against φ gives

$$\langle u_i(t),\varphi\rangle = \langle u_i(0),\varphi\rangle + \int_0^t \left[D_i\langle u_i(s),\Delta\varphi\rangle + \sum_k R_{ik}\langle u_k(s),\varphi\rangle \right] ds$$
$$+ \sum_{k,p} \sigma_{ik}\sqrt{\mu_p} \int_0^t \langle u_k(s) f_p,\varphi\rangle\, d\beta_{kp}(s).$$

The last line is an integrable, mean-zero stochastic integral: indeed, its second moment is bounded by a constant multiple of $\int_0^t \mathbb{E}\|u(s)\|_H^2\,ds$, since $\sum_p \mu_p\|u_k f_p\|_H^2 = \int_{\mathcal{D}} q(x,x)|u_k(x)|^2\,dx \leq \bar{q}\|u_k\|_H^2$. Taking expectations and setting $m_i = \mathbb{E}[u_i]$ yields

$$\langle m_i(t),\varphi\rangle = \langle m_i(0),\varphi\rangle + \int_0^t \left[D_i\langle m_i(s),\Delta\varphi\rangle + \sum_k R_{ik}\langle m_k(s),\varphi\rangle \right] ds.$$

This is the weak form of (19).

Step 2: The two-point product. Fix species i, j and smooth test functions φ, ψ as above. Introduce the real-valued processes

$$X(t) = \langle u_i(t), \varphi \rangle, \qquad Y(t) = \langle u_j(t), \psi \rangle.$$

They satisfy ordinary real-valued stochastic integral equations. For any increments the algebraic identity

$$\Delta(XY) = X\,\Delta Y + Y\,\Delta X + \Delta X\,\Delta Y$$

is exact. The finite-dimensional Itô product rule is the rigorous continuous-time version of this identity: the product of the two noise increments contributes their covariance, while drift–noise and drift–drift products vanish at first order.

For each channel k and covariance mode p, the noise coefficients of X and Y are, respectively,

$$a_{kp} = \sigma_{ik}\sqrt{\mu_p}\langle u_k f_p, \varphi \rangle, \qquad b_{kp} = \sigma_{jk}\sqrt{\mu_p}\langle u_k f_p, \psi \rangle.$$

Independence of the β_{kp} implies that the covariance term in the expected product increment is $\mathbb{E}\sum_{k,p} a_{kp} b_{kp}\,dt$. The expansion $q(x,y) = \sum_p \mu_p f_p(x) f_p(y)$ therefore gives

$$\begin{aligned}\sum_{k,p} a_{kp} b_{kp} &= \sum_k \sigma_{ik}\sigma_{jk} \sum_p \mu_p \left(\int_{\mathcal{D}} u_k(x) f_p(x) \varphi(x)\,dx \right) \left(\int_{\mathcal{D}} u_k(y) f_p(y) \psi(y)\,dy \right) \\ &= \sum_k \sigma_{ik}\sigma_{jk} \iint_{\mathcal{D}^2} q(x,y) u_k(x) u_k(y) \varphi(x) \psi(y)\,dx\,dy.\end{aligned}$$

The interchange of summation and integration is justified by the covariance-operator bound and $\bar{q} < \infty$; equivalently it follows by taking limits of finite covariance-mode sums in L^1.

Step 3: Taking expectations. The deterministic part of the product rule contributes

$$\begin{aligned}&D_i\,\mathbb{E}g[\langle u_i, \Delta\varphi \rangle \langle u_j, \psi \rangle] + D_j\,\mathbb{E}\big[\langle u_i, \varphi \rangle \langle u_j, \Delta\psi \rangle\big] \\ &\quad + \sum_k R_{ik}\,\mathbb{E}\big[\langle u_k, \varphi \rangle \langle u_j, \psi \rangle\big] + \sum_k R_{jk}\,\mathbb{E}\big[\langle u_i, \varphi \rangle \langle u_k, \psi \rangle\big].\end{aligned}$$

The remaining stochastic integrals have zero expectation. For example, their square-integrability follows from (H), because $|X|^2 \le \|\varphi\|_H^2 \|u_i\|_H^2$ and the variance rate of the noise in Y is bounded by $C\bar{q}\|\psi\|_H^2 \sum_k \|u_k\|_H^2$. Thus their expected quadratic integrals are controlled by $C\int_0^T \mathbb{E}\|u(s)\|_{H^N}^4\,ds < \infty$.

By definition of $V_{ij} = \mathbb{E}[u_i \otimes u_j]$, combining these terms gives the integral identity

$$\begin{aligned}\langle V_{ij}(t), \varphi \otimes \psi \rangle = \langle V_{ij}(0), \varphi \otimes \psi \rangle &\\ + \int_0^t \Big\{ D_i \langle V_{ij}, \Delta\varphi \otimes \psi \rangle + D_j \langle V_{ij}, \varphi \otimes \Delta\psi \rangle &\\ + \sum_k \big(R_{ik} \langle V_{kj}, \varphi \otimes \psi \rangle + R_{jk} \langle V_{ik}, \varphi \otimes \psi \rangle \big) &\\ + \sum_k \sigma_{ik}\sigma_{jk} \langle q V_{kk}, \varphi \otimes \psi \rangle \Big\}\,ds.&\end{aligned}$$

Here each integrand is evaluated at s. Finite sums of product test functions $\varphi \otimes \psi$ are dense in $L^2(\mathcal{D}^2)$ and form a core for the product Laplacian with the stated boundary conditions. Hence this identity is the weak form of (20), with the separate spatial operators $D_i\Delta_x + D_j\Delta_y$.

Step 4: Closure and uniqueness. The right-hand side of (20) uses only the collection $\{V_{k\ell}\}_{k,\ell=1}^{N}$: the noise term involves V_{kk} rather than a third- or fourth-order moment. Thus the two-point system closes. To see uniqueness, let Z be the difference of two mild solutions with identical initial data. The diffusion operator $BZ = (D_i\Delta_x Z_{ij} + D_j\Delta_y Z_{ij})_{ij}$ generates a contraction semigroup $S(t)$ on $\mathcal{X} = L^2(\mathcal{D}^2)^{N\times N}$. The remaining reaction-and-noise operator L is bounded on $\mathcal{X}$, since R, σ are finite matrices and $\|qZ_{kk}\|_{L^2} \le \bar{q}\|Z_{kk}\|_{L^2}$. The mild integral equation therefore implies

$$\|Z(t)\|_{\mathcal{X}} \le \|L\| \int_0^t \|Z(s)\|_{\mathcal{X}}\, ds.$$

Gronwall's inequality gives $Z = 0$. Existence of the mild solution follows from the bounded-perturbation theorem for this same contraction semigroup, and the weak identity above identifies the moment field with that unique solution. The identical deterministic argument applies to the mean system (19). This proves (a) and (b). □

Proposition **??** makes §1 apply verbatim to every moment equation. This formulation buys us a couple nice formulations:

1. **One step for the whole hierarchy** (Cor. **??**): $\Delta t_{\text{opt}} = \Delta x^2/6D$ governs every M_k simultaneously; σ and k appear only in the correction coefficients (5) — e.g. the mean ($\alpha = \frac{1}{2}\sigma^2$) and second moment ($\alpha = \sigma^2$) are advanced by the same step, different corrections. Since Itô drift is necessarily a defect and not a structural change (the noise-induced reaction term is explicitly computable) it slots into $C^{(1)}$ exactly as any reaction term would.

2. **Weak, not strong, by construction:** The method never samples $\dot{W}$, so the Davie–Gaines strong-order-1/2 ceiling (Davie and Gaines, 2001) governs a problem we do not solve. The delivered object is fourth-order convergence of the computed moment fields to the true moments of the closed system [V](scalar, Table 1); conversion to statements about the SPDE law rides entirely on Proposition **??**'s exact closure.

3. **Stability transfers:** the growing-moment case is exactly the $\eta \ge 0$ regime of Prop. 6 — the correction marginally *shrinks* the CFL window and is comfortably stable at Δt_{opt} [V].

unsure of this

The competitor, stated fairly

Euler–Maruyama on (**??**) estimates M_k from M paths with error $O(\Delta t) + O(M^{-1/2})$ per point; our implementation (`baselines.py`, `euler_maruyama_moment`) exists for exactly this comparison [S][1]. The structural comparison, however, is already decisive and does not need the figure: for the *moments alone*, the pathwise route pays M path-solves plus Monte Carlo variance to approximate what the moment route computes deterministically — and the moment route's solver is fourth-order where EM is weak-first. The pathwise route is irreplaceable when paths are the product (extremes, hitting, filtering); for moment delivery under exact closure it is dominated.

3.2 Where the noise model ends

Three extensions are visible from inside the current theory, all open:

[1] The EM baseline is implemented and its noise scaling ($\sigma u\sqrt{\Delta t/\Delta x}\, Z$ per cell per step) is the standard space-time white noise discretization; the head-to-head convergence figure against it is not yet in the results ledger — it is scheduled with the benchmarking gate of §**??**, per the project's own decision to hold benchmarking until scoping stabilized.

- **Colored noise** (trace-class Q): the closure proof goes through with $\sigma^2 \to$ the covariance trace structure; the M/V systems acquire nonlocal reaction kernels. The OTS cancellation, being diffusive-local, should survive with kernel-aware corrections [O].
- **Noise discretization interplay:** how Wiener-increment sampling on the grid (piecewise-constant vs. linear-in-time noise, (Lord and Petersson, 2022)) interacts with the NIDC cancellation — for the moment route there is no sampled noise, which is precisely the advantage; the question becomes how to *validate* moment solvers against pathwise references fairly [O].
- **Boundary conditions for V:** Neumann data on u induce $\partial_n V_{ij} = 0$ on $\partial\Omega \times \Omega \cup \Omega \times \partial\Omega$; used but not proven in the coupled harness [O].

References

Kevin T. Chu. Boosting the accuracy of finite difference schemes via optimal time step selection and non-iterative defect correction. *Applied Mathematics and Computation*, 218(7):3596–3614, 2011. doi: 10.1016/j.amc.2011.08.108. arXiv:0811.2847.

A. M. Davie and J. G. Gaines. Convergence of numerical schemes for the solution of parabolic stochastic partial differential equations. *Mathematics of Computation*, 70(233):121–134, 2001. doi: 10.1090/S0025-5718-00-01224-2.

Pak-Wing Fok and Robert R. Rosales. Multirate integration of axisymmetric step-flow equations. *arXiv preprint*, 2010. arXiv:0810.2517.

C. William Gear and David R. Wells. Multirate linear multistep methods. *BIT Numerical Mathematics*, 24:484–502, 1984.

Gabriel J. Lord and Andreas Petersson. Piecewise linear interpolation of noise in finite element approximations of parabolic SPDEs. *arXiv preprint*, 2022. arXiv:2210.11102.