

DEPARTMENT OF MATHEMATICS

Course: NUMBER THEORY, VECTOR CALCULUS AND COMPUTATIONAL METHODS	TEST-I	Maximum marks: 50
Course code: 22MA21C	Second semester 2022-2023 Physics Cycle Branch: AI, BT, CD, CS, CY, IS, SPARK-C	Time: 9:30AM-11:00AM Date: 10-07-2023

Scheme and Solutions

Q.No	PART -B	Marks																																																																																				
1.	Cumulative frequency table	1																																																																																				
	<table><tr><td>Marks less than (x)</td><td>30</td><td>40</td><td>50</td><td>60</td><td>70</td><td>80</td></tr><tr><td>Number of students (y)</td><td>35</td><td>84</td><td>146</td><td>220</td><td>260</td><td>280</td></tr></table>		Marks less than (x)	30	40	50	60	70	80	Number of students (y)	35	84	146	220	260	280																																																																						
	Marks less than (x)	30	40	50	60	70	80																																																																															
	Number of students (y)	35	84	146	220	260	280																																																																															
	Difference table	3																																																																																				
	<table><tr><td>x</td><td>y</td><td>Δy</td><td>Δ²y</td><td>Δ³y</td><td>Δ⁴y</td><td>Δ⁵y</td></tr><tr><td>30</td><td>35</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td></td><td></td><td>49</td><td></td><td></td><td></td><td></td></tr><tr><td>40</td><td>84</td><td></td><td>13</td><td></td><td></td><td></td></tr><tr><td></td><td></td><td>62</td><td></td><td>-1</td><td></td><td></td></tr><tr><td>50</td><td>146</td><td></td><td>12</td><td></td><td>-45</td><td></td></tr><tr><td></td><td></td><td>74</td><td></td><td>-46</td><td></td><td>105</td></tr><tr><td>60</td><td>220</td><td></td><td>-34</td><td></td><td>60</td><td></td></tr><tr><td></td><td></td><td>40</td><td></td><td>14</td><td></td><td></td></tr><tr><td>70</td><td>260</td><td></td><td>-20</td><td></td><td></td><td></td></tr><tr><td></td><td></td><td>20</td><td></td><td></td><td></td><td></td></tr><tr><td>80</td><td>280</td><td></td><td></td><td></td><td></td><td></td></tr></table>		x	y	Δy	Δ²y	Δ³y	Δ⁴y	Δ⁵y	30	35								49					40	84		13						62		-1			50	146		12		-45				74		-46		105	60	220		-34		60				40		14			70	260		-20						20					80	280					
	x		y	Δy	Δ²y	Δ³y	Δ⁴y	Δ⁵y																																																																														
	30		35																																																																																			
				49																																																																																		
	40		84		13																																																																																	
			62		-1																																																																																	
50	146			12		-45																																																																																
			74		-46		105																																																																															
60	220			-34		60																																																																																
		40		14																																																																																		
70	260		-20																																																																																			
		20																																																																																				
80	280																																																																																					
$p = \frac{x-x_0}{h} = \frac{45-30}{10} = 1.5$ and $p = \frac{x-x_n}{h} = \frac{65-80}{10} = -1.5$	1																																																																																					
Using Newton’s forward interpolation formula, we get	2																																																																																					
$y = y_0 + p\Delta y_0 + \frac{p(p-1)}{2!}\Delta^2 y_0 + \frac{p(p-1)(p-2)}{3!}\Delta^3 y_0 + \dots$																																																																																						
$y(45) = 111.15 \approx 111$	2																																																																																					
Using Newton’s backward interpolation formula, we get																																																																																						
$y = y_0 + p\nabla y_n + \frac{p(p+1)}{2!}\nabla^2 y_n + \frac{p(p+1)(p+2)}{3!}\nabla^3 y_n + \dots$	2																																																																																					
$y(65) = 246.0117 \approx 246$	1																																																																																					
The number of students with marks between 45 and 65 is $246 - 111 = 135$.																																																																																						
2(a)	$y = \frac{(x-367)(x-378)(x-387)(x-399)}{(361-367)(361-378)(361-387)(361-399)}154.9$ $+ \frac{(x-361)(x-378)(x-387)(x-399)}{(367-361)(367-378)(367-387)(367-399)}167.9 +$ $\frac{(x-361)(x-367)(x-387)(x-399)}{(378-361)(378-367)(378-387)(378-399)}191$ $+ \frac{(x-361)(x-367)(x-378)(x-399)}{(387-361)(387-367)(387-378)(387-399)}212.5$ $+ \frac{(x-361)(x-367)(x-378)(x-387)}{(399-361)(399-367)(399-378)(399-387)}244.2$ $y(x) = -0.00002x^4 + 0.0314x^3 - 17.9744x^2 + 4568.21x - 435168 = 178.1827$	2																																																																																				
		4																																																																																				

2(b)	$x = \frac{(y-24)(y-54)(y-129)}{(10-24)(10-54)(10-129)} + \frac{(y-10)(y-54)(y-129)}{(24-10)(24-54)(24-129)}^3$ $+ \frac{(y-10)(y-24)(y-129)}{(54-10)(54-24)(54-129)}^5 + \frac{(y-10)(y-24)(y-54)}{(129-10)(129-24)(129-54)}^8$ When $y = 100, x = 5.9199$	2 2																																																												
3(a)	Difference table <table><tr><td>v</td><td>p</td><td>Δp</td><td>$\Delta^2 p$</td><td>$\Delta^3 p$</td><td>$\Delta^4 p$</td></tr><tr><td>2</td><td>105</td><td></td><td></td><td></td><td></td></tr><tr><td></td><td></td><td>-62.3</td><td></td><td></td><td></td></tr><tr><td>4</td><td>42.7</td><td></td><td>44.9</td><td></td><td></td></tr><tr><td></td><td></td><td>-17.4</td><td></td><td>-36.1</td><td></td></tr><tr><td>6</td><td>25.3</td><td></td><td>8.8</td><td></td><td>32.2</td></tr><tr><td></td><td></td><td>-8.6</td><td></td><td>-3.9</td><td></td></tr><tr><td>8</td><td>16.7</td><td></td><td>4.9</td><td></td><td></td></tr><tr><td></td><td></td><td>-3.7</td><td></td><td></td><td></td></tr><tr><td>10</td><td>13</td><td></td><td></td><td></td><td></td></tr></table> $\left(\frac{dp}{dv}\right)_{v=4} = \frac{1}{h}\left[\Delta p_0 - \frac{1}{2}\Delta^2 p_0 + \frac{1}{3}\Delta^3 p_0 - \frac{1}{4}\Delta^4 p_0 + \dots\right] = \frac{1}{2}\left[-17.4 - \frac{8.8}{2} - \frac{3.9}{3}\right] = -11.55$ $\left(\frac{dp}{dv}\right)_{v=8} = \frac{1}{h}\left[\nabla p_n + \frac{1}{2}\nabla^2 p_n + \frac{1}{3}\nabla^3 p_n + \frac{1}{4}\nabla^4 p_n + \dots\right] = \frac{1}{2}\left[-8.6 + \frac{8.8}{2} - \frac{36.1}{3}\right] = -8.11$	v	p	Δp	$\Delta^2 p$	$\Delta^3 p$	$\Delta^4 p$	2	105							-62.3				4	42.7		44.9					-17.4		-36.1		6	25.3		8.8		32.2			-8.6		-3.9		8	16.7		4.9					-3.7				10	13					2 2
v	p	Δp	$\Delta^2 p$	$\Delta^3 p$	$\Delta^4 p$																																																									
2	105																																																													
		-62.3																																																												
4	42.7		44.9																																																											
		-17.4		-36.1																																																										
6	25.3		8.8		32.2																																																									
		-8.6		-3.9																																																										
8	16.7		4.9																																																											
		-3.7																																																												
10	13																																																													
3(b)	Auxiliary equations $m^3 - 8m^2 + 5m + 50 = 0$ Roots $m = -2, 5, 5$ $x = c_1 e^{-2t} + (c_2 + c_3 t)e^{5t}$	1 1 2																																																												
4	Auxiliary equations $2m^2 + m - 1 = 0$, Roots $m = \frac{1}{2}, -1$ $C.F = c_1 e^{x/2} + c_2 e^{-x}$ $PI = \frac{x e^{x/2}}{3} - \frac{1}{2} + \frac{33 \cos(4x) - 4 \sin(4x)}{2210} - (x^2 + 2x + 6)$ $y = CF + PI$	1 1 2+3+2 1																																																												
5(a)	Auxiliary equations $m^2 - 4m + 13 = 0$, Roots $m = 2 \pm 3i$ $C.F = e^{2x}(c_1 \cos(3x) + c_2 \sin(3x))$ $PI = \frac{1}{D^2 - 4D + 13} e^{2x} \sin(3x) = e^{2x} \frac{1}{D^2 + 9} \sin(3x) = -\frac{x e^{2x} \cos(3x)}{6}$ $y = CF + PI = e^{2x}(c_1 \cos(3x) + c_2 \sin(3x)) - \frac{x e^{2x} \cos(3x)}{6}$	1 1 2 1																																																												
5(b)	Auxiliary equations $m^2 + \mu = 0$, Roots $m = \pm \sqrt{\mu} i$ $x = c_1 \cos(\sqrt{\mu} t) + c_2 \sin(\sqrt{\mu} t)$ $x' = -c_1 \sqrt{\mu} \sin(\sqrt{\mu} t) + c_2 \sqrt{\mu} \cos(\sqrt{\mu} t)$ $c_1 = -a$ and $c_2 = 0$ $x = -a \cos(\sqrt{\mu} t)$	1 1 2 1																																																												