

Your Thesis Title

a Doctoral Thesis

submitted to earn the academic title

doctor rerum naturalium (Dr. rer. nat.)

submitted to

The Faculty Council,
Faculty of Mathematics and Computer Science,
Friedrich Schiller University Jena

by

Aditty Pal
born on 2000 in India.

Reviewers:

1. _____
2. _____
3. _____

Date of the public thesis defense: _____

CONTENTS

1	Toolbox	1
1.1	Directed Hypergraphs	1
1.2	Integer Hyperflows	4
2	Preliminaries	11
2.1	Generation of Reaction networks	15
2.1.1	Graph transformation rules	16
2.1.2	Rule application to a graph	19
2.1.3	Construction of the reaction network	21
2.2	Hyperflows and Fluxes	24
3	A Constraint-based Model	29
3.1	Mathematical Formulation	31
3.1.1	Constraint-Based Model	32
3.1.2	Objective Function	39
3.2	Implementation	41
3.2.1	Implementing the Objective function	42
3.2.2	Linearizing the Constraints	42
3.2.3	Implemented Model	44
3.3	Contextualizing this method	47
4	A Case Study	55
4.1	Network Generation	55
4.2	Thermodynamic Oracle	56
4.3	On assigning the Chemical Potentials	60
4.4	On the range of the concentration variable	63
4.5	Formamide Chemistry	65
5	Refinements to the model	79
5.1	From Rate Constants to Edge Weights	81
5.2	Formulating an Objective Function	84
5.3	ILP Modeling	86
5.3.1	Enumeration of solutions	89
5.3.2	Partial Ordering of Molecules	92
5.4	Comparison with other methods	95
6	Another case study	101
7	Epilogue	113

TOOLBOX

This chapter is identical to Section 2 of the published version of the preprint at [arXiv:2411.15900](#).

In this chapter, the mathematical models for chemical reaction networks and pathways in them are presented, namely directed hypergraphs and integer hyperflows, respectively. This section is based on previous work which have used these concepts [1, 2].

1.1 DIRECTED HYPERGRAPHS

A chemical reaction network consists of set of molecules V and a set of reactions E . Each reaction $e \in E$ is generally a many-to-many mapping of molecules, which can be written [3] as the formal sum

$$e: \sum_{v \in V} s_{ve}^- v \longrightarrow \sum_{v \in V} s_{ve}^+ v$$

where s_{ve}^- and s_{ve}^+ denote the stoichiometric coefficients,

s_{ve}^- is the number of molecules of v that are used

s_{ve}^+ is the number of molecules of v that are produced

by reaction e . The sum is over the entire set of molecules V in the chemical reaction network, and for molecules v that do not take part in a particular reaction e , we have $s_{ve}^- = s_{ve}^+ = 0$

A chemical reaction network is formally well represented [2] as a directed multi-hypergraph (V, E) , where

Note that the use of $^+$ and $^-$ is swapped compared to the notation in the referred previous work using hyperflows [2].




Figure 1: A directed hyperedge, which represents a reaction where one copy of the molecule represented by the vertex v_0 reacts with two copies of v_1 to produce two copies of v_2 . This hyperedge can be considered to represent the reaction with the vertices representing the molecules: $v_0 \equiv \text{O}_2$, $v_1 \equiv \text{H}_2$ and $v_2 \equiv \text{H}_2\text{O}$ and the directed hyperedge representing the chemical reaction $e : \text{O}_2 + 2 \text{H}_2 \longrightarrow 2 \text{H}_2\text{O}$.




Figure 2: A directed hypergraph, where circles are vertices and the squares are hyperedges. Hyperedges with multiple copies of a vertex as source or target are depicted with parallel arrows. See for instance e_4 , which has v_3 twice as a target and thus represents the reaction $e_4 : v_0 \rightarrow 2v_3$. Hyperedges allow the many-to-many relations in chemical reaction to be captured in themodelling unlike edges in simple directed graphs.

the set V of vertices models molecules and
the set E of directed multi-hyperedges models reactions.

A directed multi-hyperedge $e \in E$ is an ordered pair (e^-, e^+) of multisets of vertices, which represents a reaction in the chemical reaction network. The elements of e^- model the reactants of the reaction, with each reactant v appearing s_{ve}^- times, and correspondingly e^+ and s_{ve}^+ model the products of the reaction. An example hyperedge representing a chemical reaction is shown in Figure 1.

Figure 2 depicts a hypergraph with eight vertices and eleven hyperedges. As examples,

- hyperedge $e_0 \equiv (\{v_0\}, \{v_1\})$
represents the one-to-one reaction $v_0 \rightarrow v_1$,
- hyperedge $e_3 \equiv (\{v_0, v_1\}, \{v_3\})$
represents the many-to-one reaction
 $v_0 + v_1 \rightarrow v_3$,
- hyperedge $e_2 \equiv (\{v_2, v_2\}, \{v_1\})$
represents the many-to-one reaction $2v_2 \rightarrow v_1$,
- hyperedge $e_4 \equiv (\{v_3\}, \{v_0, v_1\})$
represent the one-to-many reaction
 $v_3 \rightarrow v_0 + v_1$,
- hyperedge $e_7 \equiv (\{v_3\}, \{v_5, v_5\})$
represent the one-to-many reaction
 $v_3 \rightarrow 2v_5$, and
- hyperedge $e_9 \equiv (\{v_3, v_4\}, \{v_7, v_8\})$
represents the many-to-many reaction
 $v_3 + v_4 \rightarrow v_7 + v_8$.

On a related note, standard usage of the term ‘reaction’ implies the breaking and forming of chemical bonds. However, physical processes that do not alter the set of bonds in a molecule yet involve energy transformations, thereby affecting the molecule’s ability to participate in a reaction. For instance, some reactions occur only when a molecule is excited into a higher energy state, or certain molecules (such as phosphorus or sulfur) are in specific phases. Although such transitions are not conventionally termed ‘reactions,’ they are elementary processes accompanied by a change in free energy. Consequently, in the rest of this paper we

will use the term ‘reaction’ to denote this broader set of relations between classes of molecules.

1.2 INTEGER HYPERFLOWS

A pathway is a set of reactions that transforms some available source molecules into desired target molecules. To model pathways formally, the concept of an *integer hyperflow* f in a hypergraph $\mathcal{H} = (V, E)$ was introduced by [2].

A pathway is viewed as a vector $f_E \in \mathbb{N}_0^{|E|}$, i.e., an assignment of a non-negative integer to each edge in E , denoting how many times the corresponding reaction appears in the pathway. Additionally, two vectors $f_{\text{in}} \in \mathbb{N}_0^{|V|}$ and $f_{\text{out}} \in \mathbb{N}_0^{|V|}$ denote for each molecule in V the amount of it consumed as source molecule, respectively produced as target molecule, by the pathway.

The pathway’s net effect on the number of molecules in the system is specified entirely by f_{in} and f_{out} .

$$\mathbf{S}^+ \cdot f_E - \mathbf{S}^- \cdot f_E = f_{\text{out}} - f_{\text{in}} \quad (1)$$

where

$\mathbf{S}^-$ is a $|V| \times |E|$ matrix with entries s_{ve}^-
 $\mathbf{S}^+$ is a $|V| \times |E|$ matrix with entries s_{ve}^+

(recall from definition of hypergraphs that s_{ve}^- and s_{ve}^+ are the numbers of molecule v consumed and produced by reaction e). In hypergraph terminology, $\mathbf{S}^-$ and $\mathbf{S}^+$ are the out-incidence matrix and the in-incidence matrix, respectively.¹ The matrices $\mathbf{S}^-$ and $\mathbf{S}^+$ for the hypergraph in Figure 2 are shown in Figure 4.

¹ In other settings, $\mathbf{S}^-$ and $\mathbf{S}^+$ are called the reactant stoichiometric matrix and the product stoichiometric matrix. One can also meet a single stoichiometric matrix, defined as $\mathbf{S} = \mathbf{S}^+ - \mathbf{S}^-$. This latter version is less expressive for chemical reaction network modeling, as the role of catalysts and autocatalytic molecules cannot be captured. In some contexts, f_{in} and f_{out} are called the external compounds.




Figure 3: An integer hyperflow query in the directed hypergraph from Figure ref: hypergraph, where v_0 is specified as the source and v_7 is specified as the target. The exact inflow and outflow on these vertices are not specified, and the hyperpaths in Figure 5a, 5b, 5c, 5d and 5e would be valid solutions of this flow query if the required additional sets of vertices is allowed as sources (which are $\emptyset$, $\{v_2\}$, $\{v_1\}$, $\{v_2\}$, $\{v_6\}$ respectively). If additional vertices are not allowed as sources, only the hyperpath in Figure 5a is a valid solution.

To unify the three vectors f_E , f_{in} , and f_{out} into a single flow vector on hyperedges, [2] introduced for each $v \in V$ two so-called half-edges

$$e_v^+ = (\emptyset, v) \text{ and}$$

$$e_v^- = (v, \emptyset)$$

$$\begin{array}{c}
 v_0 \\
 v_1 \\
 v_2 \\
 v_3 \\
 v_4 \\
 v_5 \\
 v_6 \\
 v_7 \\
 v_8
 \end{array}
 \begin{array}{c}
 e_0 \quad e_1 \quad e_2 \quad e_3 \quad e_4 \quad e_5 \quad e_6 \quad e_7 \quad e_8 \quad e_9 \quad e_{10} \\
 \left[\begin{array}{cccccccccccc}
 1 & & & 1 & & & & & & & \\
 & 1 & & & & 1 & & & & & \\
 & & 2 & & & 1 & & & & & 1 \\
 & & & & 1 & & & 1 & & & \\
 & & & & & & 1 & & & 1 & 1 \\
 & & & & & & & & & & \\
 & & & & & & & & & & \\
 & & & & & & & & & & \\
 & & & & & & & & & &
 \end{array} \right]
 \end{array}$$

(a) in-incidence matrix

$$\begin{array}{c}
 v_0 \\
 v_1 \\
 v_2 \\
 v_3 \\
 v_4 \\
 v_5 \\
 v_6 \\
 v_7 \\
 v_8
 \end{array}
 \begin{array}{c}
 e_0 \quad e_1 \quad e_2 \quad e_3 \quad e_4 \quad e_5 \quad e_6 \quad e_7 \quad e_8 \quad e_9 \quad e_{10} \\
 \left[\begin{array}{cccccccccccc}
 & & & & & & & & & & \\
 & 1 & & & 1 & & & & & & \\
 1 & & 1 & & & & 1 & & & & \\
 & & & & & & 1 & & & & \\
 & & & 1 & & & & & & & \\
 & & & & 1 & & & & & & \\
 & & & & & & 2 & & & & \\
 & & & & & & & 2 & & & \\
 & & & & & & & & 1 & & 1 \\
 & & & & & & & & & 1 & \\
 & & & & & & & & & 1 &
 \end{array} \right]
 \end{array}$$

(b) in
out-incidence matrix

Figure 4: A representation of the hypergraph from Figure 2 as 4b an out-incidence matrix $\mathbf{S}^-$ and 4a an in-incidence matrix $\mathbf{S}^+$. Only non-zero entries are shown. Since the hypergraph has 9 vertices and 11 hyperedges, the matrices have dimensions 9×11 .

These represent the production of v as source molecule addition to the system and the consumption of v as target molecule removal from the system, respectively. Setting

$$\begin{aligned}
 E^+ &= \{e_v^+ \mid v \in V\} \\
 E^- &= \{e_v^- \mid v \in V\} \\
 \bar{E} &= E \cup E^+ \cup E^-,
 \end{aligned}$$

[2] introduced the extended hypergraph $\overline{\mathcal{H}} = (V, \overline{E})$ and defined an *integer hyperflow* to be a vector $f \in \mathbb{N}_0^{|\overline{E}|}$ satisfying the flow conservation constraint

$$\forall v \in V: \sum_{e \in \overline{E}} s_{ve}^+ f_e - \sum_{e \in \overline{E}} s_{ve}^- f_e = 0 \quad (2)$$

where,

s_{ve}^- an entry in the $|V| \times |\overline{E}|$ out-incidence matrix of $\overline{\mathcal{H}}$
 s_{ve}^+ an entry in the $|V| \times |\overline{E}|$ in-incidence matrix of $\overline{\mathcal{H}}$
 f_e an entry in f

The constraint in Equation (2) is easily seen to be equivalent to the constraint in Equation (1).

In Figure 5, three possible hyperflows in a hypergraph are depicted. Unboxed integers are flow values on half-edges, while boxed integers are the flow values for hyperedges in E . The inflow and the outflow are depicted as half-edges in and out of the source and target vertices of the pathway. For clarity, only half-edges with non-zero flow values are shown.

One key feature of integer hyperflows is that they give a flexible and generic way to *specify* pathways via constraints on the flow on half-edges. This may be as straight-forward as specifying an overall reaction for the pathway by fixing the flow on the half-edges for the molecules in the source and target sets and setting the flow to zero for the rest of the half-edges. Or one may specify upper and lower limits for some half-edges and free values for others, for instance to ask if there is any way to produce at least some target molecules using any amount of source molecules, while at the same time allowing waste products.


Then a *search* for pathways fulfilling the specification amounts to finding flow vectors f obeying those constraints, in addition to the constraint in Equation (2). Since integer hyperflows are integer valued, it is natural to model the situation via the ILP formalism and search for f using an ILP solver. This idea has been implemented as part of the software system MØD [4]. The result is a powerful and flexible tool for specifying and searching for pathways (integer hyper-

A detailed discussion of how integer hyperflows relate to, and differ from, other similar concepts such as elementary modes, extreme pathways, and flux balance analysis can be found in Section 2.6 of [2].


flows) in a given chemical reaction network (hypergraph). The flexibility is further increased by the possibility of having flow values in the ILP objective function, such that e.g. flow solutions allowing but minimizing waste can be found. Also the flow values on non-half-edges can enter the specification and/or optimization, allowing e.g. the search for solutions minimizing the use of a given reaction deemed particularly expensive to include in the pathway.

REFERENCES


- [1] Jakob L. Andersen et al. “An intermediate level of abstraction for computational systems chemistry”. In: *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 375.2109 (2017), p. 20160354. DOI: [10.1098/rsta.2016.0354](https://doi.org/10.1098/rsta.2016.0354). URL: <https://royalsocietypublishing.org/doi/abs/10.1098/rsta.2016.0354>.
- [2] Jakob L. Andersen et al. “Chemical Transformation Motifs—Modelling Pathways as Integer Hyperflows”. In: *IEEE/ACM Trans. Comput. Biol. Bioinformatics* 16.2 (Apr. 2019), pp. 510–523. ISSN: 1545-5963. DOI: [10.1109/TCBB.2017.2781724](https://doi.org/10.1109/TCBB.2017.2781724). URL: <https://doi.org/10.1109/TCBB.2017.2781724>.
- [3] Stefan Müller, Christoph Flamm, and Peter F. Stadler. “What makes a reaction network “chemical”?” In: *J Cheminform* 14.63 (Sept. 2022). DOI: [10.1186/s13321-022-00621-8](https://doi.org/10.1186/s13321-022-00621-8). URL: <https://doi.org/10.1186/s13321-022-00621-8>.
- [4] Jakob L. Andersen. *MØD*. <http://mod.imada.sdu.dk>. 2024.




(a) hyperpath 1




(b) hyperpath 2




(c) hyperpath 3



(d) hyperpath 4



(e) hyperpath 5




(f) hyperpath 6

Figure 5: Six possible integer hyperflows [5a](#), [5b](#), [5c](#), [5d](#), [5e](#), and [5f](#) in the hypergraph from Figure 2, each representing a pathway, where hyperedges with non-zero flow are drawn in bold. The integers in the boxes represent the hyperflow assigned to the respective hyperedge in the pathway.


PRELIMINARIES

This chapter is identical to Appendix A and B of the published version of the preprint at [arXiv:2504.10609](#).


While one might choose to represent a reaction formally by (1.1), for more practical scenarios, it is just a collection of reactant molecules (with their multiplicities) that gets converted into a collection of product molecules (with some different multiplicities). For example,



This reaction is a two-to-one map which converts molecules $\{v_0, v_1\}$ to $\{v_3\}$. In a hypergraph, we choose to represent this map graphically as



This object is called an hyperedge and the individual molecules v_0 , v_2 and v_3 form the vertices. However, in a practical scenario, when the molecules v_0 and v_2 are allowed to react, reaction e_1 is not the only reaction that would occur and the system would not stop reacting after producing v_3 . Successive reactions might also be possible, some of which might be:



Some of the reactions that might occur when molecules of v_0 and v_2 are allowed to react is shown. To a chemist, not all the reactions would look feasible due to reactivity constraints on the molecules, which would disallow some of the reactions. However, all these reactions might possibly occur in the system in parallel, making it an example of a concurrent system.






Figure 6: A hypergraph representation of the reactions in the concurrent system.



(a) sequence of reactions forming glyoxal from glycolonitrile



(b) A pathway in a hypergraph with the queried molecules v_9 . The number in the squares denote the number of times the reaction is used to carry out the transformation or the flow assigned to the corresponding hyperedge.

Figure 7: The hyperflow denoting the pathway from Figure 7a is shown in the hypergraph from Figure 6.

We would represent each reaction by a separate hyperedge and each class of molecules by a distinct vertex. This results in the structure shown in Figure 6. The reverse reactions are omitted from the hypergraph in Figure 6 to reduce clutter. However, in our actual modeling, we do add separate hyperedges to represent the forward and backward directions of reversible reactions.

This data structure representing the concurrent set of reactions in the system is called a multi-directed hypergraph or for simplicity, just a hypergraph. An advantage of using this data structure is that it faithfully represents the many-to-many mappings which are present in reactions which are not unimolecular. Contrast this with a simple directed graph containing edges which map one vertex to another vertex, which are often used to represent and study diverse networks in many situations. The hypergraph in Figure 2 is an abstraction of another system of concurrent reactions.

One observes that glyoxal is represented as vertex v_9 in the hypergraph. One might query, how one inside this hypergraph can find a pathway producing glyoxal while using the input molecules v_0 and v_2 . One possible transformation pathway is shown in Scheme 7a.

Highlighting the hyperedges used in this transformation pathway in the hypergraph results in Figure 7b. This sequence of reactions in a pathway can be found by formulating it as a flow query with molecules v_0 and v_2 as the source molecules and v_9 as the target molecule. The ILP assigns integers to the hyperedges which denote the number of times the hyperedge is used in the pathway. Since the flows assigned to the hyperedges are constrained to be integers, this is called an integer hyperflow. The motivation behind using allowing only non-negative integers as permissible hyperflows on a hyperedge is discussed in Section ??.

Often, there might exist alternative solutions to the flow query. In this case, there exists another valid integer hyperflow to the same flow query in the network with molecules




Figure 8: An alternative pathway to the same flow query as in Figure 7b.

v_0 and v_2 as the source molecules and v_9 as the target molecule which is shown in Figure 8

It might so happen that the reaction $v_3 \rightarrow v_7$ is slow (because it has a high energy barrier) and the two step process $v_3 \rightarrow v_5 \rightarrow v_7$ is faster. To judge which of the two pathways is better, a score is assigned to them by evaluating an objective function (expression (24) is used in this work) on the pathway, and the one with a better score is ranked higher. This automated search for pathways in a reaction system represented as a hypergraph is the motivation behind introducing the formalism of integer hyperflows in Chapter 1.

2.1 GENERATION OF REACTION NETWORKS

There exists a diverse range of reactions. However, they are often grouped together on the basis of their mechanisms to create classes of reactions. Reactions in a class share some properties: the reactive core (type of atoms taking part in the reaction), the bonds that are broken, the movement of electrons, and the new bonds that are formed. We choose to represent these classes of reactions using reaction templates. Each template defines a change in the bondset of the molecules that take part in the reaction.

For example, in the list of reaction depicted in the hypergraph 6:

- reactions e_3 , e_5 , e_6 , e_7 and e_8 have a 1,3-shift of a hydrogen atom (often called tautomerism).
- reactions e_2 , e_4 and e_{11} are additions to an trigonal planar carbon to produce a tetrahedral carbon.
- reactions e_9 and e_{10} eliminate a small molecule from a carbon with two or more adjacent electronegative atoms to form a trigonal planar carbon.
- reaction e_1 is an addition to a linear carbon to produce a trigonal carbon.

Some might prefer more fine-grained reaction templates. For instance, one might define separate templates for keto-enol tautomerism, amide-iminol tautomerism or imine-enamine tautomerism, and similarly for addition to a trigonal planar carbonyl carbon atom and addition to a trigonal planar imine carbon atom. However, that is simply a question of the level of abstraction that the user chooses to work with.

2.1.1 *Graph transformation rules*


We choose to represent molecules using undirected labeled graphs (hereafter called molecular graphs). The nodes in the graph represent atoms labeled by the symbol of the element, and can carry attributes such as the charge on the atom. The edges in the graph represent bonds labeled by the bond type (single, double, triple or aromatic). with this representation of molecules, the reaction templates used to abstract reactions become graph transformation rules.

A graph transformation rule consists of a subgraph on the left side which can be replaced by the subgraph on the right side when a match of the left-hand side is found in an already existing molecular graph in the network. In other words, the rule converts a part of the graph (representing a fragment of a molecule) and performs a chemical reaction.

Let's consider the exchange of hydrogen atoms between oxygen atoms in a carboxylic acid.




The graph transformation rule for this class of reactions would be:



The amide-iminol conversion is a more substantial example.




The graph transformation rule for this class of reactions would be:




This rule rewrites an amide fragment when it is found in a molecule with an iminol fragment, thus effecting the required reaction.

For a keto-enol tautomerism like



the graph transformation rule for this class of reactions would be:




Note that the graph transformation rule indicates the bare minimum that is required to be present as a subgraph in


the molecule for the reaction to occur. For example, the carbon at the 2-position, might have two adjacent hydrogen atoms, but the rule specifies that at least one hydrogen atom is required for the keto isomer to convert into the enol isomer. Without any hydrogen at the 2-position, the fragment on the left is not a subgraph of the molecule and this transformation rule cannot be applied to the graph.

Also, note that the objective here is not to capture the exact mechanism of the reaction, but rather specify what the molecular fragment is before and after the reaction. In all the three reactions mentioned above, the 1,3-hydrogen transfer involves a solvent molecule. In most cases, it is not the exact same hydrogen atom that would be in the molecule before and after the reaction.

The similarity in the graph transformation rule for the three reactions mentioned above look similar. For a more elegant formulation, we define the following graph transformation rule to account for all the three cases above:




This graph transformation rule also encodes an allyl hydrogen shift, an imine-enamine tautomerism, or the hydrogen exchange in amidine.




However, all these six different reactions are expected to have differing mechanisms. This graph transformation rule introduces two variables $\mathcal{X}, \mathcal{Y}$ which can be matched with

the label C, N or O, or in other words the atoms in the place of $\mathcal{X}$, $\mathcal{Y}$ in the molecular fragment can be a carbon, nitrogen or oxygen atom.

Similarly the graph transformation rule



encodes the addition of a water or ammonia molecule to a carbonyl or imine group.



We admit that these reactions are not a simple one step process as the transformation rule make them appear to be and not all of these possible reactions is always feasible for all reactant molecules.

In all the graph transformation rules in this section, the context graph K , which comes between the left and right-hand sides, is omitted. The context graph must be added to the graph transformation rules for mathematical correctness. It ensures that logically the bonds are first removed, and then added, making the graph rewriting a two-step process.

2.1.2 Rule application to a graph

When a graph transformation rule is applied to a set of reactant molecules, a reaction might occur. First, it is checked if the left-hand side of the rule, L , is present as a subgraph




Figure 9: Two examples of the application of a graph transformation rule to a collection of molecules.

in the set of reactant molecules G . If so, then that subgraph is rewritten according to the rule to R , to generate a new set of molecules H , the product molecules. This process is illustrated in Figure 9.

If the left-hand side of the rule, L , is not present in the set of molecules G , then that reaction template cannot be applied to the given set of molecules.




Figure 10: A case where the graph transformation rule cannot be applied to a molecule

Therefore, the application of a rule to an existing set of molecules might produce new molecules, which in turn expands the molecular space. Repeated application of these rules constructs a network, which is detailed in the next section.

2.1.3 Construction of the reaction network

The construction of the reaction network requires a set of input molecules and some graph transformation rules. A match between a collection of molecules and the subgraph on the left-hand side of one of the transformation rules means that this reaction template can be applied to that collection of molecules created from the input set of molecules. Thus, when a rule is applied to a collection of molecules from the current set of molecules, new molecules are created. In a generative process, these new molecules are then added to the set of available molecules (i.e., this set is updated). Graph transformation rules can be recursively applied to this new set of molecules, effecting more reactions to be made and thus the molecular space expands. The hypergraph representation of the network keeps track of all the reactions performed and of the molecules current present in the network.




Figure 11: (top) The reaction network (with 4 molecules and 2 reactions) after one recursive application of the graph transformation rules, and (bottom) the reaction network (with 9 molecules and 10 reactions) after two recursive applications of the graph transformation rules. The part of the network already generated previously has been shaded.




Figure 12: The reaction network (with 9 molecules and 14 reactions) after three recursive applications of the graph transformation rules with the 3-cycles filtered out. The part of the network already generated previously has been shaded.

To illustrate the construction of a reaction network by the recursive application of graph transformation rules to an input set of molecules, the graph transformation rules from Figure 40 in Chapter 6 is applied to the input molecules water and 2-hydroxyethanenitrile is shown in Figures 11, 12 and 13.

Notice that some molecules in Figure 11, such as the 3-cycles, are not realistic. One can disallow their addition to the generated reaction network by using so-called right predicates. Right predicates filter out certain reactions from the network based on the properties of the molecular graphs that appear on the right-hand side of a derivation. This, it prunes the generated network and makes its growth more manageable and potentially more realistic. The effect of using the right predicate can be seen in the hypergraph in Figure 12, which has 9 molecules after three recursive expansion steps, while the hypergraph in Figure 11 already had 9 molecules after two recursive steps when the right predicate was not used.

Often, one direction of a reaction is added to the hypergraph (adding new molecules to the existing set of molecules) and the reverse direction is added in the next recursion. This adds a new reaction between reactants and products both of which are already present in the set of existing molecules. The graph transformation rules can be applied for a specified number of iterations, or until a specified limit on the size of the different molecules generated is reached which terminated the network generation process. This makes the derived hypergraph finite.

2.2 HYPERFLOWS AND FLUXES

In this section, we give a comparison of our workflow with existing methods. Constraint-based models such as flux balance analysis also use constraint programming with an objective function that is optimized. For instance, open-source tools such as COBRApy [1] perform flux analysis on a reaction network supplied to them. However, such




Figure 13: The reaction network (with 17 molecules and 26 reactions) after four recursive applications of the graph transformation rules with the 3-cycles filtered out. The part of the network that was already generated after three recursive steps has been shaded.

tools do not construct the network on the fly in a generative way— rather the reaction network has to be constructed beforehand using some other tool. Flux balance analysis generates a flux vector for the reactions in the network which is allowed to contain floating-point values, while our method accepts only integers as a valid hyperflow for a reaction, which is more aligned with the standard notion of a pathway. Formulating a flux balance analysis on the network would be equivalent to an LP relaxation of the pathway query problem using integer hyperflows for the same network. A linear program (LP) can be solved in polynomial time while an ILP is NP-hard. However, formulating the search for pathways to a queried molecule in a reaction network offers some advantages.


There exists reaction networks where the optimal solutions from the ILP formulated by our modeling and the relaxed LP from flux balance analysis differ. Consider the reaction network shown in Figure 14a with the following formulation of the problem:

$$\begin{aligned} & \max (\text{outflow}(v_3)) \\ & \text{subject to: } \text{inflow}(v_0) + \text{inflow}(v_1) + \text{inflow}(v_2) \leq 3 \end{aligned}$$


The optimal flux distribution yields an objective value of 1.5 with the assigned fluxes shown in Figure 14b while our method assigns the integer hyperflows shown in Figure 14c in one of the three optimal solutions with objective value 1. The other two solutions assign the hyperflows $(0, 1, 1)$ and $(1, 1, 0)$ to the hyperedge set (e_1, e_2, e_3) respectively.

The ILP solution space is a subset of the LP feasible reaction and therefore, the objective value for the optimal solution for the relaxed LP will always be as at least as large as the objective value of optimal solution for the corresponding ILP. This is illustrated in the example shown in Figure 14 too. As the result, the fluxes and the structure of the solution is often different from those if they were constrained to be integers.


Restricting hyperflows to integers makes physical sense because molecules occur and react in discrete quantities.



(a) An example reaction network with 4 molecules and 3 reactions.



(b) A flux distribution in the network which produces an outflow of 1.5 for v_3 .



(c) An assignment of integer hyperflows in the network which produces an outflow of 1 for v_3 .

Figure 14: Comparison of flux balance analysis and an integer hyperflow in an example reaction network.

Actually, small molecular counts in several biological settings (often involving trace amounts of enzymes and inhibitors affecting multiple iterations of the same pathway) and physical cases (as in chain reactions and cascades) do not allow arbitrary scaling of the number of molecules, as the use of real-valued fluxes by flux balance analysis assume. When a small number of molecules are encountered in a scenario as above (or in scenarios with appearances of a rare but significant molecule in the system), the discrete nature of the molecules becomes more pronounced.

Another feature of our proposed method is the enumeration of alternative pathways with disjoint set of hyperedges ranked on the basis of a chosen objective function. While flux balance analysis offers the possibility to find alternative pathways [2], they have the same objective value but a different flux distribution in the network. This differs from the enumeration of solutions in the current method which offers a list of pathways ordered by the increasing value of the objective function for the user to choose from. Enumerating solutions over variables with a continuous domain is difficult using indicator variables and biimplications as in constraints 12 and 13. The enumeration of solutions offers diverse pathways to the queried molecules and makes the method more robust to uncertainties in the weights assigned to the hyperedges.

REFERENCES


- [1] A. Ebrahim, J.A. Lerman, B.O. Palsson, et al. "COBRApy: CONstraints-Based Reconstruction and Analysis for Python". In: *Syst Biol* 7.74 (Aug. 2013). DOI: [10.1186/1752-0509-7-74](https://doi.org/10.1186/1752-0509-7-74).
- [2] COBRA Toolbox documentation. *Flux Balance Analysis: Alternate optimal solutions*. https://opencobra.github.io/cobratoolbox/stable/tutorials/tutorial_practical_alternateOptimalSolutions.html.

A CONSTRAINT-BASED MODEL


This chapter is identical to Section 3 of the published version of the preprint at [arXiv:2411.15900](https://arxiv.org/abs/2411.15900).

Analogous to typical thermodynamics problems where pathways maximizing the work done by the flow of heat between the system and the reservoir are sought (such as the cycle of a heat engine as shown in Figure 15), we seek pathways in our system to maximize the chemical work done (interpreted as maximizing the production of the target molecules) by directing the flow of molecules along specific reactions on the free energy landscape.

Classical thermodynamics—contrary to the ‘dynamics’ in the name—is used to study systems at equilibrium, or near equilibrium, when the thermodynamic driving forces are in exact balance. In a closed system, the concentrations of the molecules in the system would be such that the chemical potentials of all the molecules in the reaction network is the same. To achieve the required flow along the pathway returned by our method, one has to perturb the system from the equilibrium by tweaking the molecular concentrations in a direction returned by the solution. In practical scenarios, this is done supplying a steady inflow of the source molecules and removing the target molecules from the system to maintain a non-zero thermodynamic driving force along the pathway. For perturbations that are small enough, the equations from classical thermodynamics describing the equilibrium situation can still be used as a first-order approximation. The gradients of the chemical



(a) A heat engine attempts to maximize the work done $W = Q_H - Q_C$, while transferring energy in the form of heat between two reservoirs at temperatures T_0 and T_1 , respectively.



(b) The pathway followed by the heat engine while performing work W is illustrated in the physical $P - V$ space of the system.

Figure 15: Thermodynamics studied the pathways traversed by heat engines in the $P - V$ space which maximized the work done by it. The efficiency of such heat engines was calculated as $\eta = \frac{W}{Q_H} = \frac{Q_H - Q_C}{Q_H} = 1 - \frac{Q_C}{Q_H}$ which can be shown to be equivalent to $\eta = 1 - \frac{T_1}{T_0}$. therefore, increasing the temperature gradient in the system increased the work done by the heat engine under ideal conditions.

potentials in general, or in our case, the differences in the chemical potentials of the target molecules and the source molecules in a elementary reaction serve as the thermodynamic force driving a reaction. In the linear regime, the flow for each reaction would be directly proportional to the thermodynamic forces driving the flow. The forces driving the flows in a reaction network might not remain linear functions of the difference in the chemical potentials farther from the equilibrium, resulting in non-linear dynamics. Therefore, in this work we would restrict ourselves to analyzing the system perturbed, within acceptable limits, from the equilibrium, so that using equations from equilibrium thermodynamics does not introduce large errors into the calculations.

3.1 MATHEMATICAL FORMULATION

The first step to make the search more realistic is to assign the chemical potentials to the vertices and add constraints which allow only thermodynamically favorable reactions in the returned pathway. Chemical potentials are real-valued numbers. Hence, the inclusion of customized linear constraints and objective functions involving floating-point variables for the chemical potentials and concentrations of molecules requires us to extend the existing ILP model in MØD to one using MILP incorporating these variables. This is the key focus of the current work.

MILP attempts to model the system with linear inequalities involving some variables (both integer-values and floats, hence mixed) describing the system. An infeasible system of inequalities indicates that there is no solution to the problem under those specific constraints. When a feasible solution space exists, one often attempts to optimize the solution, by maximizing (or minimizing) the value of some objective function. In this section, we develop the structure of the MILP model, first with using logical constraints using implications in the next section on the constraint-based model, and later, in the discussion on implementing

them, we linearize it into an MILP that can be solved using a solver.

3.1.1 Constraint-Based Model

Since we are extending the existing integer hyperflow formulation defined earlier, the base ILP formulation for modeling flow conservation in pathways in reaction networks is inherited from that work.


$$\forall v \in V: \sum_{e \in \bar{E}} s_{ve}^+ f_e - \sum_{e \in \bar{E}} s_{ve}^- f_e = 0 \quad (2)$$

The working of Constraint 2 is illustrated in the four example cases in Figure 17.


Any set of integer flow values on E induces some overall reaction, which corresponds to flow values on $E^+ \cup E^-$ that make Equation (2) hold. However, the set of available source molecules is often constrained by practical considerations such as cost, ease of availability and purity. Similarly, for the flow to be of practical utility, we could be interested in flows which maximize the outflow of a particular molecule, produce minimum side-products, and perhaps limit the use of particularly expensive reactions. As explained while defining hyperflows, such conditions are modeled as constraints on the flow values f_e , in particular for e in E^+ and E^- , but potentially also in E . These constraints constitute a structural specification of the pathway searched for:

$$\text{Structural specification constraints on } f_e \text{ for } e \in \bar{E} \quad (3)$$

To also incorporate thermodynamics into the modeling, we propose assigning a chemical potentials for each vertex v , $x_v^{G^0}$, under standard physical conditions, which are calculated using a thermodynamic oracle described later in Chapter 4. The differences in the chemical potentials of the target set and the source set in a reaction e , $x_e^{\Delta G}$, serve as the thermodynamic force driving the reaction, dictating the direction of flow of matter along that hyperedge. In



(a) A pathway in a chemical reaction network is a sequence of reactions that convert the source molecules (with supplied inflow) to the target molecules (with required inflow). In a thermodynamically consistent modeling, the pathway would aim to maximize the thermodynamic driving force along the pathway producing the target molecules.



(b) The addition of constraints describing the thermodynamics of the system prunes the solution space of the system, leading to hyperflow solutions that were valid earlier being thermodynamically infeasible.

Figure 16: A thermodynamically informed modeling of pathways in reaction networks requires the addition of constraints based on thermodynamics to capture the energetics in the system. This leads to the modification of the solution space.




Figure 17: The flow conservation constraint 2 ensures steady-state mass conservation in the pathway. The induced hyperflow enforces that the number of copies of each vertex produced by the hyperedges equals the number of copies of the vertices used by the hyperedges. Case 17a has no inflows or outflows (v is neither a source vertex nor a target vertex), case 17b satisfies Constraint 2 but does not add anything physical to the pathway, case 17c involves an outflow (which balances the number of copies of target vertex v produced by the incident hyperedges), while case 17d involves an inflow (where the number of copies of source vertex v supplied is used by the out-going hyperedges).

the linear regime, not far away from the equilibrium, the thermodynamic flows on a particular hyperedge is directly proportional to the thermodynamic force driving it. The difference in chemical potential for a reaction can be manipulated by altering the concentrations of the molecules involved. We choose to represent the log concentration of a molecule v by the variable x_v^K . The nudged difference in chemical potentials for a reaction e is represented as the variable, $x_e^{\Delta G}$ after taking into account the effect of the log concentration, x_v^K , of the molecules involved in the reaction e . We would like to have an assignment of variables x_v^K (within practical limits) such that the pathway consists of reactions with only negative differences in chemical potentials, of in other words, thermodynamically favorable

MILP variable	datatype	description
f_e	integer	flow variable for a hyperedge e
$x_e^{\Delta G}$	float	non-negative differences in chemical potential of in-vertices and out-vertices of a hyperedge e after accounting for their concentrations
x_v^K	float	log concentration of a vertex v
t_v	integer	topological ordering number for vertex v , non-negative
$x_v^{G^0}$	float	chemical potential of a vertex v under standard physical conditions as assigned by the thermodynamic oracle
$x_e^{\Delta G^0}$	float	differences in chemical potential of in-vertices and out-vertices of a hyperedge e under standard physical conditions

Table 1: Variables in the proposed MILP formulation. The variables M and $x_v^{G^0}$ are assigned as input parameters, while the constants $x_e^{\Delta G^0}$ are calculated based on the given $x_v^{G^0}$.

reactions. The variables in the MILP formulation are listed in Table 1.

The flow f_e through a hyperedge e must be positive.

$$\forall e \in \bar{E}: 0 \leq f_e \quad (4)$$

Since the reactions are reversible, separate hyperedges are added for the forward and backward directions. Under specific physical conditions, thermodynamic forces dictates that either the forward or the backward direction of a reversible reaction is more likely. This is illustrated in Figure 18. The direction of the flow is in the direction in which the difference in chemical potential for a hyperedge e , $x_e^{\Delta G}$, is negative. The following implication-based constraint ensures that the pathway returned by the MILP solver consists of thermodynamically favorable reactions only.

$$\forall e \in E: f_e > 0 \implies x_e^{\Delta G} \leq 0 \quad (5)$$

The difference in chemical potential, $x_e^{\Delta G}$ under arbitrary log concentration values, x_v^K of the molecules participating in the reaction e is calculated from the difference in




Figure 18: Separate hyperedges have been added for the reaction $e_{>} : v_0 + v_1 \rightarrow v_2 + v_3$ and its reverse $e_{<} : v_2 + v_3 \rightarrow v_0 + v_1$. The difference in chemical potential for the hyperedge is calculated as $x_{e_{>}}^{\Delta G} = \Delta G(\text{RHS}) - \Delta G(\text{LHS}) = [\Delta G(v_2) + \Delta G(v_3)] - [\Delta G(v_0) + \Delta G(v_1)] = -x_{e_{<}}^{\Delta G}$. As a result of constraints 4 and 5, it is allowed to have a positive hyperflow only on the hyperedge corresponding to $e_{>}$. The hyperedge corresponding to $e_{<}$ might be assigned a hyperflow only when $x_{e_{<}}^{\Delta G} \leq 0$, by some possible assignment of the concentration variables x_v^K of the vertices involved. Therefore, thermodynamic free energy differences dictate the direction of mass flows in the network.

chemical potential under standard conditions, $x_v^{G^0}$, by the following equations.

$$\begin{aligned} \forall e \equiv (e^-, e^+) \in E: \\ x_e^{\Delta G} = & \left(\sum_{v \in e^+} x_v^{G^0} - \sum_{v \in e^-} x_v^{G^0} \right) \\ & + RT \cdot \left(\sum_{v \in e^+} x_v^K - \sum_{v \in e^-} x_v^K \right) \end{aligned} \quad (6)$$

The molecules on the left side and right side of an elementary reaction belong to e^- and e^+ respectively. R is the universal gas constant (in appropriate units) while T is the absolute temperature of the system. In a well-stirred system, RT is a constant for the system. The log concentration of the molecules, x_v^K , is allowed to vary within a range $[x_v^K]_{\min} \leq x_v^K \leq [x_v^K]_{\max}$, set by practical and physical




Figure 19: Often hyperedges with assigned $x_e^{\Delta G^0} > 0$ (which are thermodynamically not feasible under standard physical conditions) can be made thermodynamically feasible by tweaking the concentration variables x_v^K of the vertices involved. The hyperedge from Figure 19a by raising the concentrations of vertices v_0 and v_1 and lowering the concentration of the vertex v_2 as shown in Figure 19b.

constraints detailed in Chapter 4, to mimic the conditions inside a reactor.

$$\left[x_v^K \right]_{\min} \leq x_v^K \leq \left[x_v^K \right]_{\max} \quad (7)$$

The control over the concentration of the molecules in the system is used in several practical scenarios to influence the equilibrium in the system. Often, the concentration of certain molecules in the system are monitored and actively allowed to vary only within certain controlled ranges to facilitate (or inhibit) the formation of certain molecules

in the system. The variation in the concentration of the molecules allows the model to be closer to reality, because stable molecules would have a higher concentration, while reactive intermediate molecules would be expected to have lower concentrations. The variation in the concentration of the molecules can also make a reaction, that was thermodynamically infeasible under standard physical conditions, thermodynamically feasible under certain ranges of the concentration of the molecules involved as shown in Figure 19. This then allows the corresponding hyperedge to have a non-negative flow, and be included in a pathway.

Note that the dependence of difference in the chemical potentials for a reaction e on the concentrations of the molecules involved is not linear.

$$\begin{aligned}
 x_e^{\Delta G} &= x_e^{\Delta G^0} + RT \ln \left(\frac{\prod_{v \in e^+} \text{conc}(v)}{\prod_{v \in e^-} \text{conc}(v)} \right) \\
 &= x_e^{\Delta G^0} + \frac{RT}{\ln 10} \left(\log \left(\prod_{v \in e^+} \text{conc}(v) \right) \right. \\
 &\quad \left. - \log \left(\prod_{v \in e^-} \text{conc}(v) \right) \right) \\
 &= x_e^{\Delta G^0} + \frac{RT}{\ln 10} \left(\sum_{v \in e^+} \log \text{conc}(v) \right. \\
 &\quad \left. - \sum_{v \in e^-} \log \text{conc}(v) \right)
 \end{aligned}$$

where,

$$x_e^{\Delta G^0} = \sum_{v \in e^+} x_v^{G^0} - \sum_{v \in e^-} x_v^{G^0}$$

However, it can be represented as a linear expression, if one chooses to represent the log concentrations of the corresponding molecule v , $\log_{10} \text{conc}(v)$, as a variable x_v^K , in the linear program.

3.1.2 Objective Function

Formulating the search for pathways in reaction networks as an MILP problem, allows us to add an implication constraint to rule out pathways containing one or more individual reactions not favored by thermodynamics. However, it also enables us to exploit another feature of MILPs, namely the objective function to quantify how likely is the entire pathway to occur.

We would like to formulate the search for pathways as an optimization problem aimed at maximizing the outflow of the target molecule(s). To maximize the intended objective, we attempt to maximize the net thermodynamic force driving the flow. Considering the net thermodynamic forces driving the flow to be the sum of the thermodynamic driving force for each individual reaction in the pathway, we choose as objective function the minimization of the cumulative sum of the difference in chemical potentials for each constituent reaction with a non-zero flow.

$$\max \left(- \sum_{e: f_e > 0} x_e^{\Delta G} \right) \quad (8)$$

This choice of the objective function (8) is a heuristic which fits well with the cost of perturbing the system so that queried flow is effected with all reactions being thermodynamically driven in the required direction. Objective function (8) can be related to optimal control principles and considered as the minimum work required to be done to enforce the hyperflow $f : \{e | f_e > 0\}$. Once the system is in equilibrium, it has to be slightly perturbed out of that equilibrium to sustain the hyperflow f . The minimal work required to be performed to achieve the required perturbation should satisfy

$$\langle W_{\min} \rangle \geq \sum_{e: f_e > 0} x_e^{\Delta G}$$

This is similar to the Crooks fluctuation theorem [1, 2] and its implication, the Jarzynski inequality [3–5]. All in all, ranking pathways by the objective function (8) corresponds

Equation (2) in [3] and equations (3.2) and (3.4a) in [4] have a similar form involving the free energy on the right-hand side.

closely to selecting the pathway that requires the least thermodynamic work to sustain the outflow of the queried target molecule. Maximizing objective function (8) becomes equivalent to minimizing the work done to perturb the system from equilibrium to achieve the desired hyperflow.

The motivation behind choosing this particular objective function can also be argued for from the perspective of maximizing the probability of the pathway. We interpret the free energy difference to be the measure of the probability of the reaction as

$$p(e) \sim \exp(-x_e^{\Delta G}) \quad (9)$$

This probability for a reaction e can also be argued for using the principle of local detailed balance between the rates of the reaction and its reverse where

$$\frac{k_+(e)}{k_-(e)} = \exp\left(-\frac{x_e^{\Delta G}}{k_B T}\right)$$

For a Markovian process describing the sequence of reaction events along the pathway, the probability of observing the specific trajectory over time satisfies

$$\begin{aligned} p\left(\bigwedge_{e: f_e > 0} e\right) &= \prod_{e: f_e > 0} p(e) \\ &\sim \prod_{e: f_e > 0} \exp(-x_e^{\Delta G}) \\ &= \exp\left(-\sum_{e: f_e > 0} x_e^{\Delta G}\right) \end{aligned}$$

Thus, for paths biased in the direction of the pathway (compared to the reverse direction), $-x_e^{\Delta G}$ accumulates additively in the exponent for the probability. Since, $\forall(x, y): x \leq y \iff \exp(x) \leq \exp(y)$, maximizing the probability of the pathway is equivalent to maximizing the objective function (8).

$$\max \log \left[p\left(\bigwedge_{e: f_e > 0} e\right) \right] \equiv \max - \sum_{e: f_e > 0} x_e^{\Delta G}$$

Markovian trajectories might be coarse-grained into pathways (sets of reactions with positive hyperflow) under dilute limit, Markovian kinetics, and weak coupling between reactions.

Therefore, in the low-noise, steady-state, weak-coupling limit, maximizing the log-probability of sustaining a pathway with the hyperedges with assigned positive hyperflows decomposes approximately as maximizing a sum of negative of reaction-wise free-energy differences. The RHS of this expression (3.1.2) is the negation of the action associated with producing the desired hyperflow which is correspondingly minimized. The additivity of the objective function can be justified by the convexity of the rate function for perturbations in chemical reaction networks given by the Wentzell-Freidlin theory. [6]

A third way to motivate objective function (8) is to use the driving force for reactions. For a reaction e , the thermodynamic driving force is the negative of the Gibbs free energy difference [7] or $-x_e^{\Delta G}$ (which is closely related to the affinity of reactions [8] and positive for spontaneous reactions). The net driving force along a hyperflow would be $-\sum_{e: f_e > 0} x_e^{\Delta G}$ (as all hyperflows are positive, the net force adds up). Therefore, the objective corresponds to selecting pathways that are most strongly thermodynamically driven, hence most robust to perturbations.

All in all, the objective is defensible as a heuristic, and can be motivated by a physical interpretation—but only under fairly explicit assumptions, rather than a strict consequence of the laws of thermodynamics. It should be seen more as a ranking functional, rather than a dynamical law.

3.2 IMPLEMENTATION

The objective function is a conditional sum for a set of MILP variables. The value of the variable $x_e^{\Delta G}$ contributes to the objective function only if the corresponding hyperedge is used in the pathway. It needs to be converted into a simple linear sum of a set of MILP variables before the MILP solver can evaluate it. Additionally, some constraints such as (5) and (26) in the mathematical formulation of the problem have been specified as implications. Also they need to be linearized before fed into the MILP solver. This section

describes these implementation details for the formulated MILP model in the previous section.

3.2.1 Implementing the Objective function

The objective function minimizes the cumulative sum of the differences in chemical potentials along the pathway, $\min \left(\sum_{e: f_e > 0} x_e^{\Delta G} \right)$, rather than the simple sum of differences in chemical potentials. For all reactions e , $\min \left(\sum_{e \in E} x_e^{\Delta G} \right)$, in the reaction network. For each hyperedge e , a binary variable $z_e \in \{0, 1\}$ is introduced, indicating whether that hyperedge is used in the pathway, by the following constraints

$$z_e = 0 \iff f_e = 0 \quad (10)$$

$$z_e = 1 \iff f_e > 0 \quad (11)$$

We define an additional variable $\bar{x}_e^{\Delta G}$ for each hyperedge.

The assignment of the variables $\bar{x}_e^{\Delta G}$ is done by the following constraints

$$\bar{x}_e^{\Delta G} = 0 \iff z_e = 0 \quad (12)$$

$$\bar{x}_e^{\Delta G} = x_e^{\Delta G} \iff z_e = 1 \quad (13)$$

The objective function is defined as the sum of these $|E|$ variables for all hyperedges in the hypergraph; $\min \left(\sum_{e \in E} \bar{x}_e^{\Delta G} \right)$. Hyperedges that are not utilized by the pathway have $\bar{x}_e^{\Delta G} = 0$ and would not contribute to the value of the objective function, implementing the objective function as a linear expression.

3.2.2 Linearizing the Constraints

The bi-implication constraints, Equations (10) and (11), associating the boolean variable z_e to the edges e is imple-

case of f_e	(14)	(15)	implication for z_e
$f_e = 0$	$z_e \leq 0$	$M \cdot z_e \geq 0$	$z_e = 0$
$f_e = k > 0$	$z_e \leq k$	$M \cdot z_e \geq k$	$z_e = 1$

Table 2: The linear constraints in Equations (14) and (15) are equivalent to the bi-implication constraints in Equations (10) and (11).

mented using the big-M method [9], where $M > 0$ is a large positive integer constant¹

$$\forall e \in E:$$

$$z_e \leq f_e \quad (14)$$

$$M \cdot z_e \geq f_e \quad (15)$$

The expansion of how the added linear constraints in Equations (14) and (15) lead to assignment of z_e satisfying the implication constraints in Equations (10) and (11) is shown in Table 2.

The implication constraint in Equation (5) determines if an hyperedge can be used in the flow depending on the differences in the chemical potentials for the hyperedges $x_e^{\Delta G}$. This is linearized as

$$\forall e \in E:$$

$$x_e^{\Delta G} + M \cdot (z_e - 1) \leq 0 \quad (16)$$

The equivalence of the added linear constraints in Equation (16) to the implication constraint in Equation (5) is shown in Table 3.

The bi-implication constraints in Equations (12) and (13) assigns the variable $\bar{x}_e^{\Delta G}$ with the differences in the chemical potentials for the hyperedges for the evaluation of the ob-

¹ The constant M has to be larger than the possible values of the variables in the MILP formulation which include the flows f_e for the hyperedges and the chemical potentials $x_v^{G^0}$ for the vertices (in atomic units). In our present implementation $M = 100$ suffices.

case of f_e	implication for z_e	case of $x_e^{\Delta G}$	(5) satisfied?	(16) satisfied?
$f_e = 0$	$z_e = 0$	$x_e^{\Delta G} > 0$	True	True
$f_e = 0$	$z_e = 0$	$x_e^{\Delta G} \leq 0$	True	True
$f_e > 0$	$z_e = 1$	$x_e^{\Delta G} > 0$	False	False
$f_e > 0$	$z_e = 1$	$x_e^{\Delta G} \leq 0$	True	True

Table 3: The linear constraint in Equation (16) is equivalent to the implication constraint in Equation (5).

jective function. By utilizing the variables z_e , we implement the constraints as

$$\forall e \in E:$$

$$x_e^{\Delta G} - \bar{x}_e^{\Delta G} \leq M(1 - z_e) \quad (17)$$

$$x_e^{\Delta G} - \bar{x}_e^{\Delta G} \geq -M(1 - z_e) \quad (18)$$

$$\bar{x}_e^{\Delta G} \leq M \cdot z_e \quad (19)$$

$$\bar{x}_e^{\Delta G} \geq -M \cdot z_e \quad (20)$$

The expansion of how the added linear constraints in Equations (17), (18), (19), and (20) lead to desired assignment of $\bar{x}_e^{\Delta G}$ satisfying the implication constraints in Equations (12) and (13) is shown in Table 4.

Case of z_e	Case of $x_e^{\Delta G}$	Constraint on $\bar{x}_e^{\Delta G}$ by				Implication for $\bar{x}_e^{\Delta G}$
		(17)	(18)	(19)	(20)	
$z_e = 0$	$x_e^{\Delta G} > 0$			≤ 0	≥ 0	$= 0$
$z_e = 0$	$x_e^{\Delta G} \leq 0$			≤ 0	≥ 0	$= 0$
$z_e = 1$	$x_e^{\Delta G} > 0$			Cannot happen		
$z_e = 1$	$x_e^{\Delta G} \leq 0$	$\geq x_e^{\Delta G}$	$\leq x_e^{\Delta G}$			$= x_e^{\Delta G}$

Table 4: The linear constraints of Equation (17), (18), (19), and (20) assign $\bar{x}_e^{\Delta G}$ according to the bi-implication constraints in Equations (12) and (13). Note that the third case cannot happen due to Equation (5), because $x_e^{\Delta G} > 0 \implies f_e = 0 \implies z_e = 0$.

3.2.3 Implemented Model

For an overview, we here list the entire set of variables of the model in Table 6 and the constants used in the model in Table 5. We also summarize the mixed integer linear

program developed in the earlier sections. This model is what is given to the MILP solver.

$$\min \left(\sum_{e \in E} \bar{x}_e^{\Delta G} \right) \quad (21)$$

$$\forall v \in V: \sum_{e \in \bar{E}} s_{ve}^+ f_e - \sum_{e \in \bar{E}} s_{ve}^- f_e = 0 \quad (2)$$

$$\text{Structural specification constraints on } f_e \text{ for } e \in \bar{E} \quad (3)$$

$$\forall e \equiv (e^-, e^+) \in E:$$

$$0 \leq f_e \quad (4)$$

$$\begin{aligned} x_e^{\Delta G} = & \left(\sum_{v \in e^+} x_v^{G^0} - \sum_{v \in e^-} x_v^{G^0} \right) \\ & + RT \cdot \left(\sum_{v \in e^+} x_v^K - \sum_{v \in e^-} x_v^K \right) \end{aligned} \quad (6)$$

$$0 \geq x_e^{\Delta G} + M \cdot (z_e - 1) \quad (16)$$

$$M(1 - z_e) \geq x_e^{\Delta G} - \bar{x}_e^{\Delta G} \quad (17)$$

$$-M(1 - z_e) \leq x_e^{\Delta G} - \bar{x}_e^{\Delta G} \quad (18)$$

$$M \cdot z_e \geq \bar{x}_e^{\Delta G} \quad (19)$$

$$-M \cdot z_e \leq \bar{x}_e^{\Delta G} \quad (20)$$

$$\forall v \in V:$$

$$0 \leq t_v \leq |V| \quad (15)$$

$$\left[x_v^K \right]_{\min} \leq x_v^K \leq \left[x_v^K \right]_{\max} \quad (7)$$

The computational complexity of finding MILP solutions is in general NP-hard [10], and even the restricted case of finding chemical pathways remains so [11]. Despite this worst case complexity, modern MILP solvers efficiently handle most practically relevant instances in reasonable time. From a thermodynamic perspective, it is often valuable to explore *multiple* optimal or near-optimal solutions—however, MILP solvers generally only provide a single optimal solution. Some solvers, e.g., IBM ILOG CPLEX, do provide a so-called solution pool facility for enumerating

Constant	datatype	description
M	integer	a big integer used for linearizing the constraints (type promoted to float, if required)
$x_v^{G^0}$	float	chemical potential of a vertex v under standard physical conditions as assigned by the thermodynamic oracle
$x_e^{\Delta G^0}$	float	differences in chemical potential of in-vertices and out-vertices of a hyperedge e under standard physical conditions

Table 5: Constants in the proposed MILP formulation. The constants M and $x_v^{G^0}$ are provided as input, while the constants $x_e^{\Delta G^0}$ are calculated based on the given $x_v^{G^0}$.

MILP variable	datatype	description
f_e	integer	flow variable for a hyperedge e non-negative
z_e	boolean	indicator variable for a hyperedge e denoting if it is used in the flow
$x_e^{\Delta G}$	float	differences in chemical potential of in-vertices and out-vertices of a hyperedge e after accounting for their concentrations
$\bar{x}_e^{\Delta G}$	float	copy of $x_e^{\Delta G}$ used in the (linearized) objective function
x_v^K	float	log concentration of a vertex v
t_v	integer	topological ordering number for vertex v , non-negative

Table 6: Variables in the proposed MILP formulation.

multiple solutions, but there is little user-control for how it is populated, and in particular how solutions are compared for equality.

The thermodynamics module described in this contribution is implemented as an extension of the larger software package MØD [12], available at <http://mod.imada.sdu.dk/>. The package offers the option of using multiple MILP solvers, currently IBM ILOG CPLEX, Gurobi, and Coin-OR CBC. For the results presented here we have used Coin-OR CBC [13, 14]. MØD implements a MILP enumeration algorithm on top of the solver in use, where one can specify a set of integer or binary variables to enumerate over. This enumeration in MØD proceeds as tree search algorithm over the domains of this set of variables, with calls to the

MILP solver for evaluating feasibility obtaining optimal solutions of subproblems [15].

In the MILP model presented here, there are variables with a continuous domain representing the concentrations and chemical potentials of the molecules. We do not consider minor changes in these variables to represent a new solution, as long as the underlying flow is unchanged. Hence, we carry out the enumeration over the variables $f_e, \forall e \in E$. The resulting output is a list of different integral flows, listed in the order of increasing value of the objective function (21).

3.3 CONTEXTUALIZING THIS METHOD

In this short chapter, we give a detailed comparison of our methods with some existing frameworks. One of the closest is *thermodynamics-based metabolic flux analysis* (TMFA) [16] which also utilizes mixed integer linear constraints. Despite similarities in parts of the thermodynamic modeling, the problem solved is different, with some of the key differences being:

- TMFA generates *flux distributions* not containing thermodynamically infeasible reactions, while our method searches for *explicitly specified pathways* not containing thermodynamically unfavorable reactions. The flux vector generated by TMFA is real-valued as opposed to integer values for the flows in the pathways determined by our method. In many settings, the concept of a pathway involves an integer number of molecules participating in each reaction of the pathway, which aligns well with integer flows.
- TMFA uses a curated dataset of metabolites with chemical potentials determined by group contribution methods. In our case a semi-empirical method (GFN2-xTB) is used to assign chemical potentials to the molecules in the reaction network. However, it can be substituted by any suitable ‘oracle’ that maps a molecule to its Gibbs free energy of the user choice.

This provides flexibility and a wider applicability to this method.

- We allow for a ranking of the enumerated results based on thermodynamics-based metrics. In terms of the linear program modeling, restricting the flow variables to integers allows us to enumerate solutions over the integral variables (i.e., to generate not just a single solution, but many). Additionally, we can enforce a partial ordering of the vertices in the returned integer hyperflow solutions, in order to guarantee their realizability as synthesis plans.

We now turn to a comparison with some of the other existing frameworks. Formulating the problem as an MILP allows us to leverage the many algorithmic advances made in the field of MILP solvers. This may be one reason why we are able to analyze larger reaction networks than those analyzed in [17–19] involving a smaller number of distinct molecules. A further difference to these works is that they utilize kinetic rate equations derived from the law of mass action (reactions under kinetic control), while our focus is on the differences in chemical potentials of the molecules in the reaction network near equilibrium (reactions under thermodynamics control).

Weighted directed graphs are utilized in [20] to model the thermodynamic phase space of a reaction network. However, instead of a constant positive edge weight derived from the Gibbs free energy of the reaction, we utilize the free energies themselves to allow for an analysis more directly based on thermodynamics. Several other cheminformatics tools and automated protocols for computationally exploring reaction networks utilize reactive molecular dynamics simulations [21–23] or chemical heuristics [24–27]. However, to the best of our knowledge, there does not exist any computational framework for explicitly specifying and searching for pathways in reaction networks while simultaneously incorporating thermodynamic constraints in the present form.

REFERENCES

- [1] Gavin E. Crooks. “Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences”. In: *Phys. Rev. E* 60 (3 Sept. 1999), pp. 2721–2726. DOI: [10.1103/PhysRevE.60.2721](https://doi.org/10.1103/PhysRevE.60.2721). URL: <https://link.aps.org/doi/10.1103/PhysRevE.60.2721>.
- [2] Gavin E. Crooks. “Path-ensemble averages in systems driven far from equilibrium”. In: *Phys. Rev. E* 61 (3 Mar. 2000), pp. 2361–2366. DOI: [10.1103/PhysRevE.61.2361](https://doi.org/10.1103/PhysRevE.61.2361). URL: <https://link.aps.org/doi/10.1103/PhysRevE.61.2361>.
- [3] Takahiro Sagawa and Masahito Ueda. “Generalized Jarzynski Equality under Nonequilibrium Feedback Control”. In: *Phys. Rev. Lett.* 104 (9 Mar. 2010), p. 090602. DOI: [10.1103/PhysRevLett.104.090602](https://doi.org/10.1103/PhysRevLett.104.090602). URL: <https://link.aps.org/doi/10.1103/PhysRevLett.104.090602>.
- [4] Benoit Palmieri and David Ronis. “Jarzynski equality: Connections to thermodynamics and the second law”. In: *Phys. Rev. E* 75 (1 Jan. 2007), p. 011133. DOI: [10.1103/PhysRevE.75.011133](https://doi.org/10.1103/PhysRevE.75.011133). URL: <https://link.aps.org/doi/10.1103/PhysRevE.75.011133>.
- [5] Davide Carbone et al. “Efficient Training of Energy-Based Models Using Jarzynski Equality”. In: *Advances in Neural Information Processing Systems*. Ed. by A. Oh et al. Vol. 36. Curran Associates, Inc., 2023, pp. 52583–52614. URL: https://proceedings.neurips.cc/paper_files/paper/2023/file/a4ddb865e0a8ca3cca43fd7387b4b0da-Paper-Conference.pdf.
- [6] Andrea Agazzi, Amir Dembo, and Jean-Pierre Eckmann. *Large deviations theory for Markov jump models of chemical reaction networks*. 2017. eprint: [1701.02126](https://arxiv.org/abs/1701.02126). URL: <https://arxiv.org/abs/1701.02126>.
- [7] “driving force”. In: (2025). DOI: [doi:10.1351/goldbook.D01860](https://doi.org/10.1351/goldbook.D01860). URL: <https://doi.org/10.1351/goldbook.D01860>.

- [8] “affinity of reaction”. In: (2025). DOI: [doi:10.1351/goldbook.A00178](https://doi.org/10.1351/goldbook.A00178). URL: <https://doi.org/10.1351/goldbook.A00178>.
- [9] M. Cococcioni and L. Fiaschi. “The Big-M method with the numerical infinite M”. In: *M. Optim Lett* 15 (Sept. 2020), pp. 2455–2468. DOI: <https://doi.org/10.1007/s11590-020-01644-6>. URL: <https://doi.org/10.1007/s11590-020-01644-6>.
- [10] R. M. Karp. “Reducibility among combinatorial problems”. In: *Complexity of Computer Computations*. Ed. by R. E. Miller and J. W. Thatcher. NY: Plenum Press, 1972, pp. 85–103.
- [11] J.L. Andersen et al. “Maximizing output and recognizing autocatalysis in chemical reaction networks is NP-complete”. In: *J Syst Chem* 3.1 (Jan. 2012). DOI: [10.1186/1759-2208-3-1](https://doi.org/10.1186/1759-2208-3-1). URL: <https://doi.org/10.1186/1759-2208-3-1>.
- [12] Jakob L. Andersen et al. “A Software Package for Chemically Inspired Graph Transformation”. In: *Graph Transformation*. Ed. by Rachid Echahed and Mark Minas. Cham: Springer International Publishing, 2016, pp. 73–88. ISBN: 978-3-319-40530-8. DOI: [10.1007/978-3-319-40530-8_5](https://doi.org/10.1007/978-3-319-40530-8_5).
- [13] John Forrest et al. *coin-or/Cbc: Release releases/2.10.12*. Version releases/2.10.12. Aug. 2024. DOI: [10.5281/zenodo.13347261](https://doi.org/10.5281/zenodo.13347261). URL: <https://doi.org/10.5281/zenodo.13347261>.
- [14] R. Lougee-Heimer. “The Common Optimization Interface for Operations Research: Promoting open-source software in the operations research community”. In: *IBM J. Res. Dev.* 47.1 (Jan. 2003), pp. 57–66. ISSN: 0018-8646. DOI: [10.1147/rd.471.0057](https://doi.org/10.1147/rd.471.0057). URL: <https://doi.org/10.1147/rd.471.0057>.
- [15] Jakob L. Andersen et al. “Chemical Transformation Motifs—Modelling Pathways as Integer Hyperflows”. In: *IEEE/ACM Trans. Comput. Biol. Bioinformatics* 16.2 (Apr. 2019), pp. 510–523. ISSN: 1545-5963. DOI: [10.1109/TCBB.2019.2918888](https://doi.org/10.1109/TCBB.2019.2918888).

- 1109/TCBB.2017.2781724. URL: <https://doi.org/10.1109/TCBB.2017.2781724>.
- [16] Christopher S. Henry, Linda J. Broadbelt, and Vassily Hatzimanikatis. "Thermodynamics-Based Metabolic Flux Analysis". In: *Biophysical Journal* 92.5 (2007), pp. 1792–1805. ISSN: 0006-3495. DOI: <https://doi.org/10.1529/biophysj.106.093138>. URL: <https://www.sciencedirect.com/science/article/pii/S0006349507709876>.
- [17] A.M. Miangolarra and M Castellana. "On Non-ideal Chemical-Reaction Networks and Phase Separation." In: *J Stat Phys* 190.23 (Nov. 2023). DOI: [10.1007/s10955-022-03037-8](https://doi.org/10.1007/s10955-022-03037-8). URL: <https://doi.org/10.1007/s10955-022-03037-8>.
- [18] Tim Schmiedl and Udo Seifert. "Stochastic thermodynamics of chemical reaction networks". In: *The Journal of Chemical Physics* 126.4 (Jan. 2007), p. 044101. ISSN: 0021-9606. DOI: [10.1063/1.2428297](https://doi.org/10.1063/1.2428297). URL: <https://doi.org/10.1063/1.2428297>.
- [19] Anselmo García Cantú and Gregoire Nicolis. "Toward a Thermodynamic Characterization of Chemical Reaction Networks". In: *Journal of Non-Equilibrium Thermodynamics* 31.1 (2006), pp. 23–46. DOI: [doi:10.1515/JNETDY.2006.003](https://doi.org/10.1515/JNETDY.2006.003). URL: <https://doi.org/10.1515/JNETDY.2006.003>.
- [20] Matthew J. McDermott, Shyam S. Dwaraknath, and Kristin A. Persson. "A graph-based network for predicting chemical reaction pathways in solid-state materials synthesis." In: *Nat Commun* 12 (3097 Apr. 2021). DOI: [10.1038/s41467-021-23339-x](https://doi.org/10.1038/s41467-021-23339-x). URL: <https://doi.org/10.1038/s41467-021-23339-x>.
- [21] Emilio Martínez-Núñez et al. "AutoMeKin2021: An open-source program for automated reaction discovery". In: *Journal of Computational Chemistry* 42.28 (2021), pp. 2036–2048. DOI: <https://doi.org/10.1002/jcc.26734>. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/jcc.26734>.

- [22] Robin J. Shannon et al. “ChemDyME: Kinetically Steered, Automated Mechanism Generation through Combined Molecular Dynamics and Master Equation Calculations”. In: *Journal of Chemical Theory and Computation* 17.8 (2021). PMID: 34283599, pp. 4901–4912. DOI: [10.1021/acs.jctc.1c00335](https://doi.org/10.1021/acs.jctc.1c00335). URL: <https://doi.org/10.1021/acs.jctc.1c00335>.
- [23] Malte Döntgen et al. “Automated Discovery of Reaction Pathways, Rate Constants, and Transition States Using Reactive Molecular Dynamics Simulations”. In: *Journal of Chemical Theory and Computation* 11.6 (2015). PMID: 26575551, pp. 2517–2524. DOI: [10.1021/acs.jctc.5b00201](https://doi.org/10.1021/acs.jctc.5b00201). URL: <https://doi.org/10.1021/acs.jctc.5b00201>.
- [24] Mengjie Liu et al. “Reaction Mechanism Generator v3.0: Advances in Automatic Mechanism Generation”. In: *Journal of Chemical Information and Modeling* 61.6 (2021). PMID: 34048230, pp. 2686–2696. DOI: [10.1021/acs.jcim.0c01480](https://doi.org/10.1021/acs.jcim.0c01480). URL: <https://doi.org/10.1021/acs.jcim.0c01480>.
- [25] Alon Grinberg Dana et al. “Automated reaction kinetics and network exploration (Arkane): A statistical mechanics, thermodynamics, transition state theory, and master equation software”. In: *International Journal of Chemical Kinetics* 55.6 (2023), pp. 300–323. DOI: <https://doi.org/10.1002/kin.21637>. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/kin.21637>.
- [26] Tom A. Young et al. “autodE: Automated Calculation of Reaction Energy Profiles— Application to Organic and Organometallic Reactions”. In: *Angewandte Chemie International Edition* 60.8 (2021), pp. 4266–4274. DOI: <https://doi.org/10.1002/anie.202011941>. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/anie.202011941>.
- [27] Kevin M. Shebek et al. “Pickaxe: a Python library for the prediction of novel metabolic reactions.” In: *BMC Bioinformatics* 24.106 (Mar. 2023). DOI: [10.1186/](https://doi.org/10.1186/)

s12859 - 023 - 05149 - 8. URL: <https://doi.org/10.1186/s12859-023-05149-8>.

A CASE STUDY

This chapter is identical to Section 4 of the published version of the preprint at [arXiv:2411.15900](https://arxiv.org/abs/2411.15900).

In this chapter, we demonstrate the methods from Chapter 3 on a reaction network involving hydrogen cyanide, formamidic acid, and formamide in aqueous solution which has been studied in [1].

4.1 NETWORK GENERATION

The reaction network was generated using MØD by the expansion of the molecular space with the input molecules HCN, NH₃, and H₂O, by the recursive application of the graph transformation rules listed in Figure 21. The individual molecules in the generated network, such as HCN, NH₃, H₂O, formamide and so on, are represented as undirected labeled graphs. Stereo-information is not represented. Therefore, both the stereoisomers of the molecule shown in Figure 20 are represented by the same graph. Hence, they cannot be distinguished in our modeling despite them being two different chemical entities. Molecular graphs also constitute the vertices of the hypergraph representing the reaction network.

The graph transformation rules in Figure 21 represent classes of reactions, i.e. reaction templates. The left hand side of the rules is a subgraph, which may be substituted by the subgraph on the right hand side, when a match

The atoms represented as vertices, labeled by the symbol of the element it represents, such as C for a carbon atom, H for a hydrogen atom, N for a nitrogen atom, O for an oxygen atom and so on. The bonds are represented as edges, labeled by bond types such as a single, double, triple or aromatic bond. The default bondtype is single.

has been found in a molecular graphs present in the reaction network. This executes a reaction, thereby expanding the molecular space. Following the previous study of the network [1], the expansion was constrained to only generate molecules with a maximum of three carbon, three nitrogen, and three oxygen atoms. Moreover, only stable molecules ($x_v^{G^0} \leq 0$) were kept. The generated reaction network consists of 67 vertices and 202 hyperedges, and is shown in Figure 22 for completeness. A list of all molecules and transitions in the reaction network can be found in the associated GitHub repository [2].

4.2 THERMODYNAMIC ORACLE

Given a reaction network, we assign chemical potentials to the molecules using what we had earlier referred to as a ‘thermodynamic oracle’ in Chapter 3. Utilizing quantum mechanical methods based on density functional theory (DFT) as this ‘thermodynamic oracle’ quickly becomes impractical as the number of electrons in the molecule (denoted by N) increases, as indicated by the comparative computational costs of the different available methods as shown in Figure 23. Calculating the electron density in a molecule using DFT is an iterative process. Each step invokes gradient descent, which scales beyond $\mathcal{O}(N^3)$ for classical DFT based methods for a single-point calculation because it involves matrix inversions. After the electron density for the molecule has been optimized, it can then be used to calculate the electronic contribution to the energy of a molecule. Machine learnt relations for potential as-







Figure 20: Two stereoisomers of a molecule (left) and the corresponding molecular graph (right).




(a) Addition to a cyanide: in this rule $x \in \{\text{N}, \text{O}\}$, representing the addition of an amide nitrogen or hydroxyl oxygen to the nitrile bond respectively.



(b) Addition to an imine or carbonyl: in this rule $x, y \in \{\text{N}, \text{O}\}$, representing the addition of an amide nitrogen or hydroxyl oxygen to an amide or hydroxyl bond respectively. This single rule encapsulates $2^2 = 4$ individual rules depending on the instantiation of x and y .



(c) 1,3 hydrogen shift: in this rule $x, y \in \{\text{C}, \text{N}, \text{O}\}$, representing the shift of a double bond as in tautomerism. When $x \equiv \text{O}$ and $y \equiv \text{O}$ it is keto-enol tautomerism, $x \equiv \text{N}$ and $y \equiv \text{N}$ is imine-enamine tautomerism, $x \equiv \text{N}$ and $y \equiv \text{O}$ is amine-iminol tautomerism and $x \equiv \text{C}$ and $y \equiv \text{C}$ is an allylic shift of the double bond. This single rule encapsulates $3 \times 3 = 9$ individual rules depending on the instantiation of x and y .



(d) Condensation: in this rule $x \in \{\text{C}, \text{N}, \text{O}\}$, $y, z \in \{\text{N}, \text{O}\}$ representation of the addition of a double bond (of an enol, enamine) to an imine or a carbonyl group. This single rule encapsulates $3 \times 2 \times 2 = 12$ individual rules depending on the instantiation of x , y and z .

Figure 21: The graph transformation rules used to expand the molecular space and generate the network.




Figure 22: An abstract representation of the generated reaction network with 67 vertices (blue circles) and 202 hyperedges (green squares), on which the working of the method proposed in Chapter 3 was demonstrated.

segment exist, some require optimized atomic coordinates to determine potentials [4], while others just utilize the connectivity of the molecular graph [5–8] or even a string representation of it [9]. However, these machine learnt relations from a dataset might not be well generalizable; they could perform poorly when a molecule sufficiently different, from those in the dataset trained on, is encountered [10–12]. Ideally, kinetic rate parameters would have to be incorporated into the search for pathways in a reaction network as integer flows, because they dictate which reactions actually occur in the system and how fast. However, calculating activation free energies using DFT-based quantum mechanical methods to determine these kinetic parameters would demand even further time and computational resources.




Figure 23: A comparison of the computational costs of the available methods often used to assign free energies to molecules along with their accuracy. The rough ranges for the data used to draw the figure was collated from [3].




Figure 24: In a group of not too diverse reactions, the energy barrier of a reaction can be considered to be dependent on the free energy change of the reaction by some function: $G_e = f(x_e^{\Delta G})$. This holds because the reaction trajectories do not tend to cross each other when plotted normalized by the energy of the reactants.

As an alternative, we in this work adopt a compromise oracle based on equilibrium thermodynamics. This approximation has some theoretical basis behind it with the Evans–Polanyi–Semenov principle [13], which states that within a class of reactions, the thermodynamics of the reaction acts as a proxy for the kinetics of the reaction, as shown in Figure 24. It allows us to substitute the energy barrier for a reaction with the free energy difference for a reaction resulting in the relation in Equation (9). Specifically, to balance accuracy, efficiency, generalizability, and running time, we employ a semi-empirical method, GFN2-xTB [14], for potential assignment. While this method compromises on accuracy compared to DFT, the errors are acceptable for demonstrating the utility of the mathematical model on an example reaction network. We detail the thermodynamic oracle we use to assign the chemical potentials to the molecules, $x_v^{G^0}$, in the reaction network in the following section. An advantage of this modular approach is that it provides the reader the freedom to substitute the proposed thermodynamic oracle with another tool, for example pre-trained neural networks, mapping a molecule to its chemical potential (or even the reactions to their kinetics parameters) depending on the desired accuracies and available computational resources.

4.3 ON ASSIGNING THE CHEMICAL POTENTIALS

One of the main enhancements to the thermodynamic modeling of reaction networks in this work is a more robust, transparent and generic method of assigning the chemical potentials under standard physical conditions for molecules, $x_v^{G^0}$, compared to relying on thermodynamic databases such as (<https://pytfa.readthedocs.io/en/latest/thermoDB.html>) using the group contribution method, to infer the free energy change for reactions, $x_e^{G^0}$, as done in [15]. Using quantum chemical methods to assign these chemical potentials quickly becomes computationally resource-intensive because the time complexity scales as




Figure 25: The sequence of operations that was used to assign chemical potentials to the molecules, represented by vertices, in the reaction network, which is referred to thermodynamic oracle in the text.

$\mathcal{O}(N^3)$. The density-functional theory (DFT) calculations involve diagonalization of the $N \times N$ Hamiltonian matrix. The memory requirement scales as $\mathcal{O}(N^2)$ because of the number of two-electron integrals contributing to the energy of the system. Moreover, it is an iterative process which is repeated until an acceptable convergence criteria is reached.

Efficient software design necessitates striking a balance among the components in this work: expanding the molecular space, assigning chemical potentials to the molecules in the space and solving the formulated MILP for queried pathways in the network. If DFT calculations are employed, the calculation of chemical potentials required a significantly longer time compared to the remaining two components. Alternative semi-empirical methods offered a faster yet reasonable accurate way to assign chemical potentials to molecules. Therefore, we utilized xTB [14] to estimate the chemical potentials for the molecules in the expanded reaction network. In the rest of this Section, we describe how




Figure 26: OpenBabel's `WeightedRotorSearch()` iteratively samples conformers to reach towards the lowest energy conformer of the molecule. It randomly chooses from the allowed torsion angles, and reweights the choice based on the energy of the last generated lowest energy conformer. Progressively, the generated conformer has increasingly lower energy.

we estimate the chemical potentials for the molecules from their representations as graphs which is also illustrated in Scheme 25.

The molecules in the expanded reaction network were modeled as graphs where individual vertices represented atoms (labeled by the symbol of the corresponding element), while edges represented bonds. The adjacency matrix of this graph was utilized to determine the connectivity for that molecule. Given a connectivity matrix that satisfies the valences of each atom in the molecule, a three-dimensional embedding for the molecule was generated using the VSEPR theory. The `OBBuilder` class from the Open Babel [16, 17] library was employed to generate the three dimensional coordinates for the atoms utilizing a combination of VSEPR rules and commonly encountered fragments.

The generated geometry of the molecule was refined using 1000 iterations of conjugate gradient descent or until the change in energy (in atomic units, calculated using the universal force field [18]) between successive iterations is less than 10^{-4} . Next, `OBForceField::WeightedRotorSearch()` was utilized to systematically perform a tree-search for a conformer of the molecule with a low energy, by biasing the expansion of the search tree towards the current low energy conformer as shown in Figure 26. At each iterative expansion step, 25 conformers were generated, and 100 steps of conjugate descent are executed on each. This lead to the generation of conformers with successively lower energies over subsequent iterations. The conformer with the lowest energy identified at the end of this search process was further subjected to 250 iterations of conjugate descent or until the energy tolerance of 10^{-5} is reached. The energy for a conformer at each step was estimated using the universal force field [18]. The result of this search process, the configuration of the low-energy conformer, containing the three dimensional Cartesian coordinate of each atom in the molecule was written into an .xyz file.

The .xyz coordinate file generated by OpenBabel serves as an input file for xTB. A geometry optimization of the structure is performed with xTB, employing an energy tolerance of 5×10^{-8} Eh and a gradient tolerance of 5×10^{-5} Eh bohr⁻¹. This is followed by the calculation of the Hessian matrix for the optimized structure, and the chemical potentials at 298.15 K using standard translational, rotational, vibrational, and electronic partition functions from statistical mechanics under the coupled rigid-rotor and harmonic-oscillator approximations.

4.4 ON THE RANGE OF THE CONCENTRATION VARIABLE

Apart from the free energies for the molecules, $x_v^{G^0}$, assigned by the thermodynamic oracle the limits on the log concentration values of the molecules, $[x_v^K]_{\min}$ and $[x_v^K]_{\max}$

have to be assigned by the user. We provide some comments on the choice of these limits in this work. The upper limit for the concentrations is a physical limit bounded by the concentration of the pure substance. Concentration is expressed in the units of moles per liter (mol L^{-1}) for liquids and solutions while in terms of partial pressures (in Pa) for gases. For liquids, the maximum possible concentration value of the reactant would be that of the pure liquid, which is physically constrained by

$$\begin{aligned} 1 \text{ mol of pure liquid} &\equiv M \text{ kg of pure liquid} \\ 1 \text{ mol of pure liquid} &\equiv \frac{M}{\rho} \text{ L of pure liquid} \\ \text{Maximum concentration} &\equiv \frac{\rho}{M} \text{ mol L}^{-1} \end{aligned}$$

For gases, the maximum concentration is imposed by the physical constraint that it has to remain (an ideal) gas following the equation of state $PV = nRT$ so that the equations from the kinetic theory of gases might still be applied to it. The concentration of a gas under some given physical condition of pressure P and temperature T is

$$\frac{n}{V} = \frac{P}{RT}$$

As the pressure is increased, or the temperature is decreased to increase the concentration, the gas ceases to behave as an ideal gas and starts exhibiting properties of the vapor phase. A change in the physical state of the substance would have a free energy of phase reaction involved, and it can no longer be modeled only using the equations considered here.

The lower limit of the reactants is imposed by practical constraints because there are limits to the dilution that can be achieved under laboratory conditions. For a solution, infinite dilution would lead to no solute (reactant) being present, and hence the reaction would not occur. For gases, there are lower limits to pressures that can be achieved in laboratory vacuum settings. Reactions occurring in the solid phase (or on surfaces) are governed by different physical equations and cannot be expressed by this framework.





Figure 27: In the generated network, pathways to the target molecule, a trimer of formamide, were queried to with hydrogen cyanide, water and ammonia and source molecules using the method elaborated in Chapter 3.

For all practical purposes, we can assume our reactants to be solutions or gases satisfying the condition of a well-stirred reactor.

For this work, we have chosen the limits in the concentration variables as $[x_v^K]_{\min} = -3$ and $[x_v^K]_{\max} = 1$. These correspond to physical values of the concentrations between $10^{-3} \text{ mol lit}^{-1}$ and $10^1 \text{ mol lit}^{-1}$. While minimolar concentrations can be achieved under laboratory conditions using standard fluidics procedures, a concentration of tens of molar is often encountered for pure liquids (the concentration of pure water is $55.56 \text{ mol lit}^{-1}$). While we use the same range for the log concentration values for all the molecules in the reaction network, the user might choose to assign different ranges for different molecules to make the model more realistic.



vertex	molecule
v₀	HCN
v₁	H ₂ O
v₂	<chem>HO-CH=NH</chem>
v₃	<chem>O=CH-NH2</chem>
v₄	<chem>O=CH-NH-CH(OH)-NH2</chem>
v₅	<chem>O=CH-NH-CH(OH)-NH-CH(OH)-NH2</chem>

Figure 28: The pathway reported in [1] with the sum of free energies $-178.412 \text{ kJ mol}^{-1}$. This pathway requires four hyperedges (denoted by the squares, with the assigned hyperflow the integer in the square). The vertices in the pathway correspond to molecules, whose structures are listed below the pathway along with the label of the vertex.

The primary objective was to study the energetics of condensation pathways of the smaller seed molecules (HCN, NH_3 , and H_2O) into larger oligomers, which might have implication for the prebiotic synthesis of amino acids and compare it with the pathways reported in [1]. Therefore, the formulated MILP model was set up to query the pathways with HCN, NH_3 , and H_2O as the input molecules and the trimer of formamide (numbered 17 in [1]) as the target molecule as shown in Figure 27. The formulated MILP to search for pathways with the given input and output vertices had 336 variables.

As a sanity check, the enumerated list of flow solutions returned by the solver included the pathway suggested in [1], thought not as the optimal solution when scored according to Equation (21). The sum of free energies of the transitions in this pathway is $-178.412 \text{ kJ mol}^{-1}$, while the sum of free energies weighted by the number of times it is used in the pathway yields $-405.332 \text{ kJ mol}^{-1}$. This pathway uses four unique transitions (hyperedges) as depicted in Figure 28. Two of them are used multiple times, which makes seven transitions in total (sum of flows assigned to the hyperedges). The pathway requires an inflow of three molecules of v_0 (hydrogen cyanide) and three molecules of v_1 (water) to produce the target molecule v_5 . The chemical potential $x_v^{\Delta G}$ and the log concentration x_v^K values assigned to each molecule by the MILP solver is listed in Table 9, along with their molecular graphs.

While thermodynamics might provide an insight to the equilibrium flows in the reaction network, it does not describe the path to that equilibrium. Each transition has an associated barrier energy that must be overcome for the transition to occur. Enumerating pathways using our implementation of the MILP formulation of the problem provides the pathway in Figure 29 as one with the minimum value of the objective function. This pathway utilizes four unique transitions and five transitions in all to realize the pathway, which is fewer than the previously reported pathway. Since the difference in the chemical potential between the source and target molecules is fixed, choosing hyperedges with

vertex	inFlow	outFlow	$x_v^{\Delta G}$	x_v^K
v_0	3	0	-5.507	1.000
v_1	3	0	-5.068	-6.000
v_2	0	0	-10.610	-6.000
v_3	0	0	-10.625	0.195
v_4	0	0	-21.249	-1.190
v_5	0	1	-31.869	-6.000

Table 7: The MILP solver assigns the inflow for the allowed source molecules to generate an outflow of the queried target molecule, along with assigning the chemical potentials, $x_v^{\Delta G}$, and log concentration, x_v^K , values of the molecules in the pathway shown in Figure 28. The $x_v^{\Delta G}$ values are in atomic units (in Hartrees as returned by xTB) while the x_v^K values are the dimensionless equivalent of $\log_{10} \frac{[\text{conc.}]}{1 \text{ mol lit}^{-1}}$, where conc. is the concentration of the molecule in mol lit^{-1} .

hyperedge	f_e	in(v)	out(v)	$\bar{x}_e^{\Delta G}$
e_1	3	v_0, v_1	v_2	-94.141
e_2	2	v_2	v_3	-38.629
e_3	1	v_2, v_3	v_4	-0.006
e_4	1	v_3, v_4	v_5	-45.645
		$\sum_e \bar{x}_e^{\Delta G}$	=	-178.412
		$\sum_e f_e \cdot \bar{x}_e^{\Delta G}$	=	-405.332

Table 8: The MILP solution assigned hyperflows f_e to the hyperedges in the pathway, indicating the number of times they are used to effect the pathway shown in Figure 28. The set of vertices which is the inset of the hyperedge (or the reactants in the corresponding transition) is listed under in(v), while the set of vertices which is the outset of the hyperedge (or the products in the corresponding transition) is listed under out(v). $\bar{x}_e^{\Delta G}$ gives the free energy difference assigned to the hyperedge in kJ mol^{-1} .


larger free energy differences would increase the value of the objective function. Using transition with larger free energy differences to ‘cover’ the net difference in the chemical potential, lead to fewer transitions being used in the pathway.

vertex	inFlow	outFlow	$x_v^{\Delta G}$	x_v^K
v_0	3	0	-5.507	1.000
v_1	3	0	-5.068	-6.000
v_2	0	0	-10.610	1.000
v_3	0	0	-16.159	-6.000
v_4	0	0	-21.240	1.000
v_5	0	1	-31.869	-6.000

Table 9: The MILP solver assigns the inflow for the allowed source molecules to generate an outflow of the queried target molecule, along with assigning the chemical potentials, $x_v^{\Delta G}$, and log concentration, x_v^K , values of the molecules in the pathway shown in Figure 29. The units are the same as in Table 8.

hyperedge	f_e	in(v)	out(v)	$\bar{x}_e^{\Delta G}$
e_1	2	v_0, v_1	v_2	-94.141
e_2	1	v_0, v_2	v_3	-129.554
e_3	1	v_1, v_3	v_4	-38.298
e_4	1	v_2, v_4	v_5	-51.185
		$\sum_e \bar{x}_e^{\Delta G}$	=	-313.178
		$\sum_e f_e \cdot \bar{x}_e^{\Delta G}$	=	-407.319

Table 10: The MILP solution assigned hyperflows f_e to the hyperedges in the pathway, indicating the number of times they are used to effect the pathway shown in Figure 29. The set of vertices which is the inset of the hyperedge (or the reactants in the corresponding transition) is listed under in(v), while the set of vertices which is the outset of the hyperedge (or the products in the corresponding transition) is listed under out(v). $\bar{x}_e^{\Delta G}$ gives the free energy difference assigned to the hyperedge in kJ mol^{-1} .



vertex	molecule
--------	----------

v₀	<chem>HCN</chem>
----------------------	------------------

v₁	<chem>H2O</chem>
----------------------	------------------

v₂	<chem>HO-CH=NH</chem>
----------------------	-----------------------

v₃	<chem>HO-CH=N-CH=NH</chem>
----------------------	----------------------------

v₄	<chem>HO-CH=N-CH(OH)-NH2</chem>
----------------------	---------------------------------

v₅	<chem>O=C-NH-CH(OH)-NH-CH(OH)-NH2</chem>
----------------------	--

Figure 29: The pathway scoring the best when the model proposed in Chapter 3 was applied to the generated reaction network along with the objective function (8) with the simple sum of free energies $-313.718 \text{ kJ mol}^{-1}$. This pathway requires four hyperedges (denoted by the squares, with the assigned hyperflow the integer in the square). The vertices in the pathway correspond to molecules, whose structures are listed below the pathway along with the label of the vertex.




Figure 30: Stepwise trimerization of formamide by iterative addition of the monomer as hypothesised in the conventional pathway shown in Figure 28.

The main source of difference in the objective function value in the two pathways is the chemical potential of the molecule v_3 from Figure 29 that is not present in the conventional pathway. While the conventional pathway in [1] is more symmetric and appealing because of the stepwise nucleophilic attack on the carbonyl C atom by the amine N atom of an adjacent v_3 molecule, which is repeated twice to generate v_5 as illustrated in Figure 30, [1] also notes that these condensation reactions have a positive Gibbs free energy.

To quote the authors,

‘each step is endergonic although decreasingly so. Dimerization of formamide is uphill 19.8 kcal/mol, owing to the thermodynamic stability of the monomers in solution. Addition of another monomer to form the trimer, via similar nucleophilic attack, is uphill 19.0 kcal/mol’.

This can also be observed in the ILP solution in Table 8 for the conventional pathway where the hyperedges $e_3 \equiv (\{v_2, v_3\}, \{v_4\})$ have a $\bar{x}_e^{\Delta G}$ value close to zero. The constraint in Equation (5) forces the hyperedge to have $\bar{x}_e^{\Delta G} \leq 0$ for it to be included in the pathway. The MILP solver chooses the log concentration variables x_v^K for the molecules v_3 and v_4 in such a way so that $\bar{x}_e^{\Delta G}$ variables for e_3 is just below zero to realize the pathway. This might be the reason for x_v^K for these two vertices not having the extremal value as is often the case, as discussed below. Our proposed model also returns an alternative pathway involving v_3




Figure 31: The trimerization of formamide which uses v_3 in the pathway that is returned as the best scoring pathway as shown in Figure 29.

from Figure 31, where none of the $\bar{x}_e^{\Delta G}$ weights for the hyperedges is close to zero as listed in Table 10. This involved stepwise addition of of a HCN molecules followed by a H_2O molecules instead of direct addition of a foramide molecule.

We observe that the log concentration variables are always assigned extremal values decided by the constraint in Equation (7). This is likely an artifact of using an MILP solver with an objective function for the problem. An MILP is a convex optimization problem where minimizing the objective function (involving a subset of the MILP variables) results in a solution located at a vertex of the convex space defined by the constraints. The bounds for the variables are hand-crafted by the user and it might be more realistic to assign different bounds for the log concentration variable for each vertex. Alternatively, a better interpretation of the assigned values for the variables, x_v^K might be the direction in which the equilibrium concentrations have to be perturbed to achieve the desired flow. For example, the concentration values in Table 9 can be interpreted as the concentrations of v_1 , v_3 and v_5 have to be lowered and the concentrations of v_0 , v_2 and v_4 have to be raised to achieve a larger difference in the potentials while effecting a net outflow of molecule v_5 given input molecules v_0 and v_1 .

An exhaustive search of the reaction space led to the identification of 13 distinct reaction pathways [19], listed in Table 11 in the order of increasing value of objective 21. The column list the rank, the total number of reactions




Figure 32: The list of the reactions used in the pathways to the target molecules, with the sequence of specific reactions in each pathway listed in Table 11.

used, the number of unique reactions used, the number of molecules involved, the value of the objective (21) and the sum of the free energy differences of the reactions weighted by the flows in the columns respectively. Note, that the

#	$\sum_e f_e$	$\sum_e z_e$	$ v $	obj (21)	$\sum_e f_e \cdot x_e^{\Delta G}$	rxn sequence
0	5	4	6	-305.697	-399.837	1,3,4,10
1	6	5	7	-306.697	-399.837	1,3,4,5,9
2	6	5	7	-305.696	-399.836	1,2,3,4,11
3	5	3	5	-211.555	-399.836	1,6,9
4	5	3	5	-211.554	-399.836	1,8,10
5	6	4	6	-211.556	-399.836	1,8,5,9
6	6	4	6	-211.556	-399.836	1,2,7,9
7	6	4	6	-211.554	-399.836	1,2,8,11

Table 11: All enumerated pathways to the target molecule in the network, with the sequence of pathways leading to the target molecules. The specific reactions corresponding to the index listed in the sequence of reactions for each pathway listed in the last column can be read from Figure 32.

second column from the right gives the net free energy difference, which is path independent and therefore, the same for all the listed pathways. These pathways were enumerated in 866 seconds on an AMD[®] Ryzen 5 PRO 4650U CPU (2.1 GHz) utilizing 8 threads. One may consult the list of reactions in Figure 32 for the molecules and the exact set of reactions involved in these pathways.

REFERENCES

- [1] Jeremy Kua and Kyra L. Thrush. "HCN, Formamidic Acid, and Formamide in Aqueous Solution: A Free-Energy Map". In: *The Journal of Physical Chemistry B* 120.33 (2016). PMID: 27016454, pp. 8175–8185. DOI: [10.1021/acs.jpcb.6b01690](https://doi.org/10.1021/acs.jpcb.6b01690). URL: <https://doi.org/10.1021/acs.jpcb.6b01690>.
- [2] *supplementary data repository on GitHub*. <https://github.com/AdityaPal/thermodynamicsWithILP.git>. 2024.
- [3] Curtis L Janssen and Ida MB Nielsen. *Parallel computing in quantum chemistry*. CRC press, 2008.

- [4] Roman Zubatyuk et al. "Teaching a neural network to attach and detach electrons from molecules." In: *Nat Commun* 12 (4870 Aug. 2021). DOI: [10.1038/s41467-021-24904-0](https://doi.org/10.1038/s41467-021-24904-0). URL: <https://doi.org/10.1038/s41467-021-24904-0>.
- [5] Frank Noé et al. "Machine Learning for Molecular Simulation". In: *Annual Review of Physical Chemistry* 71. Volume 71, 2020 (2020), pp. 361–390. ISSN: 1545-1593. DOI: <https://doi.org/10.1146/annurev-physchem-042018-052331>. URL: <https://www.annualreviews.org/content/journals/10.1146/annurev-physchem-042018-052331>.
- [6] Juncai Li and Xiaofei Jiang. "Mol-BERT: An Effective Molecular Representation with BERT for Molecular Property Prediction". In: *Wireless Communications and Mobile Computing* 2021.1 (2021), p. 7181815. DOI: <https://doi.org/10.1155/2021/7181815>. eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1155/2021/7181815>. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1155/2021/7181815>.
- [7] Mohammadamin Tavakoli et al. "Quantum Mechanics and Machine Learning Synergies: Graph Attention Neural Networks to Predict Chemical Reactivity". In: *Journal of Chemical Information and Modeling* 62.9 (2022). PMID: 35020394, pp. 2121–2132. DOI: [10.1021/acs.jcim.1c01400](https://doi.org/10.1021/acs.jcim.1c01400). eprint: <https://doi.org/10.1021/acs.jcim.1c01400>. URL: <https://doi.org/10.1021/acs.jcim.1c01400>.
- [8] Obaidur Rahaman and Alessio Gagliardi. "Deep Learning Total Energies and Orbital Energies of Large Organic Molecules Using Hybridization of Molecular Fingerprints". In: *Journal of Chemical Information and Modeling* 60.12 (2020). PMID: 33118351, pp. 5971–5983. DOI: [10.1021/acs.jcim.0c00687](https://doi.org/10.1021/acs.jcim.0c00687). eprint: <https://doi.org/10.1021/acs.jcim.0c00687>. URL: <https://doi.org/10.1021/acs.jcim.0c00687>.
- [9] Gabriel A. Pinheiro et al. "Machine Learning Prediction of Nine Molecular Properties Based on the

- SMILES Representation of the QM9 Quantum-Chemistry Dataset". In: *The Journal of Physical Chemistry A* 124.47 (2020). PMID: 33174750, pp. 9854–9866. DOI: [10.1021/acs.jpca.0c05969](https://doi.org/10.1021/acs.jpca.0c05969). eprint: <https://doi.org/10.1021/acs.jpca.0c05969>. URL: <https://doi.org/10.1021/acs.jpca.0c05969>.
- [10] Derek van Tilborg, Alisa Alenicheva, and Francesca Grisoni. "Exposing the Limitations of Molecular Machine Learning with Activity Cliffs". In: *Journal of Chemical Information and Modeling* 62.23 (2022). PMID: 36456532, pp. 5938–5951. DOI: [10.1021/acs.jcim.2c01073](https://doi.org/10.1021/acs.jcim.2c01073). eprint: <https://doi.org/10.1021/acs.jcim.2c01073>. URL: <https://doi.org/10.1021/acs.jcim.2c01073>.
- [11] Sudarshan Vijay et al. "Limits to scaling relations between adsorption energies?" In: *The Journal of Chemical Physics* 156.23 (June 2022), p. 231102. ISSN: 0021-9606. DOI: [10.1063/5.0096625](https://pubs.aip.org/aip/jcp/article-pdf/doi/10.1063/5.0096625/16544349/231102_1_online.pdf). eprint: https://pubs.aip.org/aip/jcp/article-pdf/doi/10.1063/5.0096625/16544349/231102_1_online.pdf. URL: <https://doi.org/10.1063/5.0096625>.
- [12] Valentin Vassilev-Galindo et al. "Challenges for machine learning force fields in reproducing potential energy surfaces of flexible molecules". In: *The Journal of Chemical Physics* 154.9 (Mar. 2021), p. 094119. ISSN: 0021-9606. DOI: [10.1063/5.0038516](https://pubs.aip.org/aip/jcp/article-pdf/doi/10.1063/5.0038516/15587076/094119_1_online.pdf). eprint: https://pubs.aip.org/aip/jcp/article-pdf/doi/10.1063/5.0038516/15587076/094119_1_online.pdf. URL: <https://doi.org/10.1063/5.0038516>.
- [13] "Bell–Evans–Polanyi principle". In: (2025). DOI: [doi:10.1351/goldbook.B00628](https://doi.org/10.1351/goldbook.B00628). URL: <https://doi.org/10.1351/goldbook.B00628>.
- [14] Christoph Bannwarth et al. "Extended tight-binding quantum chemistry methods". In: *WIREs Computational Molecular Science* 11.2 (2021), e1493. DOI: <https://doi.org/10.1002/wcms.1493>. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/wcms.1493>.


- [15] Pierre Salvy et al. “pyTFA and matTFA: a Python package and a Matlab toolbox for Thermodynamics-based Flux Analysis”. In: *Bioinformatics* 35.1 (July 2018), pp. 167–169. ISSN: 1367-4803. DOI: [10.1093/bioinformatics/bty499](https://doi.org/10.1093/bioinformatics/bty499). URL: <https://doi.org/10.1093/bioinformatics/bty499>.
- [16] N. Yoshikawa and G.R. Hutchison. “Fast, efficient fragment-based coordinate generation for Open Babel.” In: *J Cheminform* 11.49 (Aug. 2019). DOI: [10.1186/s13321-019-0372-5](https://doi.org/10.1186/s13321-019-0372-5). URL: <https://doi.org/10.1186/s13321-019-0372-5>.
- [17] *Open Babel GitHub repository*. <https://github.com/openbabel/openbabel>. 2024.
- [18] A. K. Rappe et al. “UFF, a full periodic table force field for molecular mechanics and molecular dynamics simulations”. In: *Journal of the American Chemical Society* 114.25 (1992), pp. 10024–10035. DOI: [10.1021/ja00051a040](https://doi.org/10.1021/ja00051a040). eprint: <https://doi.org/10.1021/ja00051a040>. URL: <https://doi.org/10.1021/ja00051a040>.
- [19] *list of returned solutions by the MILP solver on GitHub*. <https://github.com/AdittyaPal/thermodynamicsWithILP/blob/5f7aed70fe8706bbe0ffc2efedd68316d44e0f42/listOfSolutions.pdf>. 2025.

REFINEMENTS TO THE MODEL


This chapter is identical to Section 3 of the published version of the preprint at [arXiv:2504.10609](https://arxiv.org/abs/2504.10609).

In this chapter, we develop an objective measure to rank pathways in a reaction network maximizing the probability of the entire pathway based on physical principles. Additionally, based on the notion that energy barriers of reactions are the determining factor for the kinetics of the network, we develop an automated methodology for computing such energy barriers, building on a combination of an array of existing computational tools. A key novelty is that in contrast to the standard manual intervention required to generate reaction trajectories from which the reaction barrier can be estimated, we try to automate this part of the process. Lastly, we formulate an integer linear program which allows the user to specify queries for pathways in the network in a very flexible and generic way. Solving this integer linear program returns all pathways fulfilling the criteria, ranked by the objective measure mentioned above. In short, this gives an automated, general, and flexible methodology for analyzing reaction networks, a methodology which offers a speed-up over manual search for pathways but with comparable quality.


In this chapter, we detail our proposed methodology for automating the search for kinetically informed pathways in reaction networks. As shown in Figure 33, the kinetics of the reaction can often lead to different results than those obtained from the reaction thermodynamics as done in




(a) the thermodynamically more favored reaction are generally also kinetically preferred.



(b) the thermodynamically less favored reaction with a smaller negative Gibbs free energy difference is also kinetically preferred because it has a lower energy barrier.



(c) the thermodynamically unfavored reactions (with a positive Gibbs free energy difference) generally have higher energy barriers and are kinetically not preferred.



(d) however, in fringe cases a thermodynamically unfavored reaction might have a lower energy barrier than a thermodynamically favored reaction.

Figure 33: Four of the many possible cases where the thermodynamics and kinetics of reactions might lead to different results. In cases 33b and 33d, the results based on thermodynamics are expected to be different from those obtained from kinetics.

Chapter 3. In Sections 5.1 and 5.2, we first propose an objective function for ranking pathways in a reaction network annotated with kinetic data in the form of energy barriers for each reaction. In Section 5.3, we then formulate an integer linear program (ILP) representing such networks and allowing the specification and execution of pathway searches. This part builds on the modeling described in Chapter 3. We also describe how to search for not only one, but multiple solutions to a given pathway query, ordered by an objective function such as the one presented in Section 5.2.

5.1 FROM RATE CONSTANTS TO EDGE WEIGHTS

Irrespective of whether one uses Arrhenius' theory [1, 2], collision theory or transition state theory [3], the dependence of the rate constant k_e of a reaction e on the free energy barrier G_e follows that of a negative exponential. The expression for the rate constant by Eyring's equation from transition state theory is

$$k_e = \frac{k_b T}{h} \exp\left(-\frac{G_e}{RT}\right) \quad (22)$$

where R is the universal gas constant, k_b is the Boltzmann constant, and h is the Planck constant, each in their appropriate units. In a well-stirred system where all the reactions in the reaction network are assumed to be concurrently taking place, the temperature can be considered to be the same for all reactions. Hence, it is not a factor affecting the relative rates of the reactions in the system.


Since our modeling is not aimed at capturing temporal changes in the concentrations of molecules, we make the simplifying assumption that all molecules in the reaction network are present with unit concentrations. This avoids the time propagation of the changes in the concentration of the molecules, as some of them are used up as reactants to produce other molecules. Under this assumption, the rate constant k_e can be interpreted as being proportional to the probability of reaction e happening in a fixed small volume

and time interval. A reaction with a higher rate constant (and lower energy barrier) is expected to occur more frequently, hence the probability of that reaction occurring in a fixed interval of time when a limited number of reactant molecules are present would be higher. To interpret the rate constant as probabilities of the reaction in a given reaction network, we normalize these values to sum to one:


$$\begin{aligned}
 p(e) &= \frac{k_e}{\sum_{i \in E} k_i} \\
 &= \frac{\frac{k_b T}{h} \exp\left(-\frac{G_e}{RT}\right)}{\sum_{i \in E} \frac{k_b T}{h} \exp\left(-\frac{G_i}{RT}\right)} && \text{(using 22)} \\
 &= \frac{\frac{k_b T}{h} \exp\left(-\frac{G_e}{RT}\right)}{\frac{k_b T}{h} \sum_{i \in E} \exp\left(-\frac{G_i}{RT}\right)} \\
 &= \frac{\exp\left(-\frac{G_e}{RT}\right)}{\sum_{i \in E} \exp\left(-\frac{G_i}{RT}\right)} \\
 &= \frac{\exp\left(-\frac{G_e}{RT}\right)}{D} && (23)
 \end{aligned}$$

Here, the denominator $\sum_{i \in E} \exp\left(-\frac{G_i}{RT}\right)$ is the same for all reactions e and has been shortened to D . In the current work, we assign the measure $p(e)$ as the weight of the hyperedge representing e in an attempt to score reactions with lower energy barriers favourably. However, the user is free to assign any other measures of weight to the hyperedges in the reaction network, if deemed better suited for a particular pathway search objective.

Figure 34 tries to illustrate the reasoning behind assigning the weight measure to the respective reactions. There are two reactions with energy barriers G_1 and G_2 with $G_2 > G_1$ as shown in Figure 34a. The energy distribution of the reactant molecules for both the reactions is given by the Maxwell-Boltzmann energy distribution in Figure 34b, and the fraction of molecules with the available energy to



(a) Reactions with different energy barriers compete for the limited reactant molecules to form the respective products.




(b) The reaction with the lower energy barrier is accessible to a larger fraction of the reactant molecules as more molecules have the energy required to form the products (denoted the shaded area under the curve).

Figure 34: Of the two reactions competing for the same reactant molecules, the one with a lower energy barrier is favored as more reactant molecules can form the products. Therefore, it has a higher rate (given by Equation (22)) and assigned a higher probability of occurring by Equation (23) in a finite time with limited number of reactant molecules.


overcome the energy barrier of the respective reaction and form the products is given by

$$\begin{aligned}
 f(N) &= \int_{G_i}^{\infty} P(E) dE \\
 &\sim \int_{G_i}^{\infty} \exp\left(-\frac{E}{RT}\right) dE \\
 &= \exp\left(-\frac{G_i}{RT}\right)
 \end{aligned}$$


The rate of the reaction would be expected to be proportional to this fraction of molecules with enough energy



(a) such a pathway 2 would be clearly more favored than pathway 1.



(b) it is difficult to decide which of the pathways is more favored because not all molecules in one pathway have higher energies than the other (the energy profiles of the pathways cross at the second reaction).



(c) it is also difficult to decide if pathway 1 (a shorter pathway with 4 reactions but higher barriers) is more favourable than pathway 2 (a longer pathway with 5 reactions but lower barriers).

Figure 35: An objective function scores pathways and helps to decide which would be more favourable when it is not obvious such as in the cases depicted in 35b and 35c.

to form the respective products, and might therefore, be used as a weight on a reaction to score its inclusion in the pathway. A pathway consisting of reactions with lower energy barriers would be scored higher as a larger fraction of molecules would overcome each reaction in the pathway.

5.2 FORMULATING AN OBJECTIVE FUNCTION

We are interested in pathways from the given source molecules to the desired target molecule(s) that have a high overall probability of occurring in practice. This might not always be easy to determine when there are multiple pathways to the target molecule(s) as shown in Figure 35. Considering individual reactions in a pathway as independent events

(and abusing the notation e to mean the event of reaction e happening), the probability p_{total} of the entire set of reactions in the pathway occurring in one unit of time and one unit of volume is

$$\begin{aligned}
 p_{\text{total}} &= p \left(\bigwedge_{e: f_e > 0} (e \wedge e \wedge \dots f_e \text{ times}) \right) \\
 &= \prod_{e: f_e > 0} p(e)^{f_e} \\
 &= \prod_{e: f_e > 0} \left[\frac{\exp \left(-\frac{G_e}{RT} \right)}{D} \right]^{f_e} && \text{(using 23)} \\
 &= \frac{\prod_{e: f_e > 0} \exp \left(-\frac{f_e \cdot G_e}{RT} \right)}{\prod_{e: f_e > 0} D^{f_e}} \\
 &= \exp \left(-\frac{\sum_{e \in E} f_e \cdot G_e}{RT} \right) / D^{\sum_{e \in E} f_e}
 \end{aligned}$$

Since the energy barrier G_e is weighted by the flow f_e , the conditional sum $e : f_e > 0$ can be substituted by a simple sum over all hyperedges $e \in E$ because hyperedges with $f_e = 0$ do not contribute to the value of the sum. The expression for the total probability of a pathway is a fraction and we take the logarithm of it to simplify it and make it a linear expression.

$$\begin{aligned}
 \log p_{\text{total}} &= \log \left[\exp \left(-\frac{\sum_{e \in E} f_e \cdot G_e}{RT} \right) / D^{\sum_{e \in E} f_e} \right] \\
 &= -\frac{\sum_{e \in E} f_e \cdot G_e}{RT} - \left(\sum_{e \in E} f_e \right) \cdot \log D
 \end{aligned}$$

The last line is proportional to the following expression obtained by multiplying by RT

$$-\sum_{e \in E} f_e \cdot G_e - RT \cdot \log D \cdot \sum_{e \in E} f_e$$

The probability of the set of reactions in the pathway is then maximized by maximizing the value of the obtained expression:

$$\max \left(-\sum_{e \in E} f_e \cdot (G_e + RT \cdot \log D) \right)$$

This is equivalent to the following minimization problem:

$$\min \left(\sum_{e \in E} f_e \cdot (G_e + RT \cdot \log D) \right) \quad (24)$$

This proposed objective value (24) is a linear expression because the G_e 's are constants from the input instance, T is assumed constant, and R is a physical constant. Also, the value $\log D = \log \left(\sum_{i \in E} \exp \left(-\frac{G_i}{RT} \right) \right)$ is constant when considering a fixed reaction network.

We note that via D , the objective value (24) of a pathway will depend on the underlying reaction network in which it is considered embedded. This may be regarded as a weakness. On the other hand, the objective value (24) is intuitively appealing by being composed of two natural components: Its first component $\sum_{e \in E} f_e \cdot G_e$ favors pathways with small sums of energy barriers of the reactions in the pathway, weighted by the number of times it is used. Its second component $\sum_{e \in E} f_e \cdot (RT \log D)$ favors pathways with fewer reactions. The factor $RT \log D$ can be viewed as weighting the relative importance of the two components and decides which of the two pathways is more favourable in cases such as the one shown in Figure 35c.

5.3 ILP MODELING

In this section, we express the modeling of Chapter 1 as an integer linear program (ILP). Recall the key idea that in a reaction network, pathways can be expressed as hyperflows and pathway queries can be expressed as constraints on hyperflows. Such constraints can simply be added to the ILP, after which the pathway queries can be executed by running an ILP-solver on the final ILP. We also describe how to search for multiple, structurally different solutions to a given pathway query, ordered by an objective function such as (24).

Thus, in the ILP a pathway is represented by a vector of non-negative¹ integer-valued variables f_e , one variable for each reaction e in the reaction network. These flow variables are subject to the key constraint (2), as well as to the user-supplied constraints that specify the pathway query under consideration.

To model reversible reactions, separate hyperedges were added for the forward and backward directions of the reaction. This allows us to restrict the flows to non-negative integers only. The use of the reverse of a reaction would be denoted by a positive flow on the hyperedge in the opposite direction.

$$f_e \geq 0 \quad (4)$$

For each reaction e , we add a boolean indicator variable z_e which keeps track of whether the hyperedge e is used in the pathway or not. The values of these variables are assigned according to their respective flow variables following the biimplication constraints:

$$f_e = 0 \iff z_e = 0 \quad (10)$$

$$f_e \geq 1 \iff z_e = 1 \quad (11)$$

We will use these indicator variables later when listing multiple, structurally different solutions to a given pathway query as elaborated in Section 5.3.1 and for assigning a partial order to vertices in the pathway as elaborated in Section 5.3.2.² Many ILP solvers support addition of such indicator constraints (12) and (13) directly as implications to an ILP problem (which are then solved by a nonlinear, nonconvex reformulation of the indicator constraint). If this feature is not available, one can instead use the method of linearizing these implication constraints via a large constant M .

In summary, the resulting ILP looks as follows:

- ¹ Non-negative flow values require reversible reactions to be modeled as two separate hyperedges in the reaction network, but it also allows the flexibility to model some reactions as irreversible.
- ² The indicator variables can also increase expressiveness when specifying pathway queries.

Constant	Datatype	Description
G_e	float	Weight of the hyperedge e
s_{ve}^+	integer	Count of molecule v on the LHS of a reaction e
s_{ve}^-	integer	Count of molecule v on the RHS of a reaction e
RT	float	Product of universal gas constant and temperature
M	integer	Big integer used for linearizing the constraints (type promoted to float, if required)

Table 12: Constants in the proposed ILP formulation.

ILP variable	Datatype	Description
f_e	integer	Flow variable for a hyperedge e non-negative
z_e	boolean	Indicator variable denoting if a hyperedge e is used in the flow

Table 13: Variables in the proposed ILP formulation.

$$\min \left(\sum_{e \in E} (f_e \cdot G_e + RT \cdot \log D \cdot f_e) \right) \quad (24)$$

$$\forall e \in E: 0 \leq f_e \quad (4)$$

$$\forall v \in V: \sum_{e \in \bar{E}} s_{ve}^+ f_e - \sum_{e \in \bar{E}} s_{ve}^- f_e = 0 \quad (2)$$

Constraints specifying the pathway query such as, inflow and outflow constraints

$$f_e = 0 \iff z_e = 0 \quad (10)$$

$$f_e \geq 1 \iff z_e = 1 \quad (11)$$

The constraints specifying the pathway consist of the desired outflow of the queried target molecule and the supplied source molecules. Additionally, it might involve constraints such as disallowing particular molecules as by-products (by restricting their outflow to zero), or disallowing the use of particular reactions (by specifying $f_e = 0 = z_e$




Figure 36: An example of a solution (the hyperflow of the entire figure) which contains a previous solution (the hyperflow shown in solid blue) as a subset, seen in terms of the edges (reactions) used. Adding the faded hyperedges to the integer hyperflow depicted in solid blue creates a ‘null cycle’ which creates a hyperflow solution that satisfies constraint (2) but does not create a physically different solution. Addition of constrain (25) for all previously returned solutions ensures that such solutions are not returned.

for those reactions). An overview of the variables and constants in the ILP can be found in Table 13 and 12, respectively.

5.3.1 Enumeration of solutions

We would like to enumerate multiple solutions to a pathway query in the reaction network. In this enumeration, we prefer to get a diverse set of pathways, in order to make the result more robust to uncertainties in the weights (i.e., the energy barrier values) assigned to the hyperedges. Such diversity offers a selection of alternative pathways for implementation, which is beneficial in case the single optimal solution turns out to be unattainable in practice (or just less desirable than expected due to circumstances not included in the modeling). Some ILP solvers provide so-called solution pools which allow enumeration of solutions, but the process of populating this pool of solutions

is often opaque. More control is given in MØD [4], which provides an option to enumerate multiple solutions based on a user-supplied specification of which subset of integer and binary variables to enumerate over [5]. For example, if the specification lists exactly the variables z_e for each $e \in E$, then two pathways using the same set of hyperedges (but potentially with a difference in the flow value on some edges) will be considered equal, and only the one with the best objective value is returned. That is, if we define $\text{supp}(f) = \{e \in E \mid z_e = 1\}$ then two solutions f_1 and f_2 are considered different exactly when $\text{supp}(f_1) \neq \text{supp}(f_2)$.


However, in our case initial experimentation showed that using this enumeration policy, later enumerated solutions f_2 often contained a previous solution f_1 , in the sense that $\text{supp}(f_1) \subsetneq \text{supp}(f_2)$, as exemplified in Figures 36 and 37. To increase diversity of solutions, we opted for a custom enumeration scheme where $\text{supp}(f_2)$ of new solution f_2 cannot contain $\text{supp}(f_1)$ of a previous solution f_1 as a subset. When $\text{supp}(f_1) \subsetneq \text{supp}(f_2)$ holds for a pair of solutions, the following statement must evaluate to TRUE for the solution f_1 :

$$\begin{aligned}
 & \neg \bigwedge_{e \in \text{supp}(f_2)} z_e = \text{TRUE} \\
 \text{or,} & \quad \bigvee_{e \in \text{supp}(f_2)} \neg z_e = \text{TRUE} \\
 \text{or,} & \quad \sum_{e \in \text{supp}(f_2)} (1 - z_e) \geq 1 \\
 \text{or,} & \quad |\text{supp}(f_2)| - \sum_{e \in \text{supp}(f_2)} z_e \geq 1 \\
 \text{or,} & \quad \sum_{e \in \text{supp}(f_2)} z_e \leq |\text{supp}(f_2)| - 1
 \end{aligned}$$


In more detail: Let f_i be the i th solution found. We then disallow subsequent solutions to contain $\text{supp}(f_i)$ by requiring that at least one of the hyperedges of f_i is not used. This is expressed via the following constraint:

$$\sum_{e \in \text{supp}(f_i)} z_e \leq |\text{supp}(f_i)| - 1 \tag{25}$$


This constraint ensure that not all of the hyperedges in solution f_i is used in some subsequent solution. When the k th solution is to be found, there has thus been added $k - 1$ of these constraints in total. However, this method works



(a) An integer hyperflow solution with 4 hyperedges.



(b) Another integer hyperflow solution to the same flow query using 5 hyperedges.



(c) An integer hyperflow satisfying all constraints listed in Section 5.3. However, it is just the superposition of the solutions 37a and 37b.

Figure 37: Addition of constrain (25) removes solutions like those shown in 37c from the enumerated solutions.

only when the set of variables enumerated over is boolean, as is the case with z_e .

In Figure 36, the hyperflow of the entire figure (with the part with lower opacity) has a subset of hyperedges which form a solution by itself. Therefore, $\bigwedge z_e$ evaluates to TRUE when evaluated over the set of hyperedges in the solution depicted in deep blue. Also, the left-hand side of the constraint 25 is $\sum z_e = 4$ when evaluated over all the hyperedges in the hyperflow solution in dark blue which is $\not\leq 4 - 1 = 3$. As a result, the hyperflow is not enumerated as a subsequent solution after the one in deep blue is added to the solution set. Similarly the solution in Figure 37c, would be excluded after either of the solutions in Figure 37a or 37b has been added to the solution set. The left-hand side of the constraint 25 is $\sum z_e = 4$ when evaluated over all the hyperedges in the hyperflow in Figure 37a which is $\not\leq 4 - 1 = 3$. Therefore, the hyperflow in Figure 37c is not enumerated as a subsequent solution after the one in Figure 37a is added to the solution set. However, addition if the solution in Figure 37a does not exclude the solution in 37b as the left-side of constraint 25 evaluates to $\sum z_e = 3$ (when summed over all hyperedges in the solution in Figure 37a) which is $\leq 4 - 1 = 3$.

5.3.2 Partial Ordering of Molecules

The ILP formulation of the problem [5] accounts for the conservation of the for each molecule, but not for temporal ordering of the reactions in the returned flow. In scenarios such as synthesis planning [6, 7], we would like to interpret a pathway as a directed acyclic process that generates the target molecule(s) unlike the one depicted in Figure 38. We therefore introduce constraints to enforce a partial temporal order on the found pathways.

The ordering is implemented by introducing a supplementary integer variable t_v for each vertex v that keeps track of their ordering in the pathway. The condition $t_u < t_v$ implies




Figure 38: An example of a solution which satisfies the flow conservation constraint (2) but cannot be realized if only v_0 is the allowed source and v_3 the requested target. To effect the first hyperedge, vertex v_1 is required to be present, which is in turn produced by the second hyperedge. Addition of constrain (26) remove such solution from being enumerated which are not physically realizable as a sequence of reaction to the target molecule(s).

that the vertex u is visited by the flow before the vertex v . This is ensured by the following implication constraint:

$$\begin{aligned} \forall e \equiv (e^-, e^+) \in E: \\ f_e \geq 1 \implies t_v \geq t_u + 1, \\ \forall \text{ pairs } (u, v) \text{ such that } u \in e^-, v \in e^+ \end{aligned} \quad (26)$$

That is, the constraint ensures that reactants are assigned a lower order than products of a reaction with flow. We acknowledge that it can be too restrictive to enforce constraints to return pathways with an assigned partial ordering of vertices because they would eliminate all pathways with cycles in them. Cycles are present whenever catalysts are used and in numerous metabolic networks. We merely suggest the possibility of adding such constraints as part of the larger MILP model. There might be other use cases where these partial constraints should be disabled in the model. In these cases, the realizability question for pathways can be addressed using a separate, more robust tool using the computationally more involved formalism of Petri nets on top of the MILP solver. We refer the reader to a detailed study [8] for a more thorough treatment of this issue of realizability of pathways.




Figure 39: Addition of constrain (27) ensures that the reactants are always available to effect a reaction either as products from a previous reaction or as sources. This prevents ‘back edges’ in the pathway where a molecule is used that is produced later along the pathway.

The implication constraint in Equation (26) assigns the temporal ordering t_v to each vertex v . It is implemented as a linear constraint as

$$\forall \text{ pairs } (u, v) \mid \exists e \equiv (e^-, e^+) \text{ with } u \in e^-, v \in e^+ : \\ t_v - t_u - |V|(z_e - 1) - 1 \geq 0 \quad (27)$$

$$\forall v \in V,$$

$$0 \leq t_v \leq |V| \quad (28)$$

The constraint in Equation (27) enforces the assignment $t_v > t_u$ only if the corresponding hyperedge is used in the hyperflow. The size of the set of vertices in the hypergraph $|V|$ is used instead of the large integer M in the constraint in Equation (27) and as an upper bound for the temporal ordering variable t_v for each vertex v in the constraint in Equation (28). The number of constraints added to the MILP model to determine the topological sorting of the vertices is $\mathcal{O}(|V|^2)$.

5.4 COMPARISON WITH OTHER METHODS

Previous work such as [9, 10] have utilized hyperpaths in directed hypergraphs to depict pathways in reaction networks. However, they use the notion of path length to characterize the goodness of the pathway: a shorter pathway is considered better. Often this might not be the case as the shortest pathway might contain a slow or a high-cost reaction, which becomes a bottleneck for the entire pathway. Therefore, it is important to consider some physical attributes of the reactions (such as the barrier heights) in addition to the number of reactions while searching for pathways.

Most of previous work [11–15] related to searching for pathways in reaction networks represent the underlying network as a graph with simple edges. However, this does not capture the nature of chemical reactions faithfully as it might involve multiple reactant molecules and produce multiple product molecules. Often the modeling tracks the most important molecules in the reactions [16, 17], relegating the other molecules involved in the reaction as reagents or side-products (in the style of synthesis maps of natural products). However, since the mathematical framework of hypergraphs exists to represent many-to-many mappings, we choose to use it instead of graphs. This has only one drawback: the visualization of the reaction networks becomes a little more involved and non-intuitive.

Studies focusing on large reaction networks such as [9, 10, 15, 18] often extract the network from databases or existing knowledge. In contrast, using a generative methodology which can create the network on the fly (from a set of input molecules and a set of rules representing classes of reactions) allows us to investigate regions of the molecular space what have not been studied before. Moreover, it removes biases from previous experiments and studies that might have crept in while curating the database. If a particular promising molecule is not entered into the database, then no pathways queried from that network will contain it.

While there exists prior work which includes the generation of reaction networks, these networks were mostly generated using the gradient of potential information [17, 19, 20]. The determination of the gradient information used in potential energy surface explorations is computationally expensive and might limit the size of the studied network. Such studies focus on expanding the network only in the most promising directions and do not consider the remaining molecular space. Such a choice necessitated by computational costs hinders a thorough examination of the molecular space for pathways. Chemical heuristics used by [14, 16] offers a way out of this problem, and our method uses reaction templates to generate the networks this other studies in this class. We aim to generate an exhaustive network that is an over-approximation of what actually occurs in the system. The use of reaction templates allows us to cover all reaction possibilities in the generated network with a manageable computational cost. The current method uses graph rewriting and double pushouts as the formal framework behind the expansion of the network, which to the best of our knowledge is not used in other similar works.

REFERENCES

- [1] IUPAC. *International Union of Pure and Applied Chemistry: Arrhenius Equation*. 2025. DOI: [10.1351/goldbook.A00446](https://doi.org/10.1351/goldbook.A00446). URL: <https://doi.org/10.1351/goldbook.A00446>.
- [2] IUPAC. *International Union of Pure and Applied Chemistry: Modified Arrhenius Equation*. 2025. DOI: [10.1351/goldbook.M03963](https://doi.org/10.1351/goldbook.M03963). URL: <https://doi.org/10.1351/goldbook.M03963>.
- [3] IUPAC. *International Union of Pure and Applied Chemistry: Transition State Theory*. 2025. DOI: [10.1351/goldbook.T06470](https://doi.org/10.1351/goldbook.T06470). URL: <https://doi.org/10.1351/goldbook.T06470>.

- [4] Jakob L. Andersen. *MØD*. <http://mod.imada.sdu.dk>. 2024.
- [5] Jakob L. Andersen et al. “Chemical Transformation Motifs—Modelling Pathways as Integer Hyperflows”. In: *IEEE/ACM Trans. Comput. Biol. Bioinformatics* 16.2 (Apr. 2019), pp. 510–523. ISSN: 1545-5963. DOI: [10.1109/TCBB.2017.2781724](https://doi.org/10.1109/TCBB.2017.2781724). URL: <https://doi.org/10.1109/TCBB.2017.2781724>.
- [6] John Bradshaw et al. *Barking up the right tree: an approach to search over molecule synthesis DAGs*. 2020. arXiv: [2012.11522](https://arxiv.org/abs/2012.11522) [cs.LG]. URL: <https://arxiv.org/abs/2012.11522>.
- [7] Rolf Fagerberg et al. “Finding the K best synthesis plans.” In: *J Cheminform* 10.19 (Apr. 2018). DOI: [10.1186/s13321-018-0273-z](https://doi.org/10.1186/s13321-018-0273-z). URL: <https://doi.org/10.1186/s13321-018-0273-z>.
- [8] Jakob L. Andersen et al. *On the Realisability of Chemical Pathways*. 2023. arXiv: [2309.10629](https://arxiv.org/abs/2309.10629) [q-bio.MN].
- [9] Spencer Krieger and John Kecicioglu. “Shortest Hyperpaths in Directed Hypergraphs for Reaction Pathway Inference”. In: *Journal of Computational Biology* 30.11 (2023). PMID: 37906100, pp. 1198–1225. DOI: [10.1089/cmb.2023.0242](https://doi.org/10.1089/cmb.2023.0242). eprint: <https://doi.org/10.1089/cmb.2023.0242>. URL: <https://doi.org/10.1089/cmb.2023.0242>.
- [10] M. Nakamura, T. Hachiya, Y. Saito, et al. “An efficient algorithm for de novo predictions of biochemical pathways between chemical compounds”. In: *Bioinformatics* 13 (Suppl 17).S8 (Dec. 2012). DOI: [10.1186/1471-2105-13-S17-S8](https://doi.org/10.1186/1471-2105-13-S17-S8).
- [11] Yeonjoon Kim et al. “Efficient prediction of reaction paths through molecular graph and reaction network analysis”. In: *Chem. Sci.* 9 (4 2018), pp. 825–835. DOI: [10.1039/C7SC03628K](https://doi.org/10.1039/C7SC03628K). URL: <http://dx.doi.org/10.1039/C7SC03628K>.


- [12] Ali Hashemi et al. "ReNeGate: A Reaction Network Graph-Theoretical Tool for Automated Mechanistic Studies in Computational Homogeneous Catalysis". In: *Journal of Chemical Theory and Computation* 18.12 (2022). PMID: 36321652, pp. 7470–7482. DOI: [10.1021/acs.jctc.2c00404](https://doi.org/10.1021/acs.jctc.2c00404). URL: <https://doi.org/10.1021/acs.jctc.2c00404>.
- [13] Tomonori Ida, Honoka Kojima, and Yuta Hori. "Predicting and analyzing organic reaction pathways by combining machine learning and reaction network approaches". In: *Chem. Commun.* 59 (83 2023), pp. 12439–12442. DOI: [10.1039/D3CC03890D](https://doi.org/10.1039/D3CC03890D). URL: <http://dx.doi.org/10.1039/D3CC03890D>.
- [14] Dmitrij Rappoport et al. "Complex Chemical Reaction Networks from Heuristics-Aided Quantum Chemistry". In: *Journal of Chemical Theory and Computation* 10.3 (2014). PMID: 26580168, pp. 897–907. DOI: [10.1021/ct401004r](https://doi.org/10.1021/ct401004r). URL: <https://doi.org/10.1021/ct401004r>.
- [15] Agnieszka Wołos et al. "Synthetic connectivity, emergence, and self-regeneration in the network of prebiotic chemistry". In: *Science* 369.6511 (2020), eaaw1955. DOI: [10.1126/science.aaw1955](https://doi.org/10.1126/science.aaw1955). URL: <https://www.science.org/doi/abs/10.1126/science.aaw1955>.
- [16] Michael Woulfe and Brett M. Savoie. "Chemical Reaction Networks from Scratch with Reaction Prediction and Kinetics-Guided Exploration". In: *Journal of Chemical Theory and Computation* 21.3 (2025). PMID: 39883589, pp. 1276–1291. DOI: [10.1021/acs.jctc.4c01401](https://doi.org/10.1021/acs.jctc.4c01401). URL: <https://doi.org/10.1021/acs.jctc.4c01401>.
- [17] Q. Zhao and B.M. Savoie. "Simultaneously improving reaction coverage and computational cost in automated reaction prediction tasks." In: *Nat Comput Sci* 1 (July 2021), pp. 479–490. DOI: [10.1038/s43588-021-00101-3](https://doi.org/10.1038/s43588-021-00101-3). URL: <https://doi.org/10.1038/s43588-021-00101-3>.

- [18] Amanda L. Dewyer, Alonso J. Argüelles, and Paul M. Zimmerman. “Methods for exploring reaction space in molecular systems”. In: *WIREs Computational Molecular Science* 8.2 (2018), e1354. DOI: [10.1002/wcms.1354](https://doi.org/10.1002/wcms.1354). URL: <https://wires.onlinelibrary.wiley.com/doi/abs/10.1002/wcms.1354>.
- [19] Alberton, D. et al. “Accurate ab initio spectroscopic studies of promising interstellar ethanolamine iminic precursors”. In: *A&A* 683 (2024), A198. DOI: [10.1051/0004-6361/202348196](https://doi.org/10.1051/0004-6361/202348196). URL: <https://doi.org/10.1051/0004-6361/202348196>.
- [20] Yutai Zhang, Chao Xu, and Zhenggang Lan. “Automated Exploration of Reaction Networks and Mechanisms Based on Metadynamics Nanoreactor Simulations”. In: *Journal of Chemical Theory and Computation* 19.23 (2023). PMID: 38031422, pp. 8718–8731. DOI: [10.1021/acs.jctc.3c00752](https://doi.org/10.1021/acs.jctc.3c00752). URL: <https://doi.org/10.1021/acs.jctc.3c00752>.


ANOTHER CASE STUDY

This chapter is identical to Section 4 of the published version of the preprint at [arXiv:2504.10609](https://arxiv.org/abs/2504.10609).


In this chapter, we apply the methods from Chapter 5 to a reaction network generated in silico from glycolonitrile (or more formally 2-hydroxyethanenitrile), ammonia (NH_3), and water (H_2O), a network previously studied in [1]. The choice of 2-hydroxyethanenitrile as the starting molecule was driven by the fact that it has been reported to occur in interstellar medium as a product of the condensation of methanal with hydrogen cyanide and act as a precursor to important molecules in pre-biotic systems [2]. The previous work [1] constructed a free-energy map of the molecules that might be present when 2-hydroxyethanenitrile reacts with water and ammonia using computationally-intensive density functional theory calculations. It focused on finding thermodynamic sinks and kinetic barriers in the different regions of the free energy landscape. In this chapter, the network was expanded using the graph transformation based generative framework MØD [3]. The generation of the network, starting from the three above-mentioned molecules was done through the recursive application of the three graph transformation rules shown in Figure 40. A graph transformation rules (to repeat what was already stated in Chapter 4) is a template representing a class of reactions, which share the same set of atoms involved in the reaction, with similar changes in the bond set. The left hand side of the rules is a subgraph (representing a molecular



(a) Addition to a cyanide: in this rule $x \in \{N, O\}$, representing the addition of an amide nitrogen or hydroxyl oxygen to the nitrile bond respectively.



(b) Addition to an imine or carbonyl: in this rule $x, y \in \{N, O\}$, representing the addition of an amide nitrogen or hydroxyl oxygen to an amide or hydroxyl bond respectively. This single rule encapsulates $2^2 = 4$ individual rules depending on the instantiation of x and y .



(c) 1,3 hydrogen shift: in this rule $x, y \in \{C, N, O\}$, representing the shift of a double bond as in tautomerism. When $x \equiv O$ and $y \equiv O$ it is keto-enol tautomerism, $x \equiv N$ and $y \equiv N$ is imine-enamine tautomerism, $x \equiv N$ and $y \equiv O$ is amine-iminol tautomerism and $x \equiv C$ and $y \equiv C$ is an allylic shift of the double bond. This single rule encapsulates $3 \times 3 = 9$ individual rules depending on the instantiation of x and y .

Figure 40: The graph transformation rules used to expand the molecular space and generate the network. The inverse of all the three graph transformation rules were also used while generating the reaction network.

fragment), which can be rewritten as the subgraph on the right hand side, when a match is found in one of the graphs representing molecules already present in the reaction network. This executes a reaction, adding the new product molecules and the reaction the reaction network, and expands the molecular space.






Figure 41: An abstract representation of the generated reaction network with 44 vertices (blue circles) and 116 hyperedges (green squares), on which the working of the method proposed in Chapter 5 was demonstrated.

Molecules with up to two carbon, four nitrogen, and four oxygen atoms were included in the generated network for comparability with [1], which considered molecules with two carbon atoms only. The constructed reaction network had 44 vertices and 116 hyperedges, and shown in Figure 41. A list of molecules and reactions in the generated network is available in the GitHub data repository. To find probable pathways to biologically important molecules of glycine and glycolic acid, the structures of which are shown in Figure 42a and 42b respectively, using glycolonitrile, NH_3 , and H_2O , an ILP was formulated to query the desired pathways. The formulated ILP had 336 variables and 1561 constraints. The reason for choosing glycine is that it is the simplest amino acid and can be considered as the building block of proteins which are essential for life. Glycolic acid



(a) glycine



(b) glycolic acid

Figure 42: target molecules in the flow query.

is also widespread in nature in the form of glycolate salts which are produced during photorespiration in plants.

The five best scoring pathways to the target glycine is depicted in Figure 43. The structures for the molecules in the corresponding vertices is listed in Table 44 and the weights assigned to the hyperedges listed in Table 14. The **best scoring pathway** scores 143118 on the chosen objective (24) and proceeds through 2-iminoethanal (v_6), a pre-biotic intermediate hypothesized to be of significance. A water molecule acts as a catalyst in this pathway since it is used in reaction e_1 and generated back in reaction e_4 . The **second best pathway** scores 143190 on the objective (24). The last two reactions are the same for both (in fact, for all *five*) pathways. This second best pathway uses an ammonia molecule as a catalyst, molecules with a higher chemical potential (under unit concentrations) and reactions with higher energy barriers. The sum of the energy barriers for the reactions in this pathway is $200.02 \text{ kJ mol}^{-1}$ as opposed to $128.74 \text{ kJ mol}^{-1}$ for the best pathway. One might note that the difference in the objective value of the best and the second-best solutions, that is $143190 - 143118 = 72$ is entirely due to the difference in the sum of the energy barriers of the reactions along the pathway ($(200 - 128) \text{ kJ mol}^{-1} = 72 \text{ kJ mol}^{-1}$) or the first part of the objective (24) because both the pathways have the same length. Both pathways consist of 6 reactions and the second part of the objective (24) is the same for both solutions.

The **third pathway** branches off the second pathway and merges into the first, with a different reaction e_4 eliminating an ammonia molecule. This ammonia molecules was added to glycolonitrile in the first step and therefore, acts as a catalyst in the pathway. The **fourth pathway** uses a

different molecule instead of iminoacetaldehyde, which has a higher chemical potential, hence is ranked lower than the previous three. The [fifth pathway](#) has *eight* reactions, while the previous four had seven reactions. Despite having a lower sum of the energy barriers ($152.90 \text{ kJ mol}^{-1}$), it is ranked lower than the previous three pathways because it is longer. This exemplifies how the length of the pathways affects its ranking through the second part of the objective ([24](#)). Figure [43](#) shows how interconnected the individual different pathways in the network might be, often with parallel branches and cross-connections, which leads to complex behavior of the system.

The estimated energy barriers for the reactions used in the reported pathways in Figure [43](#) is presented in Table [14](#). Table [14](#) consists of five columns that list the assigned weights for the hyperedges in the pathways shown in Figure [43](#). The shaded cells indicate hyperedges that have already been used in a previously enumerated pathway. For instance, all five pathways in Figure [43](#) share the final two reactions. Consequently, the next four pathways reuse the last two reactions of the first pathway, and the last two rows for the corresponding columns are shaded to reflect this overlap.

Ab initio molecular dynamics has been used in a previous work [[4](#), [5](#)] to simulate the Urey-Miller experiment and find possible pathways to glycine in a prebiotic setting. The reported pathways in [[4](#)] also contain reactions catalysed by water and ammonia. The reported pathways to glycine in [[4](#)] proceeds through a singlet carbene intermediate, while those in [[5](#)] has a molecule with a three-membered ring, oxiranimine, as an intermediate. However, in the current work, there were no reaction templates used which might lead to the formation of a carbene-like molecule or a three-cycle in a molecule in the generated network. Hence, such a pathway was not observed. Carbenes and three-cycles are highly reactive intermediates, and their formation is a high energy process. On the whole, the pathways presented in this work have much lower reported energy barriers than that reported in these two studies.




Figure 43: Top: A schematic depiction of the five best pathways to glycine according to the chosen scoring function (energy levels not shown exactly to scale). Bottom: The energy profile of the best-scoring pathway (in blue) from above (energy levels shown to scale) with the molecular structures explicitly shown.

Name	Molecular Structure	Name	Molecular Structure
v_0	<chem>HO-C#N</chem>	v_1	<chem>NH3-CH2-CHO</chem>
v_2	<chem>H2O-CH2-CHO</chem>	v_{13}	<chem>HO-CH2-CHO</chem>
v_3	<chem>HO-CH2-CHO</chem>		<chem>NH2-CH2-CHO</chem>
v_4	<chem>HO-CH(OH)-CHO</chem>	v_{14}	<chem>HO-CH(OH)-CHO</chem>
v_5	<chem>O=CH-CH(OH)-NH2</chem>		<chem>NH2-CH(OH)-CHO</chem>
v_6	<chem>O=CH-CH(OH)-NH2</chem>	v_{15}	<chem>O=CH-CH(OH)-NH2</chem>
v_7	<chem>HO-CH(OH)-CHO</chem>		<chem>NH2-CH(OH)-CHO</chem>
v_8	<chem>HO-CH(OH)-CHO</chem>	v_{16}	<chem>HO-CH(OH)-CHO</chem>
v_9	<chem>HO-CH(OH)-CHO</chem>		<chem>NH2-CH(OH)-CHO</chem>
		v_{36}	<chem>H2N-CH(OH)-CHO</chem>
		v_{52}	<chem>H2N-CH(OH)-CHO</chem>

Figure 44: List of the molecular structures corresponding to the the labeled vertices in the pathways in Figure 43.

Reaction count	Energy barriers (in kJ mol^{-1})				
	pathway 1	pathway 2	pathway 3	pathway 4	pathway 5
e_1	13.95 ($v_0 + v_2, v_3$)	33.18 ($v_0 + v_1, v_{13}$)	33.18 ($v_0 + v_1, v_{13}$)	13.95 ($v_0 + v_2, v_3$)	13.95 ($v_0 + v_2, v_3$)
e_2	7.67 (v_3, v_4)	64.17 (v_{13}, v_{14})	64.17 (v_{13}, v_{14})	7.67 (v_3, v_4)	18.67, 13.15 (v_3, v_{52}, v_4)
e_3	57.69 (v_4, v_5)	28.40 (v_{14}, v_{15})	28.40 (v_{14}, v_{15})	57.69 (v_4, v_5)	57.69 (v_4, v_5)
e_4	1.67 ($v_5, v_6 + v_2$)	19.86 ($v_{15} + v_2, v_{16}$)	29.35 ($v_{15}, v_6 + v_1$)	67.36 ($v_5 + v_2, v_{36}$)	1.67 ($v_5, v_6 + v_2$)
e_5	17.14 ($v_6 + v_2, v_7$)	23.79 ($v_{16}, v_7 + v_1$)	17.14 ($v_6 + v_2, v_7$)	60.84 ($v_{36}, v_7 + v_2$)	17.14 ($v_6 + v_2, v_7$)
e_6	0.19 (v_7, v_8)	0.19 (v_7, v_8)	0.19 (v_7, v_8)	0.19 (v_7, v_8)	0.19 (v_7, v_8)
e_7	30.42 (v_8, v_9)	30.42 (v_8, v_9)	30.42 (v_8, v_9)	30.42 (v_8, v_9)	30.42 (v_8, v_9)
sum	128.74	200.02	202.85	238.13	152.90

Table 14: From the left, the columns list the weights assigned to the hyperedges along with the sum of the weights, for the five pathways in Figure 43.

Reaction count	Energy barrier					
	pathway 1	pathway 2	pathway 3	pathway 4	pathway 5	pathway 6
e_1	13.95 ($v_0 + v_2, v_3$)	13.95 ($v_0 + v_2, v_3$)	13.95 ($v_0 + v_2, v_3$)	33.18 ($v_0 + v_1, v_{33}$)	13.95 ($v_0 + v_2, v_3$)	33.18 ($v_0 + v_1, v_{33}$)
e_2			7.67 (v_3, v_{24})	48.612 ($v_{33} + v_2, v_{34}$)	80.53 ($v_3 + v_1, v_{34}$)	48.61 ($v_{33} + v_2, v_{34}$)
e_3	26.61 ($v_3 + v_2, v_6$)	18.68 (v_3, v_{15})	11.76 (v_{24}, v_{15})	43.64 ($v_{34}, v_{15} + v_1$)	43.64 ($v_{34}, v_{15} + v_1$)	33.08 ($v_{34}, v_3 + v_1$)
e_4		0.62 ($v_{15} + v_2, v_6$)	0.62 ($v_{15} + v_2, v_6$)	0.62 ($v_{15} + v_2, v_6$)	0.62 ($v_{15} + v_2, v_6$)	0.62 ($v_3 + v_2, v_6$)
e_5	74.62 ($v_6, v_7 + v_1$)	74.62 ($v_6, v_7 + v_1$)	74.62 ($v_6, v_7 + v_1$)	74.62 ($v_6, v_7 + v_1$)	74.62 ($v_6, v_7 + v_1$)	74.62 ($v_6, v_7 + v_1$)
sum	118.19	107.87	108.63	200.68	213.38	219.11

Table 15: From the left, the columns list the weights assigned to the hyperedges along with the sum of the weights, for the six pathways in Figure 45.




Figure 45: The six best pathways to glyoxalic acid according to the chosen scoring function. The best scoring pathway is depicted in blue, and the next best pathways in green, brown, cyan, azure and red. The structures for the molecules for the vertices is listed in Table 46 and the weights assigned to the hyperedges listed in Table 15.

Name	Molecular Structure
v_0	<chem>NC#CCO</chem>
v_3	<chem>NC(=O)CO</chem>
v_6	<chem>NC(O)(O)CO</chem>
v_7	<chem>NC(=O)CO</chem>
v_{15}	<chem>NC(=O)C(=O)O</chem>
v_{24}	<chem>NC(=O)C(=O)O</chem>
v_{33}	<chem>NC(=O)CO</chem>
v_{34}	<chem>NC(O)(O)CO</chem>

Figure 46: List of the molecular structures corresponding to the the labeled vertices in the pathways in Figure 45.

Another molecule of importance in prebiotic systems which appears in the generated network is 2-hydroxyethanoic acid, commonly known as glycolic acid. During photorespiration plants produce glycolate. It can also be used as a monomer for various biodegradable polymers such as polyglycolic acid and poly(lactic-co-glycolic) acid. However, there have not been many studies on pathways to glycolic acid from the precursor molecules in prebiotic systems. Figure 45 shows the six best pathways to 2-hydroxyethanoic

acid along with the molecules used in the pathway in Table 46. The **best scoring pathway** has three reactions, the **second best** has four reaction, while the rest involve five reactions. Despite the third-listed pathway having a lower sum of energy barriers ($108.63 \text{ kJ mol}^{-1}$), it is ranked lower because it is longer, illustrating the effect of the second term in the objective (24).

Table 15 contains six columns that list the assigned weights for the hyperedges in the pathways illustrated in Figure 45. As in Table 14, the shaded cells indicate hyperedges that have already been used in a previously enumerated pathway. The specific hyperedge is mentioned below the energy barrier, with the reactants as the first element of the pair and the products as the second element. Individual molecules are separated by a '+', and the structures of the molecules the labels correspond to can be looked up from Table 46.

The expansion of the network shown in Figure 41 took 67 seconds. It is quite faster than comparable methods using reaction templates without using graph transformation, molecular dynamics or potential energy search as done in [1]. Annotating the 116 hyperedges with the energy barriers of the reactions as weights took around an hour. Using a neural network instead of more accurate traditional methods does lead to trade off of the accuracy for speed, but it allows scalability for larger networks. Formulating the ILP and enumerating the pathways in the network to the two target molecules using the objective function (24) took around 52 seconds. All the reported times are the average of 10 runs on an AMD[®] Ryzen 5 PRO 4650U CPU (2.1 GHz) utilizing 8 threads. One may also expand the network from methanal and hydrogen cyanide instead of 2-hydroxyethanenitrile and use a more general version of the rules where the variable heavy atoms are allowed to match with carbon, nitrogen or oxygen atoms. However, it leads to a larger network with 481 vertices and 2768 hyperedges, of which the considered network is a sub-network.

REFERENCES

- [1] Jeremy Kua and Kyra L. Thrush. "HCN, Formamidic Acid, and Formamide in Aqueous Solution: A Free-Energy Map". In: *The Journal of Physical Chemistry B* 120.33 (2016). PMID: 27016454, pp. 8175–8185. DOI: [10.1021/acs.jpcb.6b01690](https://doi.org/10.1021/acs.jpcb.6b01690). URL: <https://doi.org/10.1021/acs.jpcb.6b01690>.
- [2] G. Danger et al. "Hydroxyacetonitrile (HOCH₂CN) formation in astrophysical conditions. Competition with the aminomethanol, a glycine precursor". In: *The Astrophysical Journal* 756.1 (Aug. 2012), p. 11. DOI: [10.1088/0004-637X/756/1/11](https://dx.doi.org/10.1088/0004-637X/756/1/11). URL: <https://dx.doi.org/10.1088/0004-637X/756/1/11>.
- [3] Jakob L. Andersen. MØD. <http://mod.imada.sdu.dk>. 2024.
- [4] LP. Wang, A. Titov, R. McGibbon, et al. "Discovering chemistry with an ab initio nanoreactor." In: *Nature Chem* 6 (Nov. 2014), pp. 1044–1048. DOI: [10.1038/nchem.2099](https://doi.org/10.1038/nchem.2099). URL: <https://doi.org/10.1038/nchem.2099>.
- [5] Léon Huet et al. "A New Route to the Prebiotic Synthesis of Glycine via Ab Initio-Based Machine Learning Calculations". In: *The Journal of Physical Chemistry Letters* 15.34 (2024). PMID: 39159425, pp. 8697–8705. DOI: [10.1021/acs.jpclett.4c01954](https://doi.org/10.1021/acs.jpclett.4c01954). eprint: <https://doi.org/10.1021/acs.jpclett.4c01954>. URL: <https://doi.org/10.1021/acs.jpclett.4c01954>.

EPILOGUE

This chapter is taken from Section 6 of the published versions of the preprints at [arXiv:2411.15900](#) and [arXiv:2504.10609](#).

Thermodynamics emerged from efforts to analyze the operation of steam engines while maximizing their work. In a nice analogy, the search for pathways may be phrased as viewing reaction networks as chemical engines in which we want to maximize the output of some target molecule(s). Recognizing that thermodynamics, encapsulated by the Gibbs free energies of the reactions, play a pivotal role in determining the probability of a reaction, Chapter 3 focused on developing ways to integrate them into the search for pathways.

Exploring pathways with favorable energy profiles involves challenging navigation of a large and complex energy landscape where the driving thermodynamic forces are influenced by the concentration ratios, inflows of the source molecules, and outflows of the target molecules. Chapter 3 attempted to address part of that challenge. However, there might be several refinements possible to the model proposed in Chapter 3, some of which we list here:

1. The constraint enforcing that only reactions with negative free energies are allowed in the returned pathways (implemented through the constraint in Equation (5)) may be overly stringent. Under experimental conditions, the reactions observed to occur do not always align with those predicted to occur theoretically: the thermal decomposition of lithium carbonate is ob-

served to occur in practice despite being predicted to be thermodynamically unfavorable using computed (or experimental) free energies for the reaction. However, reactions predicted to be thermodynamically favorable do not always occur in practice because they have a high energy barrier (like the allotropic conversion of diamond into graphite). Therefore, there should be some flexibility in delineating the boundary between those reactions that are considered likely and included in the pathway and those that are not, to bring the model closer to reality. Such extensions of the current modeling should be fairly straightforward.

2. The objective function minimizes the sum of the Gibbs free energies of the reactions in the pathway. However, this approach might favor a pathway with more extreme values for the free energies rather than one with more moderate free energy values. Therefore, it might be worthwhile to explore alternative objective functions, such as the min-max, in addition to the min-sum used here.
3. The current formulation disregards the kinetics of individual reactions. However, the probability of a reaction occurring is determined to a substantial extent by the reaction barrier height, and not only by the concentration of the reactants and the products. As a result, our framework might include a reaction in the returned pathway with a favorable thermodynamic driving force but too high a reaction barrier, making it a slow and rate-limiting step for the entire pathway. On the other hand, coming up with a generic ‘kinetic oracle’ with wide applicability to estimate the barrier heights for diverse reactions poses a significant challenge. The transition states determining the barrier heights are saddle points on the potential energy surface, unlike reactants or products that are minima and can be located by gradient descent optimization methods.

Hence, we do not claim that the model proposed in Chapter 3 to be the final in the quest to incorporate thermody-

namics into the search for pathways in reaction networks. Still, we believe that it constitutes a valuable step forward, and we end by summarizing its virtues. We provide a systematic and thermodynamically informed approach to optimize the computational search for pathways in reaction networks.

Modeling pathways as integer hyperflows gives results that can be interpreted in mechanistic terms, as each molecule is present in integral quantities. Formulating the problem as an MILP problem and utilizing an MILP solver allows us to obtain solutions with reasonable computational resources and time. It allows enumeration of pathways over the integral flow variables, facilitating a systematic exploration of pathways in the reaction network. Returning several solutions to pathway queries is an asset, as the answer becomes more robust to possible errors in the calculations of chemical potentials of the molecules and to other artifacts of the simplification of reality inherent in any modeling. Moreover, it offers alternatives to choose from if the optimal solution turns out to be unattainable under controlled industrial reactor or laboratory conditions.

Finally, the application of the framework formulated in Chapter 3 to a reaction network, in Chapter 4, based on real-life chemistry acts as a proof-of-concept and illustrates the computational feasibility and analytic potential of our methods.

In Chapter 5, we have described a methodology for kinetically informed exploration of pathways in reaction networks building on the model proposed in Chapter 3. One part of the methodology is an automated workflow for annotating individual reactions with their energy barriers, computed using a diverse selection of tools—RDKit [1], xTB [2], ASE [3] and NeuralNEB [4]. The goal of this workflow is to minimize the laborious manual interventions and intensive expert knowledge often necessary when executing quantum chemical calculations to determine energy barriers. As demonstrated in Chapter 6, the workflow can be applied on networks of rather large scale, can work on the fly with generative network construction, and uses rela-

tively moderate computational resources. The method does not take into account the interactions of solvent molecules while calculating the energy barriers. This and the question of how well models based on machine learning generalize past their training sets may lead to reasonable reservations on the precision of the estimates of the energy barriers for some use cases. However, the proposed methodology offers the flexibility that this component can be substituted with alternatives, if desired.

As the other part of the proposed methodology, we formulated an ILP to query pathways in the annotated network and designed an objective function for pathway evaluation. The query formalism is at the same time both precise and very flexible in what queries can be formulated. The objective function is made under the simplifying assumption that all molecules in the reaction network are present with unit concentrations at all times. Additionally, the costs assigned to pathways depend on the underlying reaction network considered, so comparisons across networks are not directly possible. However, the question of how to evaluate pathways is complex and the reader of course has the option to substitute a different objective function, based on other cost metrics. We do note, though, that our simplification and the formulation via ILP allows us to analyze much larger networks and using much less computational resources than alternatives like stochastic simulations or potential energy surface exploration. We also note that by a progressive constraining of the ILP, we in the end can return a *list* of high-scoring solutions, each representing structurally different pathways. Giving a list of good, structurally different solutions, rather than a single solution, helps mitigate disagreements about the details of pathway cost metrics.

Finally, the efficiency and utility of the method described in Chapter 5 was shown on a concrete reaction network in Chapter 6, studying pathways to pre-biotically significant molecules, such as glycine and 2-hydroxyethanoic acid.

In summary, this work contributes to field of computationally assisted analysis and design of pathways in reaction networks represented as directed hypergraphs.

REFERENCES

- [1] RDKit documentation. *embedMolecule()* function in RDKit. <https://www.rdkit.org/docs/source/rdkit.Chem.rdDistGeom.html#rdkit.Chem.rdDistGeom.EmbedMolecule>.
- [2] Christoph Bannwarth et al. “Extended tight-binding quantum chemistry methods”. In: *WIREs Computational Molecular Science* 11.2 (2021), e1493. DOI: <https://doi.org/10.1002/wcms.1493>. URL: <https://wires.onlinelibrary.wiley.com/doi/abs/10.1002/wcms.1493>.
- [3] Ask Hjorth Larsen et al. “The atomic simulation environment - a Python library for working with atoms”. In: *Journal of Physics: Condensed Matter* 29.27 (2017), p. 273002. URL: <http://stacks.iop.org/0953-8984/29/i=27/a=273002>.
- [4] Mathias Schreiner et al. “NeuralNEB—neural networks can find reaction paths fast”. In: *Machine Learning: Science and Technology* 3.4 (Dec. 2022), p. 045022. DOI: [10.1088/2632-2153/aca23e](https://doi.org/10.1088/2632-2153/aca23e). URL: <https://dx.doi.org/10.1088/2632-2153/aca23e>.

