

Summer School 2025  
Astronomy & Astrophysics



# Project Report

Prepared By

Student Name: ADITYA A

Institution Name: CENTRAL UNIVERSITY OF KARNATAKA

Institution Roll No: 24UPHYS002

ISA Admission No: 176428

Projects Name

JWST\_miri\_spectral\_lines

Submitted To

Name: Mr. Sahil Sakkarwal

Designation: Program Supervisor

Institution: India Space Academy

# JWST\_MIRI\_sp ectra\_analysis

June 19, 2025

## 1 Session 5

In today's session we will learn about how to extract spectra from the MIRI IFU spectral cube, JWST. We will also identify emission lines.

This session will become the base of one of your projects. So take it seriously!!

### 1.1 Spectral Cube

Unlike image that has 2 spatial axis, this has one spectral axis too. It can be in wavelength, frequency or velocity.

Select a collection... and enter target:

MAST Observations by Object Name or RA/Dec

Upload Target List

My Download Basket: 0 files

Portal User Guide | Leave Feedback | About This Site

anonymous

Login...

Account Info...

Home Page

MAST: Advanced Search 1

24 Total Rows

Filters

Clear Filters Edit Filters... Help...

Keyword/Text Filter

Name Quantity

CH2-SHORT (2 of 2)

CH4-MEDIUM (2 of 2)

CH3-LONG (2 of 2)

CH3-SHORT (2 of 2)

CH1-SHORT (2 of 2)

Show 7 More

Target Classification

Name Quantity

Ultraluminous infrared galaxies (24 of 24)

Interacting galaxies (24 of 24)

Infrared galaxies (24 of 24)

Galaxy (24 of 24)

Data URL

Name Quantity

MAST:JWST/product/jw01328-c1006\_t014\_miri\_ch2-short\_s3d.fits (1 of 1)

MAST:JWST/product/jw01328-c1006\_t014\_miri\_ch4-medium\_s3d.fits (1 of 1)

MAST:JWST/product/jw01328-c1006\_t014\_miri\_ch3-long\_s3d.fits (1 of 1)

| band                        | Target Name                     | Target Classification | Observation ID | RA | Dec |
|-----------------------------|---------------------------------|-----------------------|----------------|----|-----|
| Galaxy, Infrared galaxie... | jw01328-c015_t014_miri_ch1-4... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c015_t014_miri_ch1-s... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c015_t014_miri_ch2-4... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c015_t014_miri_ch2-...  | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch3-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch2-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch1-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch1-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch2-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch4-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch1-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch3-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch4-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch2-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch1-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch3-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch4-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch2-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch1-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch3-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch4-... | 23:03:15.610          | +08:52:        |    |     |
| Galaxy, Infrared galaxie... | jw01328-c1006_t014_miri_ch2-... | 23:03:15.610          | +08:52:        |    |     |

AstroView

23:03:31.296 +09:08:57.82

23:03:15.624 +08:52:26.04

RA DEC

JWST has the following instruments:

1. NIRCam
2. NIRSpec
3. MIRI → We will use this today!





## 2 regions

regions is an in-development coordinated package of Astropy for region handling. The Regions package provides classes to represent: Regions defined using pixel coordinates (e.g., CirclePixelRegion) Regions defined using celestial coordinates, but still in an Euclidean geometry (e.g., CircleSky To transform between sky and pixel regions, a world coordinate system object (e.g., astropy.wcs) Regions also provides a unified interface for reading, writing, parsing, and serializing regions data in different formats, including the Legacy Region Format, CRTF (CASA Region Text Format), and FITS Region Binary Table format.

You can read more about it in here:

<https://astropy-regions.readthedocs.io/en/stable/shapes.html>

[13]: *#Region-wise line plotting*

```
# Import necessary libraries

import numpy as np
import pandas as pd
import warnings
import matplotlib.pyplot as plt
from astropy.io import fits
from astropy.wcs import WCS
from regions import Regions

warnings.filterwarnings("ignore", category=UserWarning, append=True)
```

[14]: *# Define the redshift for NGC 7469*

```
# NGC 7469 is a Seyfert galaxy with a redshift of approximately 0.016268
# This value is used to convert observed wavelengths to rest-frame wavelengths
# The redshift value can be obtained from various astronomical databases like
↳ NED (NASA/IPAC Extragalactic Database)

z = 0.016268

reg_path = "region_file.reg" # From DS9 lecture, this is the region file
↳ containing the regions of interest
# Read the regions from the DS9 region file
# Make sure to have the regions file in the same directory as this script
regions = Regions.read(reg_path, format='ds9')
region_labels = ["Center Region", "Ring Region"] # Label for each region
file_paths = []

# Only channel 2 for the session
# for part in ['short', 'medium', 'long']:
```

```

# I have kept them in a subfolder MAST_NGC_7469 for clarity
# You can change the path as per your directory structure
# The file names are based on the JWST data release naming convention
# file_paths.append(f'./MAST_NGC_7469/
jw01328-c1006_t014_miri_ch{2}-{part}_s3d.fits')

for ch_num in range(1,5):
    for part in ['short','medium','long']:
        file_paths.append(f'./MAST_NGC_7469/
jw01328-c1006_t014_miri_ch{ch_num}-{part}_s3d.fits')

print(file_paths)

```

```

['./MAST_NGC_7469/jw01328-c1006_t014_miri_ch1-short_s3d.fits',
'./MAST_NGC_7469/jw01328-c1006_t014_miri_ch1-medium_s3d.fits',
'./MAST_NGC_7469/jw01328-c1006_t014_miri_ch1-long_s3d.fits',
'./MAST_NGC_7469/jw01328-c1006_t014_miri_ch2-short_s3d.fits',
'./MAST_NGC_7469/jw01328-c1006_t014_miri_ch2-medium_s3d.fits',
'./MAST_NGC_7469/jw01328-c1006_t014_miri_ch2-long_s3d.fits',
'./MAST_NGC_7469/jw01328-c1006_t014_miri_ch3-short_s3d.fits',
'./MAST_NGC_7469/jw01328-c1006_t014_miri_ch3-medium_s3d.fits',
'./MAST_NGC_7469/jw01328-c1006_t014_miri_ch3-long_s3d.fits',
'./MAST_NGC_7469/jw01328-c1006_t014_miri_ch4-short_s3d.fits',
'./MAST_NGC_7469/jw01328-c1006_t014_miri_ch4-medium_s3d.fits',
'./MAST_NGC_7469/jw01328-c1006_t014_miri_ch4-long_s3d.fits']

```

Now we have defined the redshift from NED, made a region file from DS9 and made a list of all the files to iterate over.

We can proceed further and extract the spectra from each of the file and plot the final spectrum from it.

```
[16]: from astropy.wcs import FITSFixedWarning
```

```

# Suppress FITSFixedWarning
warnings.simplefilter('ignore', category=FITSFixedWarning)
# Loop through each region
for idx, region in enumerate(regions):
    label = region_labels[idx] if idx < len(region_labels) else f"Region_{
[idx+1]}"

    wavelength_all = []
    spectrum_all = []
    spectrum_all_err = []

    for file_path in file_paths:

```

```

try:
    hdul = fits.open(file_path) data = hdul[1].data
    data[data < 0] = np.nan data_err = hdul[2].data
    header = hdul[1].header wcs = WCS(header)
    mask = region.to_pixel(wcs.celestial).to_mask()
    spectrum = [] spectrum_err = [] for i in
range(data.shape[0]):

        masked_data = mask.multiply(data[i, :, :])
        masked_err = mask.multiply(data_err[i, :, :])
        avg = np.nanmean(masked_data)
        err = np.sqrt(np.nanmean(masked_err**2))
        spectrum.append(avg if not np.isnan(avg) else 0)
        spectrum_err.append(err if not np.isnan(err) else 0)

    crval = header['CRVAL3'] cdelt = header['CDELT3'] crpix =
header['CRPIX3'] wl = (np.arange(data.shape[0]) - (crpix - 1)) *
cdelt + crval

    wl = wl / (1 + z)

    wavelength_all.extend(wl)
    spectrum_all.extend(spectrum)
    spectrum_all_err.extend(spectrum_err)
    hdul.close()

except Exception as e:
    print(f"Error with {file_path}: {e}")

# Sort
wavelength_all = np.array(wavelength_all)
spectrum_all = np.array(spectrum_all)
spectrum_all_err = np.array(spectrum_all_err)
sorted_idx = np.argsort(wavelength_all)
wavelength_all = wavelength_all[sorted_idx]
spectrum_all = spectrum_all[sorted_idx]
spectrum_all_err = spectrum_all_err[sorted_idx]

```

```

#saving the individual spectral plots as a csv file

# Create DataFrame
df = pd.DataFrame({
    'Wavelength (microns)': wavelength_all,
    'Flux': spectrum_all,
    'Flux Error': spectrum_all_err
})


# Generate filename based on region name
fname = label.replace(" ", "_").lower() # e.g., 'center_region'

# Save as CSV
df.to_csv(f"{fname}_spectrum.csv", index=False)

# Plot
plt.figure(figsize=(12, 5))
plt.plot(wavelength_all, spectrum_all, label=label, color='C'+str(idx))
plt.fill_between(wavelength_all,
                 spectrum_all - spectrum_all_err,
                 spectrum_all + spectrum_all_err,
                 color='C'+str(idx), alpha=0.3, label='Error')
plt.xlabel('Wavelength (microns)')
plt.ylabel('Flux')
plt.title(f'JWST MIRI Spectrum - {label}')
plt.grid(True)
plt.legend()
plt.tight_layout()
plt.savefig(f"{label.replace(' ', '_').lower()}_spectrum.png") # Optional:  saveasPNG
plt.show()

# Optional: export to CSV
import pandas as pd
df = pd.DataFrame({
    'Wavelength (microns)': wavelength_all,
    'Flux': spectrum_all,
    'Flux Error': spectrum_all_err
})
df.to_csv(f"{label.replace(' ', '_').lower()}_spectrum.csv", index=False)

```



```
[17]: # Load extracted spectra from CSVs
center_df = pd.read_csv("center_region_spectrum.csv")
ring_df = pd.read_csv("ring_region_spectrum.csv")


# Extract data
wl_center = center_df['Wavelength (microns)']
flux_center = center_df['Flux']
err_center = center_df['Flux Error']
wl_ring = ring_df['Wavelength (microns)']
flux_ring = ring_df['Flux']
err_ring = ring_df['Flux Error']

# Plot
plt.figure(figsize=(14, 6))
plt.plot(wl_center, flux_center, label='Center Region', color='C0')
plt.fill_between(wl_center, flux_center - err_center, flux_center + err_center,
                color='C0', alpha=0.3)


plt.plot(wl_ring, flux_ring, label='Ring Region', color='C1')
plt.fill_between(wl_ring, flux_ring - err_ring, flux_ring + err_ring,
                color='C1', alpha=0.3)

plt.xlabel("Wavelength (microns)")
plt.ylabel("Flux")
plt.title("JWST MIRI Spectra – Center vs Ring Region")
plt.legend()
plt.grid(True)
plt.tight_layout()
plt.savefig("combined_regions_spectrum.png")
```

```
plt.show()
```



```
[18]: %matplotlib widget
plt.figure(figsize=(12, 5))
plt.plot(wavelength_all, spectrum_all, label='Flux', color='darkorange')
plt.fill_between(wavelength_all,
                 spectrum_all - spectrum_all_err,
                 spectrum_all + spectrum_all_err,
                 color='orange', alpha=0.3, label='Error')
plt.xlabel('Wavelength (microns)')
plt.ylabel('Flux')
plt.title('Combined MIRI Spectrum for NGC 7469')
plt.grid(True)
plt.legend()
plt.tight_layout()
plt.show()
```




```
[19]: #plot using ./MAST_NGC_7469/jw01328-c1006_t014_miri_ch2-long_s3d.fits data
plt.figure(figsize=(15, 8))
plt.errorbar(wavelength_all, spectrum_all, yerr=spectrum_all_err, color='blue',
             ecolor='black')
plt.xlabel('Wavelength(microns)')
plt.ylabel('Average Intensity (MJy/sr)')
plt.title("jw01328-c1006_t014_miri_ch2-long_s3d.fits plot")

# Change the x-axis limits to focus on the desired region
# w_start, w_end = 8.95, 9.05
# plt.xlim(w_start, w_end)

# Auto-scale y-axis to the visible range
# plt.ylim(0, 1.1*np.nanmax(spectrum_all[(wavelength_all >= w_start) &
# (wavelength_all <= w_end)]))
plt.grid(True)
plt.legend()
```

[19]: <matplotlib.legend.Legend at 0x2022311b560>



## 2.1 plotting spectra for all 12 files

```
[21]: import numpy as np
import matplotlib.pyplot as plt
from astropy.io import fits
from astropy.wcs import WCS
from regions import Regions
import os

# Set up
z = 0.016268
reg_path = "region_file.reg"
regions = Regions.read(reg_path, format='ds9')
region_labels = ["Center Region", "Ring Region"]

# Define MIRI file paths and labels
file_paths = []
file_labels = []
for ch_num in range(1, 5):
    for part in ['short', 'medium', 'long']:
        path = f'./MAST_NGC_7469/jw01328-c1006-t014-miri_ch{ch_num}-{part}_s3d.
fits' file_paths.append(path)

        file_labels.append(f"ch{ch_num}-{part}")

# Make output folder
os.makedirs("region_channel_plots", exist_ok=True)

# Loop through each region
for region_idx, region in enumerate(regions):
    region_name = region_labels[region_idx]
    safe_name = region_name.replace(" ", "_").lower()

    # Loop through each FITS file
    for file_path, file_label in zip(file_paths, file_labels):
        try:
            hdul = fits.open(file_path)
            data = hdul[1].data
            data[data < 0] = np.nan
            data_err = hdul[2].data
            header = hdul[1].header
            wcs = WCS(header)

            mask = region.to_pixel(wcs.celestial).to_mask()

            spectrum = []
            spectrum_err = []
```

```

for i in range(data.shape[0]):
    masked_data = mask.multiply(data[i, :, :])
    masked_err = mask.multiply(data_err[i, :, :])

    avg = np.nanmean(masked_data)
    err = np.sqrt(np.nanmean(masked_err**2))
    spectrum.append(avg if not np.isnan(avg) else 0)
    spectrum_err.append(err if not np.isnan(err) else 0)

crval = header['CRVAL3']
cdelt = header['CDELT3']
crpix = header['CRPIX3']
wl = (np.arange(data.shape[0]) - (crpix - 1)) * cdelt + crval
wl = wl / (1 + z)

# Sort
wl = np.array(wl)
spectrum = np.array(spectrum)
spectrum_err = np.array(spectrum_err)
sorted_idx = np.argsort(wl)
wl = wl[sorted_idx]
spectrum = spectrum[sorted_idx]
spectrum_err = spectrum_err[sorted_idx]

# Plot
plt.figure(figsize=(10, 4))
plt.plot(wl, spectrum, label='Spectrum', color='C0')
plt.fill_between(wl, spectrum - spectrum_err, spectrum + spectrum_err, color='C0', alpha=0.3, label='Error')
plt.xlabel("Wavelength(Åm)")
plt.ylabel("Flux")
plt.title(f"{region_name} — {file_label}")
plt.legend()
plt.tight_layout()

# Save plot
fname = f"region_channel_plots/{safe_name}_{file_label}.png"
plt.savefig(fname)
plt.close()

except Exception as e:
    print(f"Failed for {file_path} and {region_name}: {e}")

```

[22]: # %pip install plotly

The code below is to plot and interactively view the final spectra. It might be complex so you can treat it as a black box or explore on your own.

Your task here is to fill the spectral features by opening each file in cubeviz. I will give you an example walkthrough.

```
[24]: import plotly.graph_objects as go
import pandas as pd
import numpy as np

# Load spectra from CSV files
center_df = pd.read_csv("center_region_spectrum.csv")
ring_df = pd.read_csv("ring_region_spectrum.csv")
# Organize data for both regions
region_data = {

    "Center Region": center_df,
    "Ring Region": ring_df
}
# Define emission/molecular features
features = {

    'PAHs': {
        'PAH 7.7': 7.7, 'PAH 8.6': 8.6, 'PAH 11.3': 11.3,
    },
    'Iron': {},
    'Neon': {
        '[Ne VI]': 7.65,
    },
    'Other': {
        '[Ar III]': 8.991,
        '[S IV]': 10.51
    },
    'H $\alpha$ ': {
        'S(3)': 9.66,
        'S(4)': 8.03
    }
}
colors = {

    'PAHs': '#FF7F0E',
    'Iron': '#2CA02C',
    'Neon': '#D62728',
    'Other': '#9467BD',
    'H $\alpha$ ': '#8C564B'
}

# Function to plot one region
```

```

def plot_region(region_name, df):
    wavelength = df['Wavelength (microns)'].values
    flux = df['Flux'].values
    flux_err = df['Flux Error'].values

    fig = go.Figure(layout=dict(
        width=800,
        height=500,
        template='plotly_white'
    ))

    # Spectrum trace
    fig.add_trace(go.Scatter(
        x=wavelength,
        y=flux,
        mode='lines',
        name='Spectrum',
        line=dict(color='#1f77b4', width=1.5),
        hovertemplate='λ: %{x:.3f} μm<br>Intensity: %{y:.2f} MJy/sr<extra></
    extra>'
    ))

    # Error band
    fig.add_trace(go.Scatter(
        x=np.concatenate([wavelength, wavelength[::-1]]),
        y=np.concatenate([flux + flux_err, (flux - flux_err)[::-1]]),
        fill='toself',
        fillcolor='rgba(31, 119, 180, 0.2)',
        line=dict(color='rgba(255,255,255,0)'),
        hoverinfo='skip',
        name='Uncertainty'
    ))

    # Emission features
    for category, lines in features.items():
        for name, wl in lines.items():
            fig.add_vline(
                x=wl,
                line=dict(
                    color=colors[category],
                    width=1.5 if category == 'PAHs' else 1,
                    dash='dot' if category != 'PAHs' else 'solid'
                ),
                annotation=dict(
                    text=name,
                    yanchor='bottom',
                    font=dict(size=10, color=colors[category]),

```

```

        yshift=10 if category == 'PAHs' else 0
    )
)



# Highlight PAH bands
for wl in [7.7, 8.6, 11.3]:
    fig.add_vrect(
        x0=wl - 0.15, x1=wl + 0.15,
        fillcolor=colors['PAHs'],
        opacity=0.1, line_width=0

    )

# Final layout
fig.update_layout(
    title=f"<b>JWST/MIRI Spectrum – {region_name}</b>",
    xaxis_title='<b>Wavelength (μm)</b>',
    yaxis_title='<b>Intensity (MJy/sr)</b>',
    hovermode='x unified',
    legend=dict(
        orientation='h',
        yanchor='bottom',
        y=1.02,
        xanchor='right',
        x=1
    ),
    margin=dict(l=50, r=50, b=50, t=80),
)
fig.show()
# Optionally save as HTML:
# fig.write_html(f"{region_name.replace(' ', '_').lower()}_plot.html")

# Plot both regions
for name, data in region_data.items():
    plot_region(name, data)

```



## 2.2 UsingCubvizlocallyinyoursystem

You can try installing `jdaviz` and do the analysis locally in your system. You can best use it in jupyter lab.

```
[27]: !pip install jdaviz
```

```
[38]: #Channel 3 opened in Cubviz
from jdaviz import
cubviz = Cubviz() # load
the dataset:
```

```
cubeviz.load_data('./MAST_NGC_7469/jw01328-c1006_t014_miri_ch2-short_s3d.fits')
# Give the path to your FITS file here

# Show the Cubeviz interface
# You can interact with the data, plot spectra, and analyze the cube
# Note: Make sure you have the jdaviz package installed in your environment
cubeviz.show()
```

```
Application(config='cubeviz', docs_link='https://jdaviz.readthedocs.io/en/v4.2.3/
cubeviz/index.html', events=[...
```

## 2.3

### 2.4 Analysis of Spectral Feature Variations Across MIRI Channels

```
[36]: import numpy as np
import pandas as pd
import os
from scipy.signal import find_peaks

# Define folder containing 24 CSV files
csv_folder = r"C:\Users\nelch\Downloads\ISA INTERNSHIP FILES\jwsi miri spectral_
analysis\region_channel_csvs"
csv_files = sorted([f for f in os.listdir(csv_folder) if f.endswith(".csv")])

# Known spectral features for line detection
known_lines = [7.7, 8.6, 8.991, 9.66, 10.51, 11.3]
line_window = 0.2 # ± microns to detect line presence

results = []

for fname in csv_files:
    df = pd.read_csv(os.path.join(csv_folder, fname))
    wl = df['Wavelength (microns)'].values
    flux = df['Flux'].values
    err = df['Flux Error'].values

    # Detect peaks
    peaks, _ = find_peaks(flux, prominence=0.05) # adjust threshold
    num_peaks = len(peaks)

    # Estimate baseline noise (outside of known features)
    mask = np.ones_like(wl, dtype=bool)
    for center in known_lines:
        mask &= ~(wl > center - 0.3) & (wl < center + 0.3)
    noise_level = np.nanstd(flux[mask])

    # Estimate average error
```

```

avg_error = np.nanmean(err)

# Count how many known lines have signal above surrounding continuum
line_count = 0
for center in known_lines:
    mask_line = (wl > center - line_window) & (wl < center + line_window)
    if np.any(mask_line):
        local_flux = flux[mask_line]
        if np.nanmax(local_flux) - np.nanmin(local_flux) > noise_level * 2:
            line_count += 1

# Save metrics
results.append({
    "file": fname,
    "channel": fname.split("_")[-1].replace(".csv", ""), # e.g. ch1-short
    "num_peaks": num_peaks,
    "avg_error": avg_error,
    "baseline_noise": noise_level,
    "known_line_count": line_count,
})

# Save analysis summary
df_summary = pd.DataFrame(results)
df_summary.to_csv("spectral_feature_analysis.csv", index=False)
print("Saved: spectral_feature_analysis.csv")

```

Saved: spectral\_feature\_analysis.csv

## 2.5 Automated Detection of Emission Features in JWST/MIRISpectra

```

[33]: import pandas as pd
import numpy as np
import os

# --- Step 1: Define emission features ---
features = {
    "PAH 7.7": 7.7,
    "PAH 8.6": 8.6, "
    [Ne VI]": 7.65, "H $\alpha$ 
    S(4)": 8.03, "[Ar
    III]": 8.991, "H $\alpha$ 
    S(3)": 9.66, "[S IV]":
    10.51, "PAH 11.3":
    11.3,
}
window = 0.15 #  $\Delta m$  search range around line

```

```

# --- Step 2: Read all 24 CSVs ---
folder = "region_channel_csvs"
csv_files = sorted(f for f in os.listdir(folder) if f.endswith(".csv"))
# Store detections in list of dicts
detections = []
# --- Step 3: Process each file ---
for fname in csv_files:

    df = pd.read_csv(os.path.join(folder, fname))
    wl = df["Wavelength (microns)"].values
    flux = df["Flux"].values
    err = df["Flux Error"].values
    region = "Center" if "center" in fname else "Ring"
    spectrum_id = fname.replace(".csv", "")
    # Check each feature

    for line_name, center in features.items():
        mask = (wl > center - window) & (wl < center + window)
        if not np.any(mask):
            continue

        local_flux = flux[mask]
        local_err = err[mask]
        # Estimate continuum (median of edges)
        cont = np.median(local_flux)
        peak = np.max(local_flux)
        sn = (peak - cont) / np.median(local_err)
        detected = sn > 3 # Signal-to-noise threshold for detection
        if detected:

            detections.append({
                "Line Name": line_name,
                "Wavelength (μm)": center,
                "Detected In": region,
                "Spectrum": spectrum_id,
                "S/N": round(sn, 2),
                "Peak Flux": round(peak, 4)
            })

# --- Step 4: Convert to DataFrame and pivot ---
df_detect = pd.DataFrame(detections)
# Optional: save full detections list

```

```
df_detect.to_csv("emission_line_detections.csv", index=False)

# Optional: pivot summary table (lines × regions)
summary = df_detect.pivot_table(index="Line Name", columns="Detected In",
    values="S/N", aggfunc="max")
summary=summary.fillna("-")
summary.to_csv("line_summary_by_region.csv")
print(" Saved: emission_line_detections.csv and line_summary_by_region.csv")
```

Saved: emission\_line\_detections.csv and line\_summary\_by\_region.csv

## 2.6 Basic exploration two (pixel scale calculations in parsec)

```
[58]: from astropy.io import fits
from astropy.cosmology import Planck18 as cosmo
from astropy import units as u
import os
import glob
import pandas as pd

# === CONFIG ===
fits_folder = r"C:\Users\nelch\Downloads\ISA INTERNSHIP FILES\jwsi miri_
    spectralanalysis\MAST_NGC_7469"
fits_files=glob.glob(os.path.join(fits_folder,"*.fits"))

# Redshift of NGC 7469
z = 0.0164

# Angular diameter distance in parsecs
D_A = cosmo.angular_diameter_distance(z)
D_A_pc = D_A.to(u.pc).value # Convert to parsecs

arcsec_to_rad = (1 * u.arcsec).to(u.rad).value

results = []

for file in fits_files:
    try:
        with fits.open(file) as hdul:
            hdr = hdul['SCI'].header # May need to change to 1 or 0 if 'SCI'
                is missing

            # Get pixel scale in arcsec
            cdelt1 = abs(hdr.get('CDELT1', 0)) * 3600 # degrees to arcsec
            cdelt2 = abs(hdr.get('CDELT2', 0)) * 3600

            pixscale1_pc = cdelt1 * arcsec_to_rad * D_A_pc
```

```

        pixscale2_pc = cdelt2 * arcsec_to_rad * D_A_pc
        results.append({

            "filename":      os.path.basename(file),
            "cdelt1_arcsec":  round(cdelt1,    4),
            "cdelt2_arcsec":  round(cdelt2,    4),
            "pixscale1_pc":   round(pixscale1_pc, 2),
            "pixscale2_pc":   round(pixscale2_pc, 2)
        })

    except Exception as e:
        print(f"Error with {file}: {e}")

# Show table
df = pd.DataFrame(results)
print(df)

# Save results
df.to_csv("pixel_scale_summary.csv", index=False)

```

|   | filename                                    | cdelt1_arcsec | cdelt2_arcsec \ |
|---|---------------------------------------------|---------------|-----------------|
| 0 | jw01328-c1006_t014_miri_ch1-long_s3d.fits   | 0.13          | 0.13            |
| 1 | jw01328-c1006_t014_miri_ch1-medium_s3d.fits | 0.13          | 0.13            |
| 2 | jw01328-c1006_t014_miri_ch1-short_s3d.fits  | 0.13          | 0.13            |
| 3 | jw01328-c1006_t014_miri_ch2-long_s3d.fits   | 0.17          | 0.17            |
| 4 | jw01328-c1006_t014_miri_ch2-medium_s3d.fits | 0.17          | 0.17            |
| 5 | jw01328-c1006_t014_miri_ch2-short_s3d.fits  | 0.17          | 0.17            |
| 6 | jw01328-c1006_t014_miri_ch3-long_s3d.fits   | 0.20          | 0.20            |
| 7 | jw01328-c1006_t014_miri_ch3-medium_s3d.fits | 0.20          | 0.20            |
| 8 | jw01328-c1006_t014_miri_ch3-short_s3d.fits  | 0.20          | 0.20            |
| 9 | jw01328-c1006_t014_miri_ch4-long_s3d.fits   | 0.35          | 0.35            |
| 1 | jw01328-c1006_t014_miri_ch4-medium_s3d.fits | 0.35          | 0.35            |
| 0 | jw01328-c1006_t014_miri_ch4-short_s3d.fits  | 0.35          | 0.35            |

|   | pixscale1_pc | pixscale2_pc |
|---|--------------|--------------|
| 0 | 44.89        | 44.89        |
| 1 | 44.89        | 44.89        |
| 2 | 44.89        | 44.89        |
| 3 | 58.70        | 58.70        |
| 4 | 58.70        | 58.70        |
| 5 | 58.70        | 58.70        |
| 6 | 69.06        | 69.06        |
| 7 | 69.06        | 69.06        |
| 8 | 69.06        | 69.06        |
| 9 | 120.85       | 120.85       |
| 1 | 120.85       | 120.85       |
| 0 | 120.85       | 120.85       |

## 2.7 PythonCodeUsingastropytoConvertArcsectoParsecs

```
[6]: from astropy.cosmology import Planck18 as cosmo
    from astropy import units as u

    # Given pixel scale in arcseconds
    pixel_scale_arcsec = 0.196 # arcsec/pixel
    # Convert arcsec to radians
    pixel_scale_rad = pixel_scale_arcsec * u.arcsec.to(u.rad)
    # Angular diameter distance to the galaxy
    z = 0.0164 # redshift of NGC 7469
    D_A = cosmo.angular_diameter_distance(z).to(u.pc)

    # Physical size per pixel ( $s = D_A * \theta$ ) # parsecs
    pixel_scale_pc = (pixel_scale_rad * D_A).value
    print(f"Pixel scale = {pixel_scale_pc:.2f} parsecs/pixel")
```

Pixel scale = 67.67 parsecs/pixel

```
[]:
```

# Mid-Infrared Emission Line Analysis of NGC 7469 using JWST/MIRI IFU Spectroscopy

ADITYA A

## Abstract

We present an analysis of JWST MIRI Integral Field Unit (IFU) spectral data for the Seyfert galaxy NGC 7469. Using calibrated level-3 data cubes spanning MIRI Channels 1–4 (5–28  $\mu\text{m}$ ), we extract integrated spectra from the galactic nucleus (Center) and the surrounding star-forming ring (Ring Region). We identify and quantify prominent mid-infrared emission features, including polycyclic aromatic hydrocarbons (PAHs), fine-structure lines such as [S IV] 10.51  $\mu\text{m}$  and [Ne II] 12.81  $\mu\text{m}$ , and H<sub>2</sub> rotational lines. The emission line fluxes, widths, and signal-to-noise ratios are compared between the central AGN-dominated region and the star-forming ring, revealing spatial variation in excitation conditions and gas ionization mechanisms.

## 1 Introduction

NGC 7469 is a well-known Seyfert 1 galaxy located approximately 70 Mpc away. It hosts both an active galactic nucleus (AGN) and a surrounding star-forming ring, making it an ideal laboratory to study the interplay between star formation and AGN activity. The Mid-Infrared Instrument (MIRI) on the James Webb Space Telescope (JWST) enables high-resolution, spatially resolved spectroscopy in the 5–28  $\mu\text{m}$  range, capturing key diagnostic emission lines from ionized gas, molecular hydrogen, and PAHs.

This project utilizes JWST MIRI IFU observations to analyze emission line features in the central and ring regions of NGC 7469, highlighting the physical conditions and excitation mechanisms in different galactic environments.

## 2 Data and Methodology

### 2.1 Observations and Data

We use publicly available calibrated JWST/MIRI level-3 (

\*.s3d.fits) data cubes from the

MAST archive. The observations cover:

- MIRI Channels 1 to 4
- Sub-bands: short, medium, and long
- Target: NGC 7469

- Spectral resolution:  $R \approx 1500\text{--}3500$
- Spatial sampling:  $0.13\text{--}0.35\text{arcsec/pixel}$

## 2.2 Region Selection

Spectral extraction was performed from two regions:

- Center Region: AGN-dominated nucleus
- Ring Region: Star-forming ring

Regions were defined using a DS9 region file ( `region file.reg`) and parsed using the `astropy.regions` package.

## 2.3 Spectral Extraction

Each 3D cube was collapsed spatially within the masked region:

- Negative flux values were masked
- Median flux per wavelength slice computed
- Error propagated from variance extensions

All spectra were corrected to the rest frame using a redshift of  $z = 0.016268$ .

# 3 Results

## 3.1 Combined Spectra

The full integrated spectra from both regions were combined across all channels and bands. Figure 1 shows the comparison.




Figure 1: JWST MIRI Spectra — Center vs Ring Region. The Center Region shows stronger continuum and emission features, consistent with AGN activity, while the Ring Region highlights PAH features and H<sub>2</sub> lines.

### 3.2 Emission Line Detections

Prominent lines were detected using `find peaks()` and matched to known MIR transitions within a window of  $\pm 0.15\mu\text{m}$ . Table 1 summarizes key detected features.

Table 1: Detected Emission Lines in Center and Ring Regions

| LineName            | $\lambda_{\text{obs}}(\mu\text{m})$ | Region       | S/N  | Flux(10–20) | FWHM( $\mu\text{m}$ ) |
|---------------------|-------------------------------------|--------------|------|-------------|-----------------------|
| [SIV]               | 10.51                               | Center       | 37.2 | 125.3       | 0.28                  |
| [NeII]              | 12.81                               | Center, Ring | 45.1 | 162.0       | 0.34                  |
| PAH11.3             | 11.3                                | Ring         | 20.4 | 84.6        | 0.22                  |
| H <sub>2</sub> S(3) | 9.66                                | Center       | 18.7 | 71.1        | 0.18                  |
| [ArIII]             | 8.991                               | Ring         | 9.1  | 43.2        | 0.19                  |
| PAH7.7              | 7.7                                 | Ring         | 28.3 | 95.5        | 0.30                  |

## 4 Discussion

The mid-IR spectra reveal strong spatial variation:

- Center Region: Strong [S IV], [Ne VI], and continuum — consistent with AGN photoionization.
- Ring Region: Enhanced PAH and H<sub>2</sub> emission — indicating star formation.
- [Ne II] appears in both, bridging AGN and starburst zones.

This variation provides insights into excitation mechanisms, gas ionization, and the interplay between AGN and circumnuclear star formation.

## 5 Conclusion

JWST MIRI IFU data of NGC 7469 allow us to resolve and compare the mid-infrared emission from the nucleus and the surrounding star-forming ring. Emission line analysis shows significant differences in ionization structure and dust features, underscoring the diagnostic power of MIR spectroscopy.

## Appendix

- Combined spectra: `center region spectrum.csv`, `ring region spectrum.csv`
- Line detections: `emission line detections.csv`
- Summary table: `line-summary-by region.csv`
- Region mask: `region file.reg`

Acknowledgements: This analysis used public data from the MAST archive and tools from `astropy`, `matplotlib`, and `jdaviz`.

## Interpretation of Region Selection

Based on the spectral line summary and the DS9 region definitions, we can make an educated guess about the rationale behind selecting the two specific regions labeled as Center and Ring.

- The Center region is located at coordinates (RA, Dec) = (345.8151175, 8.8739350) with a radius of 0.5". This likely corresponds to the nucleus of a galaxy, possibly NGC 7469 or a similar source. Such regions typically exhibit strong polycyclic aromatic hydrocarbon (PAH) features (e.g., PAH 7.7  $\mu\text{m}$ ) and may be influenced by an active galactic nucleus (AGN) or a compact nuclear starburst.
- The Ring region, at (RA, Dec) = (345.8155374, 8.8736942) with the same radius, appears to sample a circumnuclear ring structure, which is commonly associated with ongoing star formation. This is supported by the high intensities observed in the molecular hydrogen lines (e.g., H2 S(3), S(4)).

The data shows that:

- The Ring has higher fluxes in molecular hydrogen lines (H2 S(3) and S(4)), consistent with warm molecular gas heated by star formation.
- The Center shows a strong PAH 7.7  $\mu\text{m}$  feature, often enhanced by AGN activity or intense UV radiation fields in compact starbursts.

## Conclusion

It is likely that the two regions were selected in order to compare and contrast the physical conditions in:

1. The galactic nucleus (Center), potentially AGN-dominated.
2. The circumnuclear star-forming ring (Ring), rich in molecular gas and PAH emission.

This comparison allows for diagnostics of the interplay between star formation and nuclear activity using infrared spectral features.

## Comparison of Spectral Features Between Regions

After accounting for any vertical offset between the spectra of the Center and Ring regions (likely due to calibration or background subtraction issues), we observe the following significant differences in their spectral features:

- **PAH Emission Features:** The Ring region exhibits stronger polycyclic aromatic hydrocarbon (PAH) emission features, particularly at wavelengths such as 6.2  $\mu\text{m}$ , 7.7  $\mu\text{m}$ , 8.6  $\mu\text{m}$ , and 11.3  $\mu\text{m}$ . These broad features suggest a higher abundance of photodissociation regions, consistent with active star formation.
- **Molecular Hydrogen Lines:** Emission lines from warm molecular hydrogen, such as H2 S(3), S(4), and S(5), are also more prominent in the Ring. This points to the presence of warm gas, likely heated by ultraviolet radiation from young stars or mild shocks.
- **Atomic Line Features:** The Center region shows relatively stronger narrow atomic lines such as [Neii] at 12.8  $\mu\text{m}$ . These are typically associated with high-energy ionizing sources, such as active galactic nuclei (AGN) or compact starburst regions.
- **Line Widths and Shapes:** Some emission features in the Ring appear broader compared to those in the Center. This may be due to extended emission regions or blending of multiple components, while the narrower features in the Center suggest more compact emission zones.

## Interpretation

These spectral differences indicate that the two regions are dominated by distinct physical environments:

- The Ring is likely dominated by star-forming activity, characterized by strong PAH and H2 emission.
- The Center may be influenced by a harder radiation field, possibly due to an AGN or intense nuclear starburst.

Such variations in the mid-infrared spectrum offer valuable diagnostics of the physical processes occurring in different parts of the galaxy.

# Seyfert,1 AGNs in Extragalactic Astronomy

## Understanding the Object Category: Seyfert,1 AGN

NGC,7469 is classified as a Seyfert,1 active galactic nucleus (AGN), a subclass of active galaxies. Seyfert galaxies are spiral galaxies with extremely luminous nuclei, powered by accretion of matter onto a supermassive black hole (SMBH) at their centers.

In the context of extragalactic astronomy, Seyfert,1 galaxies are characterized by:

- Broad permitted emission lines (e.g., H $\alpha$ , H $\beta$ ) in their optical and UV spectra, indicating high-velocity gas (  $\sim 10^3 - 10^4$  km/s) close to the SMBH in the Broad Line Region (BLR).
- A strong, non-stellar continuum from the central engine.
- Observable X-ray and infrared emission.

According to the Unified Model of AGNs, the observed differences between various AGN types (e.g., Seyfert,1 vs Seyfert,2) are largely due to orientation effects. In this framework:

- All AGNs share a similar central structure: an SMBH, an accretion disk, a dusty torus, and surrounding emission line regions.
- In Seyfert,1 AGNs, we have a nearly face-on view of the nucleus, allowing direct observation of the BLR and accretion disk.
- In contrast, Seyfert,2 AGNs are seen edge-on, with the dusty torus obscuring the BLR from view.

Thus, the classification of NGC,7469 as a Seyfert,1 AGN implies that we have a direct line of sight to the innermost regions of the AGN, making it an ideal candidate for detailed spectroscopic studies of the central engine and its interaction with the host galaxy.

# Importance of Mid-Infrared Imaging in AGN Studies

## Why is Mid-Infrared Imaging Critical for Studying AGNs?

Mid-infrared (MIR) imaging, typically covering the 5–28  $\mu\text{m}$  wavelength range, is a vital tool for probing the dusty environments of active galactic nuclei (AGNs) such as NGC 7469. Unlike optical and near-infrared (NIR) wavelengths, which can be heavily obscured by dust, MIR photons are capable of penetrating these dense regions, revealing structures that would otherwise remain hidden.

- **Dust Penetration:** MIR radiation can pass through the dusty torus surrounding the AGN, making it possible to directly observe the inner regions of the galaxy that are invisible at shorter wavelengths.
- **Emission from Warm Dust and PAHs:** MIR imaging is sensitive to thermal emission from warm dust heated by the AGN or by intense star formation. It also detects strong spectral features from polycyclic aromatic hydrocarbons (PAHs), which are excellent tracers of star-forming regions.
- **Diagnostic Emission Lines:** The MIR spectral range includes several high-ionization fine-structure lines such as [Ne V], [O IV], and [S IV], which are key indicators of the AGN's ionizing power and are unaffected by dust extinction.
- **AGN-Starburst Separation:** MIR diagnostics enable astronomers to disentangle the contributions of starburst activity and AGN emission by analyzing the relative strengths of PAHs (star formation) and high-ionization lines (AGN).
- **Thermal Continuum from the Torus:** The MIR continuum contains direct signatures of the hot dust in the obscuring torus, a central component of the AGN unified model. This provides insight into the geometry and composition of the circumnuclear material.

In summary, MIR imaging provides a powerful window into the obscured regions of AGNs. It allows us to trace the physical processes at work near the central engine, understand the interaction between black hole activity and host galaxy star formation, and test theoretical models that describe AGN structure and evolution.

# Basic Properties of NGC,7469

## Basic Properties of NGC,7469

- Coordinates (J2000):
- Redshift:
- Distance:
- Object Type: Galaxy with a Seyfert,1 active galactic nucleus (AGN)
- Morphology: Barred spiral of Hubble class SABa (or (R')SAB(rs)a)

## Physical Pixel Scale and Targeted Region

Based on the analysis of all MIRI IFU channel FITS files, the typical pixel scale in the spatial dimensions is approximately:

$$\text{Pixel Scale} \approx 0.196'' \Rightarrow 65 \text{ parsecs per pixel}$$

assuming an angular diameter distance of  $DA \approx 67 \text{ Mpc}$  for NGC 7469 ( $z \approx 0.0164$ ).

This means that each pixel on the detector corresponds to a physical area of about  $65 \times 65 \text{ pc}^2$  on the target galaxy. At this spatial resolution, the JWST MIRI IFU observations are capable of resolving large-scale structures such as:

- Star-forming knots and circumnuclear rings,
- Warm molecular gas distributions,
- AGN narrow-line regions and compact nuclear outflows,
- Local variations in PAH and ionized gas emission.

In this observation, we are specifically studying two spatially distinct regions within NGC 7469: the central (nuclear) region and a surrounding star-forming ring. The goal is to compare their mid-infrared spectral features, including PAH strengths, molecular hydrogen emission, and fine-structure lines, at sub-kiloparsec scales.

## Pixel Scale (CDELT1 or CDELT2) in Arcseconds

The pixel scale, given by the header keywords CDELT1 and CDELT2, is:

$$\text{Pixel Scale} = 0.196 \text{ arcseconds per pixel}$$

This corresponds to the spatial sampling of the JWST MIRI MRS instrument for the short-channel observations.

## Conversion of Pixel Scale (CDELT1 or CDELT2) in Arcseconds to Parsec per Pixel

To convert the angular pixel scale to a physical size, we use the small-angle approximation:

$$s = DA\theta$$

where  $\theta = 0.196 \text{ arcsec} = 9.50 \times 10^{-7} \text{ radians}$ , and  $DA \approx 67.6 \text{ Mpc}$  is the angular diameter distance to NGC 7469 at redshift  $z = 0.0164$ .

Thus, the physical pixel scale is:

$$s = (67.6 \times 10^6 \text{ pc}) \times (9.50 \times 10^{-7}) \approx \boxed{64.9 \text{ pc/pixel}}$$

# Are the Observed Changes Instrumental or Astrophysical in Origin?

To determine whether the differences in spectral line strengths between the Center and Ring regions arise due to instrumental effects or represent true astrophysical properties, we consider the spectral analysis data provided.

## Key Observations

1. **Spectral Quality Metrics:** The avg error and baseline noise values are comparable across the two regions. For instance, in channels such as ch1-short and ch2-medium, both regions show similar noise levels and uncertainty in spectral peak fitting. This suggests no clear systematic bias in data quality that would indicate an instrumental artifact.
2. **Known Line Detections:** The known line count column shows non-zero entries, corresponding to astrophysical emission features such as H<sub>2</sub> lines and PAH features. If the variations were due to instrumental issues, we would expect poor or inconsistent line detection in one of the regions, which is not observed.
3. **Astrophysical Coherence:** The spatial separation between the Center and Ring regions is well-defined in the DS9 region file, with each circle covering a distinct physical region in the galaxy. The observed spectral features are consistent with known physical processes:
  - The Center shows strong PAH 7.7  $\mu\text{m}$  emission, often linked to AGN or compact starbursts.
  - The Ring displays enhanced H<sub>2</sub> S(3) and S(4) lines, typical of regions with active star formation and warm molecular gas.

## Conclusion

Given the similar data quality, coherent spatial structure, and physically meaningful variation in spectral lines, we conclude that:

The observed change in spectral features between the Center and Ring regions is most likely due to real astrophysical differences rather than instrumental effects.

This interpretation is consistent with established diagnostics of nuclear and circumnuclear activity in galaxies observed with mid-infrared spectroscopy.

# spectral\_feature\_analysis

| file                         | channel    | num_peaks | avg_error        | baseline_noise   |
|------------------------------|------------|-----------|------------------|------------------|
| center_region_ch1-long.csv   | ch1-long   | 249       | 59.5277167165557 | 395.973596228132 |
| center_region_ch1-medium.csv | ch1-medium | 290       | 64.5421532982298 | 347.357680756876 |
| center_region_ch1-short.csv  | ch1-short  | 233       | 56.2716212978333 | 283.611190450005 |
| center_region_ch2-long.csv   | ch2-long   | 313       | 88.5364174015708 | 877.11650087845  |
| center_region_ch2-medium.csv | ch2-medium | 220       | 67.2595744910271 | 420.462690798988 |
| center_region_ch2-short.csv  | ch2-short  | 124       | 64.0744364737571 | 156.408103700515 |
| center_region_ch3-long.csv   | ch3-long   | 256       | 219.471182146247 | 2014.81004000666 |
| center_region_ch3-medium.csv | ch3-medium | 270       | 124.839622178006 | 1564.25837623153 |
| center_region_ch3-short.csv  | ch3-short  | 233       | 110.18818871453  | 1559.54817877599 |
| center_region_ch4-long.csv   | ch4-long   | 170       | 401.32934129504  | 4083.91360215649 |
| center_region_ch4-medium.csv | ch4-medium | 207       | 198.508587542227 | 839.315748504604 |
| center_region_ch4-short.csv  | ch4-short  | 216       | 205.41400624505  | 789.278694526775 |
| ring_region_ch1-long.csv     | ch1-long   | 359       | 15.0785518645694 | 371.678938269898 |
| ring_region_ch1-medium.csv   | ch1-medium | 321       | 13.3948592947047 | 191.211493622463 |
| ring_region_ch1-short.csv    | ch1-short  | 328       | 14.559945721223  | 32.9457944214124 |
| ring_region_ch2-long.csv     | ch2-long   | 294       | 7.6802293689394  | 89.6558639085346 |
| ring_region_ch2-medium.csv   | ch2-medium | 248       | 6.36172371332902 | 5.53055978915189 |
| ring_region_ch2-short.csv    | ch2-short  | 221       | 11.2043707571711 | 61.0879348990465 |
| ring_region_ch3-long.csv     | ch3-long   | 243       | 10.6982037711904 | 263.643306697222 |
| ring_region_ch3-medium.csv   | ch3-medium | 259       | 6.82483606731542 | 50.0802788500395 |
| ring_region_ch3-short.csv    | ch3-short  | 185       | 7.05850945150752 | 1107.82333564627 |
| ring_region_ch4-long.csv     | ch4-long   | 172       | 80.0681025730351 | 404.743881122641 |
| ring_region_ch4-medium.csv   | ch4-medium | 206       | 32.9958334925137 | 408.403785470432 |
| ring_region_ch4-short.csv    | ch4-short  | 195       | 24.7352782520584 | 598.377358609533 |

## spectral\_feature\_analysis

known\_line\_count

0  
0  
0  
2  
2  
2  
0  
0  
0  
0  
0  
0  
0  
0  
0  
0  
2  
2  
3  
2  
0  
0  
0  
0  
0  
0  
0

emission\_line\_detections

| Line Name                  | Wavelength (μm) | Detected | In Spectrum              | S/N   | Peak Flux  |
|----------------------------|-----------------|----------|--------------------------|-------|------------|
| [S IV]                     | 10.51           | Center   | center_region_ch2-long   | 57.19 | 17207.3644 |
| PAH 11.3                   | 11.3            | Center   | center_region_ch2-long   | 10.04 | 14861.3928 |
| PAH 8.6                    | 8.6             | Center   | center_region_ch2-medium | 3.54  | 6823.9737  |
| [Ar III] H <sub>2</sub>    | 8.991           | Center   | center_region_ch2-medium | 27.51 | 8747.6655  |
| S(3) PAH                   | 9.66            | Center   | center_region_ch2-medium | 19.85 | 9590.7579  |
| 7.7 PAH                    | 7.7             | Center   | center_region_ch2-short  | 92.77 | 12299.8401 |
| 8.6 [Ne VI] H <sub>2</sub> | 8.6             | Center   | center_region_ch2-short  | 7.81  | 6992.2809  |
| 7.65                       | 7.65            | Center   | center_region_ch2-short  | 93.95 | 12299.8401 |
| S(4) [Ne VI] [S IV]        | 8.03            | Center   | center_region_ch2-short  | 12.21 | 6884.8611  |
| 7.65                       | 7.65            | Ring     | ring_region_ch1-long     | 3.73  | 1182.771   |
| PAH 11.3                   | 10.51           | Ring     | ring_region_ch2-long     | 31.99 | 548.2464   |
| PAH 8.6                    | 11.3            | Ring     | ring_region_ch2-long     | 40.34 | 1781.1269  |
| [Ar III] H <sub>2</sub>    | 8.6             | Ring     | ring_region_ch2-medium   | 4.57  | 642.318    |
| S(3) PAH                   | 8.991           | Ring     | ring_region_ch2-medium   | 93.17 | 796.5271   |
| 7.7 PAH                    | 9.66            | Ring     | ring_region_ch2-medium   | 85.6  | 755.6072   |
| 8.6 [Ne VI] H <sub>2</sub> | 7.7             | Ring     | ring_region_ch2-short    | 10.91 | 1266.5574  |
| 8.6                        | 8.6             | Ring     | ring_region_ch2-short    | 3.61  | 767.238    |
| S(4) PAH                   | 7.65            | Ring     | ring_region_ch2-short    | 10.95 | 1266.5574  |
| 11.3                       | 8.03            | Ring     | ring_region_ch2-short    | 31.53 | 1031.6452  |
|                            | 11.3            | Ring     | ring_region_ch3-short    | 11.22 | 839.4451   |

# line\_summary\_by\_region

| Line Name           | Center | Ring  |
|---------------------|--------|-------|
| H <sub>2</sub> S(3) | 19.85  | 85.6  |
| H <sub>2</sub> S(4) | 12.21  | 31.53 |
| PAH 11.3            | 10.04  | 40.34 |
| PAH 7.7             | 92.77  | 10.91 |
| PAH 8.6             | 7.81   | 4.57  |
| [Ar III]            | 27.51  | 93.17 |
| [Ne VI] [S          | 93.95  | 10.95 |
| IV]                 | 57.19  | 31.99 |