

ANSWER KEYS

- | | | | | | | | |
|---------|---------|---------|---------|---------|---------|---------|---------|
| 1. (2) | 2. (1) | 3. (4) | 4. (1) | 5. (3) | 6. (2) | 7. (1) | 8. (1) |
| 9. (4) | 10. (1) | 11. (3) | 12. (4) | 13. (2) | 14. (1) | 15. (1) | 16. (1) |
| 17. (1) | 18. (3) | 19. (2) | 20. (1) | 21. (1) | 22. (1) | 23. (2) | 24. (1) |
| 25. (3) | 26. (1) | 27. (2) | 28. (1) | 29. (4) | 30. (4) | | |

1. (2)

$$y^2 = a \left(x + \frac{\sqrt{a}}{2} \right) = ax + \frac{a^{3/2}}{2} \dots (1)$$

$$\Rightarrow 2yy' = a$$

Put in equation (1) we get,

$$y^2 = (2yy')x + \frac{(2yy')^{3/2}}{2}$$

$$(y^2 - 2xyy') = \frac{(2yy')^{3/2}}{2}$$

Squaring both the sides we get,

$$(y^2 - 2xyy')^2 = 2y^3(y')^3$$

\therefore Order = 1

Degree = 3

Degree - order = 3 - 1 = 2

2. (1)

$$\frac{dy}{dx} = y + 7 \Rightarrow \frac{dy}{dx} - y = 7$$

I.F. = e^{-x}

$$ye^{-x} = \int 7e^{-x} dx$$

$$\Rightarrow ye^{-x} = -7e^{-x} + c$$

$$\Rightarrow y = -7 + ce^x$$

$$-7 + 7e^x = -7 + 8e^x$$

$$\Rightarrow e^x = 0$$

3. (4)

$$\text{Given, } \frac{dy}{dx} + \frac{2^x - y(2^y - 1)}{2^x - 1} = 0 \quad x, y > 0, y(1) = 1$$

Now rearranging and integrating both side of $\frac{dy}{dx} = -\frac{2^x(2^y - 1)}{2^y(2^x - 1)}$, we get

$$\Rightarrow \int \frac{2^y}{2^y - 1} dy = - \int \frac{2^x}{2^x - 1} dx$$

$$\Rightarrow \frac{1}{\ln 2} \int \frac{2^y \ln 2}{2^y - 1} dy = - \frac{1}{\ln 2} \int \frac{2^x \ln 2}{2^x - 1} dx$$

$$\Rightarrow \frac{1}{\ln 2} \ln|2^y - 1| = - \frac{1}{\ln 2} \ln|2^x - 1| + C$$

$$\text{At } x = 1, y = 1$$

Putting this values in above equation we get $C = 0$

$$\text{So, } \ln|2^y - 1| + \ln|2^x - 1| = 0$$

$$\Rightarrow (2^x - 1)(2^y - 1) = 1$$

$$\Rightarrow 2^y - 1 = \frac{1}{2^x - 1}$$

At $x = 2$

$$\Rightarrow 2^y = \frac{1}{3} + 1 = \frac{4}{3}$$

$$y = \log_2 \frac{4}{3} = \log_2 4 - \log_2 3 = 2 - \log_2 3$$

4. (I)athongo // mathongo // mathongo // mathongo // mathongo // mathongo // mathongo // mathongo //

Given,

$$\frac{dy}{dx} = \frac{x+y-2}{x-y} = \frac{(x-1)+(y-1)}{(x-1)-(y-1)}$$

Now let $x - 1 = X, y - 1 = Y$

$$\text{So, } \frac{dy}{dx} = \frac{X+Y}{Y-X} \dots\dots\dots(1)$$

Now let $Y = VX$ $\frac{dY}{dX} = V + X \frac{dV}{dX}$

Putting the value in equation (1) we get,

$$V + X \frac{dV}{dX} = \frac{1+V}{1-V}$$

$$\Rightarrow X \frac{dV}{dX} = \frac{V^{2+1}}{1-V}$$

$$\Rightarrow \int \frac{1-V}{1+V^2} dV = \int \frac{dX}{X}$$

$$\Rightarrow \int \frac{dV}{1+V^2} - \frac{1}{2} \int \frac{2VdV}{1+V^2} = \int \frac{dX}{X}$$

$$\Rightarrow \tan^{-1} V - \frac{1}{2} \ln(1 + V^2) = \ln X + c$$

$$\Rightarrow \tan^{-1}\left(\frac{Y}{X}\right) - \frac{1}{2}\ln\left(1 + \frac{Y^2}{X^2}\right) = \ln(X) + c$$

$$\Rightarrow \tan^{-1}\left(\frac{y-1}{x-1}\right) - \frac{1}{2}\ln\left(1 + \frac{(y-1)^2}{(x-1)^2}\right) = \ln(x-1) + c$$

Now given curve passes through $(2, 1)$

So, $0 - \frac{1}{2} \ln 1 = \ln 1 + c \Rightarrow c = 0$

Now given curve also passes through $(k + 1, 2)$

$$\text{So, } \tan^{-1}\left(\frac{1}{2}\right) - \frac{1}{2} \ln\left(1 + \frac{1}{2}\right) = \ln k$$

$$\Rightarrow 2 \tanh^{-1}\left(\frac{1}{k}\right) = \ln\left(\frac{1+k^2}{k^2}\right) + 2 \ln k$$

$$\Rightarrow 2 \tan^{-1} \left(\frac{1}{k} \right) = \ln(1 + k^2)$$

5. (3) mathongo // mathongo // mathongo // mathongo // mathongo // mathongo // mathongo // mathongo // mathongo

Given,

$$(1 + e^{2x}) \frac{dy}{dx} + 2(1 + y^2)e^x = 0$$

$$\Rightarrow \frac{dy}{1+y^2} + \frac{2e^x}{1+e^{2x}} dx = 0 \dots\dots\dots(i)$$

Now integrating both side we get,

$$\Rightarrow \int \frac{dy}{y} + \int \frac{2e^x}{1+2e^x} dx = \int 0$$

$$\Rightarrow \tan^{-1} y + 2 \tan^{-1} e^x = c$$

$$\therefore y(0) = 0$$

so, $C = \frac{\pi}{2} \Rightarrow \tan^{-1} y + 2 \tan^{-1} e^x = \frac{\pi}{2} \dots (ii)$

Now from equation (i), we get $\left(\frac{dy}{dx}\right)_{x=0} = -1$

From equation (ii) we get, $y(\ln \sqrt{3}) = -\frac{1}{\sqrt{3}}$

So, $6[y'(0) + (y \ln \sqrt{3})^2] = 6\left[-1 + \frac{1}{3}\right] = -4$.

We have,

Put $\cos^{-1}(e^{-x}) = \theta, \theta \in [0, \pi]$

$$\Rightarrow \cos \theta = e^{-x}$$

$$\Rightarrow 2 \cos^2 \frac{\theta}{2} - 1 = e^{-x}$$

$$\Rightarrow \cos\left(\frac{\theta}{2}\right) = \sqrt{\frac{e^{-x}+1}{2}} = \sqrt{\frac{e^x+1}{2e^x}}$$

Hence, by (i), we have

$$\cos\left(\frac{\theta}{2}\right)dx = \left(\sqrt{e^{2x}-1}\right)dy$$

$$\Rightarrow \sqrt{\frac{e^x+1}{2e^x}}dx = \sqrt{e^{2x}-1}dy$$

$$\Rightarrow \left(\sqrt{\frac{e^x + 1}{2e^x}} \right) dx = (\sqrt{e^x - 1})(\sqrt{e^x + 1}) dy$$

$$\Rightarrow \frac{1}{\sqrt{2}} \int \frac{dx}{\sqrt{e^x} \sqrt{e^x - 1}} = \int dy$$

Put $e^x = t \Rightarrow dt = e^x dx$

$$\frac{1}{\sqrt{2}} \int \frac{dt}{e^x \sqrt{e^x} \sqrt{e^x - 1}} = \int dy$$

$$\int \frac{dt}{t\sqrt{t^2-1}} = \sqrt{2} \int dy$$

$$\text{Put } t = \frac{1}{z} \Rightarrow \frac{dt}{dz} = -\frac{1}{z^2}$$

$$\int \frac{-\frac{dz}{z^2}}{\sqrt{\quad}} = \sqrt{2} \int dy$$

$$\Rightarrow - \int \frac{dz}{\sqrt{1-z}} = \sqrt{2} \int dy$$

$$\Rightarrow \frac{-2(1-z)^{1/2}}{-1} = \sqrt{2}y + c$$

$$\Rightarrow 2\left(1 - \frac{1}{t}\right)^{1/2} = \sqrt{2}y + c$$

$$\Rightarrow 2(1 - e^{-x})^{1/2} = \sqrt{2}y + c$$

Now, it meets y -axis at $(0, -1)$, hence

$$0 = -\sqrt{2} + c \Rightarrow c = \sqrt{2}$$

Hence,

$$2(1 - e^{-x})^{1/2} = \sqrt{2}(y + 1)$$

It passes through $(\alpha, 0)$

$$2(1 - e^{-\alpha})^{1/2} = \sqrt{2}$$

$$\Rightarrow \sqrt{1 - e^{-\alpha}} = \frac{1}{\sqrt{2}}$$

$$\Rightarrow 1 - e^{-\alpha} = \frac{1}{2}$$

$$\Rightarrow e^{-\alpha} = \frac{1}{2} \Rightarrow e^{\alpha} = 2$$

7. (I) We have,

$$\frac{dy}{dx} = \frac{x \left(\frac{y}{x} \cdot \tan \left(\frac{y}{x} \right) - 1 \right)}{x \tan \left(\frac{y}{x} \right)}$$

$$\Rightarrow \frac{dy}{dx} = \frac{y}{x} - \cot \left(\frac{y}{x} \right)$$

Put $y = vx$

$$\Rightarrow \frac{dy}{dx} = v + x \left(\frac{dv}{dx} \right)$$

Now, we get

$$v + x \left(\frac{dv}{dx} \right) = v - \cot v$$

$$\Rightarrow \int \tan v dv = - \int \frac{dx}{x}$$

$$\Rightarrow \ln |\sec v| = - \ln |x| + c$$

$$\Rightarrow \ln \left| \sec \left(\frac{y}{x} \right) \right| = - \ln |x| + c$$

$$\Rightarrow \ln \left| \sec \left(\frac{y}{x} \right) \right| + \ln |x| = c$$

Now, $y \left(\frac{1}{2} \right) = \frac{\pi}{6}$, then

$$\ln \left| \sec \left(\frac{\pi}{3} \right) \right| + \ln \left(\frac{1}{2} \right) = c$$

$$\Rightarrow \ln 2 + \ln \left(\frac{1}{2} \right) = c$$

$$\Rightarrow \ln 2 - \ln 2 = c$$

$$\Rightarrow c = 0$$

Hence,

$$\therefore \sec \left(\frac{y}{x} \right) = \frac{1}{x}$$

$$\Rightarrow \cos \left(\frac{y}{x} \right) = x$$

$$\Rightarrow y = x \cos^{-1}(x)$$

So, required bounded area

$$= \int_0^{\frac{1}{\sqrt{2}}} \left(\underbrace{\cos^{-1} x}_I \right) \underbrace{xdx}_{II}$$

$$= \left[\cos^{-1} x \cdot \left(\frac{x^2}{2} \right) \right]_0^{\frac{1}{\sqrt{2}}} + \frac{1}{2} \int_0^{\frac{1}{\sqrt{2}}} \left(\frac{x^2}{\sqrt{1-x^2}} \right) dx$$

$$= \frac{\pi}{16} - \frac{1}{2} \int_0^{\frac{1}{\sqrt{2}}} \left(\frac{1-x^2-1}{\sqrt{1-x^2}} \right) dx$$

$$= \frac{\pi}{16} - \frac{1}{2} \int_0^{\frac{1}{\sqrt{2}}} \left(\sqrt{1-x^2} - \frac{1}{\sqrt{1-x^2}} \right) dx$$

$$= \frac{\pi}{16} - \frac{1}{2} \left[\frac{x}{2} \sqrt{1-x^2} + \frac{1}{2} \sin^{-1} x - \sin^{-1} x \right]_0^{\frac{1}{\sqrt{2}}}$$

$$= \frac{\pi}{16} - \frac{1}{2} \left[\frac{x}{2} \sqrt{1-x^2} - \frac{1}{2} \sin^{-1} x \right]_0^{\frac{1}{\sqrt{2}}}$$

$$= \frac{\pi}{16} - \frac{1}{2} \left[\frac{1}{4} - \frac{\pi}{8} \right]$$

$$= \left(\frac{\pi-1}{8} \right) \text{ sq. units}$$

8. (I) $\frac{dy}{dx} - \frac{y+3x}{\ln(y+3x)} + 3 = 0$

$$\frac{dy}{dx} + 3 = \frac{y+3x}{\ln(y+3x)}$$

$$\frac{d}{dx} (y+3x) = \frac{y+3x}{\ln(y+3x)}$$

$$\int \frac{\ln(y+3x)}{(y+3x)} d(y+3x) = \int \frac{dx}{x}$$

Let $\ln(y+3x) = t$

$$\frac{1}{(y+3x)} d(y+3x) = dt$$

$$\int t dt = \int \frac{dx}{x}$$

$$\frac{t^2}{2} = x + c$$

$$\frac{(\ln(y+3x))^2}{2} = x + c$$

9. (4) $\frac{dy}{dx} = \frac{1 + \left(\frac{y}{x}\right)^2}{2\left(\frac{y}{x}\right)}$

Let $y = tx$

$$\Rightarrow t + x \frac{dt}{dx} = \frac{1+t^2}{2t}$$

$$\Rightarrow x \frac{dt}{dx} = \frac{1-t^2}{2t}$$

$$\Rightarrow \int \frac{2t}{1-t^2} dt = \int \frac{dx}{x}$$

$$\Rightarrow \ln|1-t^2| = \ln x + \ln c$$

$$\Rightarrow (1-t^2)(cx) = 1$$

$$\Rightarrow \left(1 - \frac{y^2}{x^2}\right)cx = 1$$

$$y(2) = 0 \Rightarrow c = \frac{1}{2}$$

$$\left(1 - \frac{y^2}{x^2}\right) \cdot \frac{1}{2}x = 1$$

at $x = 8$

$$\left(1 - \frac{y^2}{64}\right) \times \frac{8}{2} = 1$$

$$y = \pm 4\sqrt{3}$$

10. (1)

$$\frac{dy}{dx} = \frac{y}{x} + \frac{y^2}{2x^2}$$

$$\Rightarrow y^2 \frac{dy}{dx} - \frac{1}{y} \cdot \frac{1}{x} = \frac{1}{2x^2}$$

$$\text{Put } -\frac{1}{y} = t \Rightarrow \frac{1}{y^2} \frac{dy}{dx} = \frac{dt}{dx}$$

$$\Rightarrow \frac{dt}{dx} + \left(\frac{1}{x}\right)t = \frac{1}{2x^2}$$

This is a linear differential equation,

with Integrating Factor : $e^{\int \frac{1}{x} dx} = e^{\ln(x)} = x$

So, solution of the linear differential equation is $tx = \int \frac{1}{2x^2} \cdot x dx + C$

$$\Rightarrow -\frac{x}{y} = \frac{1}{2} \ln(x) + C$$

The curve passes through $(1, 2)$

$$\Rightarrow -\frac{1}{2} = \frac{1}{2} \ln(1) + C \Rightarrow C = -\frac{1}{2}$$

Hence, the particular solution to the differential equation is $-\frac{x}{y} = \frac{1}{2} \ln(x) - \frac{1}{2}$

$$\Rightarrow \frac{x}{y} = \frac{1 - \ln(x)}{2} \Rightarrow y = \frac{2x}{1 - \ln(x)}$$

$$\Rightarrow f\left(\frac{1}{2}\right) = \frac{2 \times \frac{1}{2}}{1 - \ln\left(\frac{1}{2}\right)} = \frac{1}{1 + \ln(2)} = \frac{1}{1 + \log_e(2)}$$

11. (3)

Given equation is:

$$(x^2 - 3y^2)dx + 3xydy = 0$$

We can re-write equation as

$$\frac{dy}{dx} = -\frac{(x^2 - 3y^2)}{3xy}$$

$$\Rightarrow \frac{dy}{dx} = \frac{y}{x} - \frac{1}{3} \frac{x}{y} \dots (1)$$

Put $y = vx$

$$\frac{dy}{dx} = v + x \frac{dv}{dx}$$

Equation (1) can be written as

$$v + x \frac{dv}{dx} = v - \frac{1}{3} \frac{1}{v}$$

$$\Rightarrow v dv = -\frac{1}{3x}$$

Integrating both sides, we get

$$\frac{v^2}{2} = -\frac{1}{3} \ln(x) + c$$

$$\Rightarrow \frac{y^2}{2x^2} = -\frac{1}{3} \ln(x) + c \dots (2)$$

$$\therefore y(1) = 1 \quad (\text{given})$$

$$\therefore c = \frac{1}{2} \quad (\text{from equation 2})$$

Equation (2) can be written as

$$\frac{y^2}{2x^2} = -\frac{1}{3} \ln(x) + \frac{1}{2}$$

$$\Rightarrow y^2 = -\frac{2}{3} x^2 \ln(x) + x^2$$

$$\text{Now } y^2(e) = -\frac{2}{3} e^2 \ln(e) + e^2 = -\frac{2}{3} e^2 + e^2 = \frac{e^2}{3}$$

$$\Rightarrow 6y^2(e) = 2e^2$$

12. (4)

$$\text{Given: } \frac{y}{x} \frac{dy}{dx} = \left[\frac{y^2}{x^2} + \frac{\phi\left(\frac{y^2}{x^2}\right)}{\phi'\left(\frac{y^2}{x^2}\right)} \right] \dots (1)$$

$$\text{Let } \frac{y}{x} = t$$

$$\Rightarrow y = xt$$

$$\Rightarrow \frac{dy}{dx} = t + x \cdot \frac{dt}{dx}$$

$$\therefore t \left(t + x \frac{dt}{dx} \right) = \left(t^2 + \frac{\phi(t^2)}{\phi'(t^2)} \right)$$

$$\Rightarrow xt \frac{dt}{dx} = \frac{\phi(t^2)}{\phi'(t^2)}$$

$$\Rightarrow \frac{t \cdot \phi'(t^2)}{\phi(t^2)} dt = \frac{1}{x} dx$$

Integrating both sides

$$\int \frac{t \cdot \phi'(t^2)}{\phi(t^2)} dt = \int \frac{1}{x} dx$$

$$\text{Let } \phi(t^2) = p$$

$$\Rightarrow \phi'(t^2) \cdot 2t = dp$$

$$\Rightarrow \frac{1}{2} \int \frac{1}{p} dp = \int \frac{1}{x} dx$$

$$\Rightarrow \frac{1}{2} \ln p = \ln x + C$$

$$\Rightarrow \frac{1}{2} \ln \phi(t^2) = \ln x + C$$

$$\Rightarrow \frac{1}{2} \ln \left(\phi \left(\frac{y^2}{x^2} \right) \right) = \ln x + C \dots (2)$$

$$\text{If } x = 1, y = -1 \text{ then } C = \frac{1}{2} \ln(\phi(1))$$

Substituting value of C in (2)

$$\frac{1}{2} \ln \left(\phi \left(\frac{y^2}{x^2} \right) \right) = \ln x + \frac{1}{2} \ln(\phi(1))$$

$$\Rightarrow \ln \left(\phi \left(\frac{y^2}{x^2} \right) \right) = \ln x^2 + \ln(\phi(1))$$

If $x = 2$ then

$$\ln \left(\phi \left(\frac{y^2}{4} \right) \right) = \ln 4 + \ln(\phi(1))$$

$$\text{SO, } \phi \left(\frac{y^2}{4} \right) = 4\phi(1)$$

13. (2)

Given, $\frac{dy}{dx} \propto \frac{-y}{x}$

$\Rightarrow \frac{dy}{dx} = \frac{-Ky}{x}$ (where K is proportionality constant)

Now integrating both side,

$\Rightarrow \int \frac{dy}{y} = -K \int \frac{dx}{x}$

$\Rightarrow \ln|y| = -K \ln|x| + C$

If the above equation satisfy (1, 2)

$\Rightarrow \ln 2 = -K \times 0 + C \Rightarrow C = \ln 2$

So, $\ln|y| = -K \ln|x| + \ln 2$

Now it also passes through (8, 1)

$\Rightarrow \ln 1 = -K \ln 8 + \ln 2 \Rightarrow K = \frac{1}{3}$

So, equation becomes $\ln|y| = -\frac{1}{3} \ln|x| + \ln 2$

So, at $x = \frac{1}{8}$

$\Rightarrow \ln|y| = -\frac{1}{3} \ln\left(\frac{1}{8}\right) + \ln 2 = 2 \ln 2$

$\Rightarrow |y| = 4$

14. (1)

Given:

$\frac{dy}{dx} + y \tan x = x \sec x$

This is a linear differential equation.

I. F. = $e^{\int \tan x dx} = \sec x$

Then solution of differential equation is

$y(\sec x) = \int x \sec^2 x dx$

$\Rightarrow y(\sec x) = x \tan x - \int \tan x dx$

$\Rightarrow y(\sec x) = x \tan x - \ln(\sec x) + C$

Given:

$y(0) = 1 \Rightarrow c = 1$

$\therefore y(\sec x) = x \tan x - \ln(\sec x) + 1$

At $x = \frac{\pi}{6}$, we get

$y\left(\sec\left(\frac{\pi}{6}\right)\right) = \left(\frac{\pi}{6}\right) \tan\left(\frac{\pi}{6}\right) - \ln\left(\sec\left(\frac{\pi}{6}\right)\right) + 1$

$\Rightarrow y\left(\frac{2}{\sqrt{3}}\right) = \left(\frac{\pi}{6}\right) \left(\frac{1}{\sqrt{3}}\right) - \ln\left(\frac{2}{\sqrt{3}}\right) + 1$

$\Rightarrow y = \frac{\pi}{12} - \frac{\sqrt{3}}{2} \ln\left(\frac{2}{\sqrt{3}}\right) + \frac{\sqrt{3}}{2}$

$\Rightarrow y = \frac{\pi}{12} - \frac{\sqrt{3}}{2} \left[\ln\left(\frac{2}{\sqrt{3}}\right) - \ln e \right]$

$\Rightarrow y = \frac{\pi}{12} - \frac{\sqrt{3}}{2} \log_e\left(\frac{2}{e\sqrt{3}}\right)$

15. (1)

Given,

$y(x+1)dx - x^2 dy = 0$

$\Rightarrow \frac{x+1}{x^2} dx = \frac{dy}{y}$

$\Rightarrow \left(\frac{1}{x} + \frac{1}{x^2}\right) dx = \frac{dy}{y}$

Now integrating both side we get,

$\log_e x - \frac{1}{x} = \log_e y + c$

Now on using $y(1) = e$ we get, $c = -2$

So, the equation of curve becomes $\log_e x - \frac{1}{x} = \log_e y - 2$

$\Rightarrow y = e^{\ln x - \frac{1}{x} + 2}$

Hence, $\lim_{x \rightarrow 0^+} e^{\ln x - 1 - \frac{1}{x} + 2} = e^{-\infty} = 0$.

16. (I) $\frac{dy}{dx} = \frac{x^2 - 2y}{x}$
 $\Rightarrow \frac{dy}{dx} + \frac{2}{x}y = x$. It is in linear form.
 I.F. = $e^{\int \frac{2}{x} dx} = e^{2 \ln x} = x^2$
 $\therefore y \cdot x^2 = \int x \cdot x^2 dx = \frac{x^4}{4} + c$
 \therefore At $x = 1, y = -2$,
 $\therefore -2 \times 1^2 = \frac{(1)^4}{4} + c$
 $\Rightarrow c = -2 - \frac{1}{4} = -\frac{9}{4}$
 \therefore curve is $y \cdot x^2 = \frac{x^4}{4} - \frac{9}{4}$
 \therefore It passes through $(\sqrt{3}, 0)$.
17. (I) $(1 + x^2) dy = y(x - y) dx$
 $y(0) = 1 \cdot y(2\sqrt{2}) = \beta$
 $\frac{dy}{dx} = \frac{yx - y^2}{1 + x^2}$
 $\frac{dy}{dx} + y \left(\frac{-x}{1 + x^2} \right) = \left(\frac{-1}{1 + x^2} \right) y^2$
 $\frac{1}{y^2} \frac{dy}{dx} + \frac{1}{y} \left(\frac{-x}{1 + x^2} \right) = \frac{-1}{1 + x^2}$
 put $\frac{1}{y} = t$ then $\frac{-1}{y^2} \frac{dy}{dx} = \frac{dt}{dx}$
 $\frac{dt}{dx} + t \frac{x}{1 + x^2} = \frac{1}{1 + x^2}$
 $I.F. = e^{\int \frac{x}{1 + x^2} dx} = e^{\frac{1}{2} \ln(1 + x^2)} = \sqrt{1 + x^2}$
 $t \sqrt{1 + x^2} = \int \frac{1}{\sqrt{1 + x^2}} dx$
 $\frac{\sqrt{1 + x^2}}{y} = \ln(x + \sqrt{x^2 + 1}) + c$
 $y(0) = 1 \Rightarrow c = 1$
 $\Rightarrow \sqrt{1 + x^2} = y \ln(e(x + \sqrt{x^2 + 1}))$
 $\beta = \frac{3}{\ln(e(3 + 2\sqrt{2}))} \Rightarrow \frac{3}{\beta} = \ln(e(3 + 2\sqrt{2}))$
 $e^{\frac{3}{\beta}} = e(3 + 2\sqrt{2})$
18. (3)
 Given,
 $\frac{dy}{dx} - y = 2 - e^{-x}$
 Linear differential equation,
 So, $I.F. = e^{-\int dx} = e^{-x}$
 Now solution of differential equation is given by
 $y \times IF = \int (2 - e^{-x}) \times IF dx$
 $\Rightarrow ye^{-x} = \int (2 - e^{-x}) e^{-x} dx$
 $\Rightarrow ye^{-x} = \int (2e^{-x} - e^{-2x}) dx$
 $\Rightarrow ye^{-x} = -2e^{-x} + \frac{e^{-2x}}{2} + C$
 $\Rightarrow y = -2 + \frac{e^{-x}}{2} + Ce^x$
 Given $\lim_{x \rightarrow \infty} y$ is finite
 So, $\lim_{x \rightarrow \infty} \left(-2 + \frac{e^{-x}}{2} + C \cdot e^x \right) \rightarrow \text{finite}$
 This is possible only when $C = 0$
 Hence $y = y(x) = -2 + \frac{e^{-x}}{2}$
 Now finding the slope $\frac{dy}{dx} = -\frac{1}{2}e^{-x}$
 $\frac{dy}{dx} \Big|_{x=0} = -\frac{1}{2} = m, y(0) = -2 + \frac{1}{2} = \frac{-3}{2}$
 Now equation of tangent will be,
 $y + \frac{3}{2} = -\frac{1}{2}(x - 0)$
 $\Rightarrow x + 2y = -3$
 x-intercept $a = -3$, y-intercept $b = \frac{-3}{2}$
 So, $a - 4b = -3 + 6 = 3$

19. (2) mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

$$\text{Given } 2x^2 \frac{dy}{dx} - 2xy + 3y^2 = 0$$

Dividing by $2x^2y^2$, we get,

$$\frac{1}{y^2} \frac{dy}{dx} + \frac{1}{xy} = \frac{3}{2x^2}$$

$$\text{Let } \frac{1}{y} = p, \frac{1}{y^2} \frac{dy}{dx} = \frac{dp}{dx}$$

$$\Rightarrow \frac{dp}{dx} + \frac{1}{x}(p) = \frac{3}{2x^2}$$

$$\text{General solution will be } p \cdot e^{\int \frac{1}{x} dx} = \int \frac{3}{2x^2} \cdot e^{\int \frac{1}{x} dx} dx + c$$

$$\Rightarrow \frac{x}{y} = \int \frac{3}{2x} dx + c \Rightarrow \frac{x}{y} = \frac{3}{2} \ln x + c$$

$$\therefore y(e) = \frac{e}{3} \Rightarrow c = \frac{3}{2} \Rightarrow \frac{x}{y} = \frac{3}{2} \ln x + \frac{3}{2} \text{ is the particular solution.}$$

$$\text{when } x = 1; y(1) = \frac{2}{3}$$

20. (1) mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

$$\text{Given } x(1-x^2) \frac{dy}{dx} + (3x^2y - y - 4x^3) = 0$$

$$(x-x^3) \frac{dy}{dx} + (3x^2-1)y = 4x^3$$

$$\frac{dy}{dx} + \frac{(3x^2-1)}{(x-x^3)} y = \frac{4x^3}{(x-x^3)}$$

This is a linear differential equation of the form $\frac{dy}{dx} + Py = Q$

$$\text{Here } IF = e^{\int P dx} = e^{\int \frac{3x^2-1}{x-x^3} dx}$$

$$\text{Let } x-x^3 = t \Rightarrow IF = e^{\int \frac{dt}{t}}$$

$$= e^{-\ln t} = \frac{1}{t}$$

$$\therefore IF = \frac{1}{x-x^3}$$

So the general solution of the differential equation will be

$$y \times IF = \int Q \times IF dx$$

$$y \left(\frac{1}{x-x^3} \right) = \int \frac{4x^3}{x-x^3} \times \frac{1}{(x-x^3)} dx$$

$$= \int \frac{4x^3}{(x-x^3)^2} dx$$

$$= \int \frac{4x}{(1-x^2)^2} dx$$

$$\text{Now let } I = \int \frac{4x}{(1-x^2)^2} dx$$

$$\text{Substituting } 1-x^2 = z$$

$$I = 2 \int \frac{-dz}{z^2}$$

$$= -2 \left(-\frac{1}{z} \right) + c = \frac{2}{1-x^2} + c$$

Hence the solution of the differential equation becomes

$$\frac{y}{x-x^3} = \frac{2}{1-x^2} + c$$

$$\text{At } x = 2, y = -2$$

$$\frac{-2}{2-8} = \frac{2}{1-4} + c$$

$$\frac{1}{3} = \frac{-2}{3} + c$$

$$\therefore c = 1$$

$$\text{Hence the particular solution will be } \frac{y}{x-x^3} = \frac{2}{1-x^2} + 1$$

$$\text{Put } x = 3$$

$$\frac{y}{3-27} = \frac{2}{1-9} + 1$$

$$\frac{y}{-24} = -\frac{1}{4} + 1$$

$$-\frac{y}{24} = \frac{3}{4}$$

$$y = \frac{3}{4}(-24) = -18$$

21. (I) mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

Given,

$$x \ln x \frac{dy}{dx} + y = x^2 \ln x$$

$$\Rightarrow \frac{dy}{dx} + \frac{y}{x \ln x} = x$$

This is a linear differential equation.

Now, finding integrating factor we get

$$I. F = e^{\int \frac{1}{x \ln x} dx} = e^{\ln(\ln x)} = \ln x$$

Now, solution of the differential equation is given by

$$y \times I. F = \int (x \times I. F) dx$$

$$\Rightarrow y \times \ln x = \int x \ln x \, dx$$

$$\Rightarrow y \times \ln x = \ln x \cdot \frac{x^2}{2} - \frac{x^2}{4} + c$$

Now given $y(2) = 2$, mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

$$2 \times \ln 2 = \ln 2 \cdot \frac{2^2}{2} - \frac{2^2}{4} + c$$

$$\Rightarrow c = 1$$

So, equation will be $y \times \ln x = \ln x \cdot \frac{x^2}{2} - \frac{x^2}{4} + 1$ mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

Now finding $y(e)$ we get,

$$y \times \ln e = \ln e \cdot \frac{e^2}{2} - \frac{e^2}{4} + 1$$

$$\Rightarrow y \times 1 = 1 \cdot \frac{e^2}{2} - \frac{e^2}{4} + 1$$

$$\Rightarrow y = \frac{4+e^2}{4}$$

22. (I) mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

Given differential equation can be rewritten as,

$$\frac{dy}{dx} - \frac{y}{x} = y^3(1 + \log_e x)$$

$$\frac{1}{y^3} \frac{dy}{dx} - \frac{1}{xy^2} = 1 + \log_e x$$

$$\text{Let } \left(-\frac{1}{y^2}\right) = t \Rightarrow \frac{2}{y^3} \frac{dy}{dx} = \frac{dt}{dx}$$

$$\therefore \frac{dt}{2dx} + \frac{t}{x} = (1 + \log_e x)$$

$$\Rightarrow \frac{dt}{dx} + \frac{2t}{x} = 2(1 + \log_e x) \dots\dots\dots (1)$$

We know the solution of the differential equation $\frac{dy}{dx} + Py = Q$ is given by,

$$y e^{\int P dx} = \int Q e^{\int P dx} + C \quad (\text{Where } I. F. = e^{\int P dx})$$

Therefore, on solving equation(1) we get,

$$I.F. = e^{\int \frac{2}{x} dx} = x^2$$

$$\frac{-x^2}{y^2} = \frac{2}{3} \left((1 + \log_e x) x^3 - \frac{x^3}{3} \right) + C \dots\dots\dots (2)$$

Given, $y(1) = 3$

$$\Rightarrow \frac{1}{9} = -2 \left(\frac{1}{3} - \frac{1}{9} \right) + C$$

$$\therefore C = \frac{5}{9}$$

Now putting value of C in equation(2) we get,

$$\frac{y^2}{9} = \frac{x^2}{5-2x^3(2+\log_e x^3)}$$

23. (2) Differentiate the given equation

$$\Rightarrow 2xf(x) + x^2 f'(x) - 1 = 4xf(x)$$

$$\Rightarrow x^2 \frac{dy}{dx} - 2xy = 1$$

$$\Rightarrow \frac{dy}{dx} + \left(-\frac{2}{x}\right)y = \frac{1}{x^2}$$

$$I. F. = e^{\int -\frac{2}{x} dx} = \frac{1}{x^2}$$

$$\therefore y \left(\frac{1}{x^2}\right) = \int \frac{1}{x^4} dx$$

$$\Rightarrow \frac{y}{x^2} = \frac{-1}{3x^3} + c$$

$$\Rightarrow y = -\frac{1}{3x^3} + c$$

$$\Rightarrow y = -\frac{1}{3x} + cx^2$$

$$\therefore f(1) = \frac{2}{3} = -\frac{1}{3} + c \Rightarrow c = 1$$

$$f(x) = -\frac{1}{3x} + x^2$$

$$18f(3) = 160$$

24. (1)

Let

$$\int_0^2 f(t) dt = k$$

$$y = f(x)$$

Now, we have

$$f(x) + f'(x) = \int_0^2 f(t) dt$$

$$\Rightarrow \frac{dy}{dx} + y = k$$

This is a linear differential equation.

$$I. F. = e^{\int dx} = e^x$$

Solution of the given differential equation is

$$ye^x = k \int e^x dx$$

$$\Rightarrow ye^x = ke^x + C$$

Now, $f(0) = e^{-2}$, so

$$C = e^{-2} - k$$

Hence,

$$ye^x = ke^x + (e^{-2} - k)$$

$$\Rightarrow y = f(x) = k + (e^{-2} - k)e^{-x} \dots (i)$$

Now,

$$k = \int_0^2 f(t) dt$$

$$\Rightarrow k = \int_0^2 [k + (e^{-2} - k)e^{-t}] dt$$

$$\Rightarrow k = [kt - (e^{-2} - k)e^{-t}]_0^2$$

$$\Rightarrow k = 2k - (e^{-2} - k)(e^{-2} - 1)$$

$$\Rightarrow (e^{-2} - k)(e^{-2} - 1) = k$$

$$\Rightarrow k = (e^{-2} - 1)$$

So,

$$f(x) = (e^{-2} - 1) + e^{-x}$$

$$\therefore f(2) = 2e^{-2} - 1$$

$$f(0) = e^{-2}$$

So,

$$2f(0) - f(2) = 1$$

25. (3)

Given

$$\frac{dy}{dx} + e^x(x^2 - 2)y = (x^2 - 2x)(x^2 - 2)e^{2x}$$

It is linear differential equation so,

$$IF = e^{\int (x^2 - 2)e^x dx}$$

$$= e^{x^2 e^x - 2 \int x e^x dx - 2e^x} = e^{x^2 e^x - 2[xe^x - e^x] - 2e^x}$$

$$IF = e^{(x^2 - 2x)e^x}$$

Now solution is given by

$$y \times e^{(x^2 - 2x)e^x} = \int e^{(x^2 - 2x)e^x} \times (x^2 - 2x)(x^2 - 2)e^{2x} dx$$

$$= \int e^{(x^2 - 2x)e^x} \times (x^2 - 2x)e^x (x^2 - 2)e^x dx$$

$$\text{Now let } (x^2 - 2x)e^x = t$$

$$\text{We get } e^x(x^2 - 2x) + e^x(2x - 2)dx = dt$$

$$e^x(x^2 - 2x + 2x - 2)dx = dt$$

$$e^x(x^2 - 2)dx = dt$$

$$y \times e^{(x^2 - 2x)e^x} = \int e^t \times t \times dt$$

$$y \times e^{(x^2 - 2x)e^x} = e^t(t - 1) + c$$

$$y \times e^{(x^2 - 2x)e^x} = e^{(x^2 - 2x)e^x}((x^2 - 2x)e^x - 1) + c$$

$$\text{Now at } x = 0, y = 0$$

$$0 \times e^0 = e^0(0xe^x - 1) + c$$

$$0 = -1 + c \quad c = 1$$

Putting the value of $c = 1$ we get the equation as

$$\Rightarrow y \times e^{(x^2 - 2x)e^x} = e^{(x^2 - 2x)e^x}((x^2 - 2x)e^x - 1) + 1$$

Now putting $x = 2$ in equation we get

$$y \times e^{(4 - 4)e^2} = e^{(4 - 4)e^2}((4 - 4)e^2 - 1) + 1$$

$$\Rightarrow y \times e^0 = e^0((0)e^2 - 1) + 1$$

$$\Rightarrow y = -1 + 1 = 0$$

26. (1)

$$\text{We have, } xdy = (y + x^3 \cos x)dx$$

$$\Rightarrow xdy = ydx + x^3 \cos x dx$$

$$\Rightarrow \frac{xdy - ydx}{x^2} = \frac{x^3 \cos x dx}{x^2}$$

$$\Rightarrow \frac{d}{dx}\left(\frac{y}{x}\right) = \int x \cos x dx$$

$$\Rightarrow \frac{y}{x} = x \sin x - \int 1 \cdot \sin x dx$$

$$\text{Therefore, } \frac{y}{x} = x \sin x + \cos x + C$$

$$\text{At } x = \pi, y = 0,$$

$$0 = -1 + C$$

$$\Rightarrow C = 1, x = \pi, y = 0$$

$$\text{So, } \frac{y}{x} = x \sin x + \cos x + 1$$

$$\Rightarrow y = x^2 \sin x + x \cos x + x$$

$$\text{Hence, } y\left(\frac{\pi}{2}\right) = \frac{\pi^2}{4} + \frac{\pi}{2}.$$

27. (2)

$$(x - x^3)dy = (y + yx^2 - 3x^4)dx$$

$$\Rightarrow xdy - ydx = (yx^2 - 3x^4)dx + x^3 dy$$

$$\Rightarrow \frac{xdy - ydx}{x^2} = (ydx + xdy) - 3x^2 dx$$

$$\Rightarrow d\left(\frac{y}{x}\right) = d(xy) - d(x^3)$$

Integrate

$$\Rightarrow \frac{y}{x} = xy - x^3 + c$$

$$\text{given } f(3) = 3$$

$$\Rightarrow \frac{3}{3} = 3 \times 3 - 3^3 + c$$

$$\Rightarrow c = 19$$

$$\therefore \frac{y}{x} = xy - x^3 + 19$$

$$\text{at } x = 4, \frac{y}{4} = 4y - 64 + 19$$

$$15y = 4 \times 45$$

$$\Rightarrow y = 12$$

28. (I) mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

Given differential equation, $\sin(2x^2)\ln(\tan x^2)dy + (4xy - 4\sqrt{2}x \sin(x^2 - \frac{\pi}{4}))dx = 0$

$$\Rightarrow \ln(\tan x^2)dy + \frac{4xydx}{\sin(2x^2)} - \frac{4\sqrt{2}x \sin(x^2 - \frac{\pi}{4})}{\sin(2x^2)}dx = 0$$

$$\Rightarrow d(y \cdot \ln(\tan x^2)) - 4\sqrt{2}x \frac{(\sin x^2 - \cos x^2)}{\sqrt{2}(2 \sin x^2 \cos x^2)}dx = 0$$

$$\Rightarrow d(y \ln(\tan x^2)) - \frac{4x(\sin x^2 - \cos x^2)}{(\sin x^2 + \cos^2 x)^2 - 1}dx = 0$$

Now integrating both side,

$$\Rightarrow \int d(y \ln(\tan x^2)) - \int \frac{4x(\sin x^2 - \cos x^2)}{(\sin x^2 + \cos^2 x)^2 - 1}dx = \int 0$$

Now let $\sin x^2 + \cos^2 x = t \Rightarrow -2x(\sin x^2 - \cos^2 x) = dt$

$$\Rightarrow \int d(y \ln(\tan x^2)) + 2 \int \frac{dt}{t^2 - 1} = \int 0$$

$$\Rightarrow y \ln(\tan x^2) + 2 \cdot \frac{1}{2} \ln \left| \frac{t-1}{t+1} \right| = c$$

$$y \ln(\tan x^2) + \ln \left(\frac{\sin x^2 + \cos x^2 - 1}{\sin x^2 + \cos x^2 + 1} \right) = c$$

$$\text{Put } y = 1 \text{ and } x = \sqrt{\frac{\pi}{6}}$$

$$1 \ln \left(\frac{1}{\sqrt{3}} \right) + \ln \left(\frac{\left(\frac{1}{2} + \frac{\sqrt{3}}{2} - 1 \right)}{\left(\frac{1}{2} + \frac{\sqrt{3}}{2} + 1 \right)} \right) = c$$

$$\text{Now at } x = \sqrt{\frac{\pi}{3}}$$

$$\Rightarrow y \left(\ln \sqrt{3} \right) + \ln \left(\frac{\left(\frac{\sqrt{3}}{2} + \frac{1}{2} - 1 \right)}{\left(\frac{\sqrt{3}}{2} + \frac{1}{2} + 1 \right)} \right) = \ln \left(\frac{1}{\sqrt{3}} \right) + \ln \left(\frac{\sqrt{3}-1}{\sqrt{3}+3} \right)$$

$$\Rightarrow y \left(\ln \sqrt{3} \right) = \ln \left(\frac{1}{\sqrt{3}} \right)$$

$$\Rightarrow y = -1$$

$$\text{So, } |y| = 1$$

29. (4) mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

Given,

$$2ye^{\frac{x}{y^2}}dx + (y^2 - 4xe^{\frac{x}{y^2}})dy = 0$$

$$2e^{\frac{x}{y^2}}[ydx - 2xdy] + y^2dy = 0$$

$$2e^{\frac{x}{y^2}} \left[\frac{y^2dx - x \cdot (2y)dy}{y} \right] + y^2dy = 0$$

Divide by y^3

$$2e^{\frac{x}{y^2}} \left[\frac{y^2dx - x \cdot (2y)dy}{y^4} \right] + \frac{1}{y}dy = 0$$

$$2e^{\frac{x}{y^2}} d \left(\frac{x}{y^2} \right) + \frac{1}{y}dy = 0$$

Now integrating both side we get,

$$\int 2e^{\frac{x}{y^2}} d \left(\frac{x}{y^2} \right) + \int \frac{1}{y}dy = 0$$

$$2e^{\frac{x}{y^2}} + \ln y + c = 0$$

Given, (0, 1) lies on it,

$$\text{So, } 2e^0 + \ln 1 + c = 0 \Rightarrow c = -2$$

$$\text{Hence required curve: } 2e^{\frac{x}{y^2}} + \ln y - 2 = 0$$

For $x(e)$

$$2e^{\frac{x}{e^2}} + \ln e - 2 = 0 \Rightarrow x = -e^2 \log_e 2$$

mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo mathongo

30. (4)

The given differential equation is $\left((x+2)e^{\left(\frac{y+1}{x+2}\right)} + (y+1)\right)dx = (x+2)dy$, $y(1)=1$

Let, $y+1=Y \Rightarrow dy=dY$ and $x+2=X \Rightarrow dx=dX$ and at $y=1$, $Y=2$ and at $x=1$, $X=3$.

Thus, the differential equation becomes $\left(Xe^{\frac{Y}{X}} + Y\right)dX = XdY$

$$\Rightarrow XdY - YdX = Xe^{\frac{Y}{X}}dX$$

$$\Rightarrow \frac{(XdY - YdX)}{X^2} = \left(\frac{Xe^{\frac{Y}{X}}}{X^2}\right)dX$$

$$\Rightarrow \frac{d}{dX}\left(\frac{Y}{X}\right) = \left(\frac{Xe^{\frac{Y}{X}}}{X^2}\right)dX$$

$$\Rightarrow d\left(\frac{Y}{X}\right)e^{-\frac{Y}{X}} = \frac{dX}{X}$$

Integrating both sides w.r.to X , we get $\int d\left(\frac{Y}{X}\right)e^{-\frac{Y}{X}} = \int \frac{dX}{X}$

$$\Rightarrow -e^{-\frac{Y}{X}} = \log_e |X| + c$$

Given, at $X=3$, $Y=2$

$$\Rightarrow -e^{-\frac{2}{3}} = \log_e |3| + c$$

$$\Rightarrow c = -e^{-\frac{2}{3}} - \log_e |3|$$

$$\Rightarrow -e^{-\frac{Y}{X}} = \log_e |X| - e^{-\frac{2}{3}} - \log_e 3$$

$$e^{-\frac{Y}{X}} = e^{\frac{2}{3}} + \log_e 3 - \log_e |X| > 0$$

$$\log_e |X| < e^{\frac{2}{3}} + \log_e 3$$

Let, $\lambda = \left(e^{\frac{2}{3}} + \log_e 3\right)$ then, we have $|x+2| < e^\lambda$

$$\Rightarrow -e^\lambda < x+2 < e^\lambda$$

$$\Rightarrow -e^\lambda - 2 < x < e^\lambda - 2$$

Thus, the domain of the function is $(-e^\lambda - 2, e^\lambda - 2)$.

Given, the domain of the function is (α, β) , hence $\alpha = -e^\lambda - 2$ & $\beta = e^\lambda - 2$

$$\Rightarrow \alpha + \beta = -e^\lambda - 2 + e^\lambda - 2 = -4$$

$$\Rightarrow |\alpha + \beta| = 4.$$