



Mock Tests for NTA

JEE MAIN

2021 | 5 in Book
& 5 Online

Corporate Office

DISHA PUBLICATION

45, 2nd Floor, Maharishi Dayanand Marg, Corner Market,
Malviya Nagar, New Delhi - 110017
Tel : 49842349 / 49842350

© Copyright Author

No part of this publication may be reproduced in any form without prior permission of the publisher. The author and the publisher do not take any legal responsibility for any errors or misrepresentations that might have crept in. We have tried and made our best efforts to provide accurate up-to-date information in this book.

All Right Reserved

Get free access to Online Test(s)?

INSTRUCTIONS

1. You can access your test on any Window based Desktop, android tablets or ipads and mobile phones absolutely free.
2. Visit the given link or scan the QR code.
3. Click on "Attempt Free Mock Tests", a Registration window pops up, enter all the details in the form & click "Sign UP".
4. User is now logged in the account & all the Mock Tests appears in the grid. User can attempt the Free Mock Test(s) by clicking the "Start" button.
5. Contact us at support@mylearninggraph.com for any support.

5 Mock Tests for
JEE Main
Exam 2021



12 Chapter wise Tests for
NTA JEE Main Exam

Physics - 4 Tests | Chemistry - 4 Tests



<http://bit.ly/5MocktestJEE>

<http://bit.ly/12MockTestJEE>

Typeset by Disha DTP Team



www.dishapublication.com

Books &
ebooks for
School &
Competitive
Exams



www.mylearninggraph.com

Etests
for
Competitive
Exams



INDEX

- **Mock Test-1** **1 – 12**
- **Mock Test-2** **13 – 24**
- **Mock Test-3** **25 – 34**
- **Mock Test-4** **35 – 46**
- **Mock Test-5** **47 – 60**
- **Solutions Mock Test-1** **61 – 82**
- **Solutions Mock Test-2** **83 – 98**
- **Solutions Mock Test-3** **99 – 114**
- **Solutions Mock Test-4** **115 – 137**
- **Solutions Mock Test-5** **138 – 160**

MOCK TEST

1

INSTRUCTIONS

1. This test will be a 3 hours Test.
2. This test consists of Physics, Chemistry and Mathematics questions with equal weightage of 100 marks.
3. Each question is of 4 marks.
4. There are three sections in the question paper consisting of Physics (Q.no.1 to 25), Chemistry (Q.no.26 to 50) and Mathematics (Q. no.51 to 75). Each section is divided into two parts, Part I consists of 20 multiple choice questions & Part II consists of 5 Numerical value type Questions.
5. There will be only one correct choice in the given four choices in Part I. For each question 4 marks will be awarded for correct choice, 1 mark will be deducted for incorrect choice for Part I Questions and zero mark will be awarded for not attempted question. For Part II Questions 4 marks will be awarded for correct answer and zero for unattempted and incorrect answer.
6. Any textual, printed or written material, mobile phones, calculator etc. is not allowed for the students appearing for the test.
7. All calculations / written work should be done in the rough sheet provided.

PHYSICS

PART-I (Multiple Choice Questions)

1. Two stars each of mass M and radius R are approaching each other for a head-on collision. They start approaching each other when their separation is $r \gg R$. If their speeds at this separation are negligible, the speed v with which they collide would be

$$(a) \quad v = \sqrt{GM\left(\frac{1}{R} - \frac{1}{r}\right)}$$


$$(b) \quad v = \sqrt{GM\left(\frac{1}{2R} - \frac{1}{r}\right)}$$

$$(c) \quad v = \sqrt{GM\left(\frac{1}{R} + \frac{1}{r}\right)}$$

$$(d) \quad v = \sqrt{GM\left(\frac{1}{2R} + \frac{1}{r}\right)}$$

2. A block of mass M is kept on a platform which is accelerated upward with a constant acceleration ' a ' during the time interval T . The work done by

normal reaction between the block and platform is



(a) $-\frac{MgaT^2}{2}$

(b) $\frac{1}{2}M(g+a)aT^2$

(c) $\frac{1}{2}Ma^2T$

(d) Zero

3. A large number of water drops each of radius r combine to have a drop of radius R . If the surface tension is T and the mechanical equivalent of heat is J , then the rise in temperature will be

(a) $\frac{2T}{rJ}$


(b) $\frac{3T}{RJ}$

(c) $\frac{3T}{J}\left(\frac{1}{r}-\frac{1}{R}\right)$

(d) $\frac{2T}{J}\left(\frac{1}{r}-\frac{1}{R}\right)$

4. Three charges are placed at the vertices of an equilateral triangle of side ' a ' as shown in the following figure. The force experienced by the charge placed

at the vertex A in a direction normal to BC is



(a) $Q^2/(4\pi\epsilon_0 a^2)$

(b) $-Q^2/(4\pi\epsilon_0 a^2)$

(c) Zero

(d) $Q^2/(2\pi\epsilon_0 a^2)$

5. Axis of a solid cylinder of infinite length and radius R lies along y -axis, it carries a uniformly distributed current i along $+y$ direction. Magnetic field at a point

$$\left(\frac{R}{2}, y, \frac{R}{2}\right)$$

(a) $\frac{\mu_0 i}{4\pi R}(\hat{i} - \hat{k})$

(b) $\frac{\mu_0 i}{2\pi R}(\hat{j} - \hat{k})$

(c) $\frac{\mu_0 i}{4\pi R} \hat{j}$

(d) $\frac{\mu_0 i}{4\pi R}(\hat{i} + \hat{k})$

6. Two identical short bar magnets, each having magnetic moment of 10 Am^2 , are arranged such that their axial lines are perpendicular to each other and their centres be along the same straight line in a horizontal plane. If the distance between their centres is 0.2 m , the

resultant magnetic induction at a point midway between them is

$$(\mu_0 = 4\pi \times 10^{-7} \text{ Hm}^{-1})$$

(a) $\sqrt{2} \times 10^{-7}$ tesla

(b) $\sqrt{5} \times 10^{-7}$ tesla

(c) $\sqrt{2} \times 10^{-3}$ tesla

(d) $\sqrt{5} \times 10^{-3}$ tesla

7. Two boys are standing at the ends A and B of a ground where $AB = a$. The boy at B starts running in a direction perpendicular to AB with velocity v_1 . The boy at A starts running simultaneously with velocity v and catches the other boy in a time t , where t is





(a) $a / \sqrt{v^2 + v_1^2}$

(b) $a / (v + v_1)$

(c) $a / (v - v_1)$

(d) $\sqrt{a^2 / (v^2 - v_1^2)}$

8. A block is placed on a rough horizontal plane. A time dependent horizontal force $F = kt$ acts on the block. Here, k is a positive constant. The acceleration-time graph of the block is



9. A new system of units is proposed in which unit of mass is α kg, unit of length is β m and unit of time is γ s. What will be value of 5 J in this new system?

(a) $5\alpha\beta^2\gamma^{-2}$

(b) $5\alpha^{-1}\beta^{-2}\gamma^2$

(c) $5\alpha^{-2}\beta^{-1}\gamma^{-2}$

(d) $5\alpha^{-1}\beta^2\gamma^2$

10. Television signals on earth cannot be received at distances greater than 100 km from the transmission station. The reason behind this is that

(a) the receiver antenna is unable to detect the signal at a distance greater than 100 km

(b) the TV programme consists of both audio and video signals

(c) the TV signals are less powerful than radio signals


(d) the surface of earth is curved like a sphere

11. A sinusoidal voltage of amplitude 25 volt and frequency 50Hz is applied to a half wave rectifier using P-n junction diode. No filter is used and the load resistor is 1000Ω . The forward resistance R_f of ideal diode is 10Ω . The percentage rectifier efficiency is
 (a) 40% (b) 20%
 (c) 30% (d) 15%

12. When photon of energy 4.25 eV strike the surface of a metal A, the ejected photoelectrons have maximum kinetic energy $T_A\text{ eV}$ and de-Brolie wavelength λ_A . The maximum kinetic energy of photoelectrons liberated from another metal B by photon of energy 4.70 eV is $T_B = (T_A - 1.50)\text{ eV}$. If the de-Brolie wavelength of these photoelectrons is $\lambda_B = 2\lambda_A$, then
 (a) the work function of A is 3.40 eV
 (b) the work function of B is 6.75 eV
 (c) $T_A = 2.00\text{ eV}$
 (d) $T_B = 2.75\text{ eV}$


13. Given is the graph between $\frac{PV}{T}$ and P for 1 g of oxygen gas at two different temperatures T_1 and T_2 , as shown in figure. Given, density of oxygen = 1.427 kg m^{-3} . The value of PV/T at the point A and the relation between T_1 and T_2 are respectively

$$\frac{PV}{T}\text{ J k}^{-1}$$




- (a) 0.259 J K^{-1} and $T_1 < T_2$
 (b) $8.314\text{ J mol}^{-1}\text{ K}^{-1}$ and $T_1 > T_2$
 (c) 0.259 J K^{-1} and $T_1 > T_2$
 (d) 4.28 g JK^{-1} and $T_1 < T_2$
14. An observer moves towards a stationary source of sound with a speed $1/5$ th of the speed of sound. The wavelength and frequency of the sound emitted are λ and f respectively. The apparent frequency and wavelength recorded by the observer are respectively.
 (a) $0.8f, 0.8\lambda$ (b) $1.2f, 1.2\lambda$
 (c) $1.2f, \lambda$ (d) $f, 1.2\lambda$
15. The figure shows a system of two concentric spheres of radii r_1 and r_2 are kept at temperatures T_1 and T_2 , respectively. The radial rate of flow of heat in a substance between the two concentric spheres is proportional to

- (a) $\ln\left(\frac{r_2}{r_1}\right)$
 (b) $\frac{(r_2 - r_1)}{(r_1 r_2)}$
 (c) $(r_2 - r_1)$
 (d) $\frac{r_1 r_2}{(r_2 - r_1)}$



16. A gas is compressed isothermally to half its initial volume. The same gas is compressed separately through an adiabatic process until its volume is again reduced to half. Then :
- Compressing the gas isothermally will require more work to be done.
 - Compressing the gas through adiabatic process will require more work to be done.
 - Compressing the gas isothermally or adiabatically will require the same amount of work.
 - Which of the case (whether compression through isothermal or through adiabatic process) requires more work will depend upon the atomicity of the gas.
17. A ray PQ incident on the refracting face BA is refracted in the prism BAC as shown in the figure and emerges from the other



refracting face AC as RS such that $AQ = AR$. If the angle of prism $A = 60^\circ$ and the refractive index of the material of prism is $\sqrt{3}$, then the angle of deviation of the ray is

- 60°
- 45°
- 30°
- None of these


18. Which of the following has/have zero average value in a plane electromagnetic wave ?
- Both magnetic and electric field
 - Electric field only
 - Magnetic energy
 - Electric energy
19. Two inductors L_1 (inductance 1 mH, internal resistance 3Ω) and L_2 (inductance 2 mH, internal resistance 4Ω), and a resistor R (resistance 12Ω) are all connected in parallel across a 5V battery. The circuit is switched on at time $t = 0$. The ratio of the maximum to the minimum current (I_{\max}/I_{\min}) drawn from the battery is
- 8
 - 10
 - 12
 - 14
20. In a diffraction pattern due to a single slit of width 'a', the first minimum is observed at an angle 30° when light of wavelength 5000 \AA is incident on the slit. The first secondary maximum is observed at an angle of :
- $\sin^{-1}\left(\frac{1}{4}\right)$
 - $\sin^{-1}\left(\frac{2}{3}\right)$
 - $\sin^{-1}\left(\frac{1}{2}\right)$
 - $\sin^{-1}\left(\frac{3}{4}\right)$

PART-II (Numerical Answer Questions)

21. Figure shows use of potentiometer for comparison of two resistances. The balance point with standard resistance $R = 10\Omega$ is at 58.3 cm, while that with unknown resistance X is 68.5 cm. Find X (in Ω).

CHEMISTRY

PART-I (Multiple Choice Questions)



- 22.** An automobile moves on a road with a speed of 54 km h^{-1} . The radius of its wheels is 0.45 m and the moment of inertia of the wheel about its axis of rotation is 3 kg m^2 . If the vehicle is brought to rest in 15s , the magnitude of average torque (in $\text{kgm}^2\text{s}^{-2}$) transmitted by its brakes to the wheel is :
- 23.** A coil of effective area 4 m^2 is placed at right angles to the magnetic induction B . The e.m.f. of 0.32 V is induced in the coil. When the field is reduced to 20% of its initial value in 0.5 sec . Find B (in wb/m^2).
- 24.** A disc of radius $R = 10 \text{ cm}$ oscillates as a physical pendulum about an axis perpendicular to the plane of the disc at a distance r from its centre. If $r = \frac{R}{4}$, the approximate period of oscillation (in second) is (Take $g = 10 \text{ m s}^{-2}$)
- 25.** Taking the wavelength of first Balmer line in hydrogen spectrum ($n = 3$ to $n = 2$) as 660 nm , the wavelength (in nm) of the 2^{nd} Balmer line ($n = 4$ to $n = 2$) will be;
- 26.** Which of the following has the highest $p\pi - p\pi$ bonding tendency?
- N
 - P
 - As
 - Sb
- 27.** Among the following, the compound that is both paramagnetic and coloured, is
- KMnO_4
 - CuF_2
 - $\text{K}_2\text{Cr}_2\text{O}_7$
 - All are coloured
- 28.** The bond angle between two hybrid orbitals is 105° . The percentage of s -character of hybrid orbital is between
- 50-55%
 - 9-12%
 - 21-23%
 - 11-12%
- 29.** Identify Z in the following sequence of reactions –
- $$\text{CH}_3 - \text{COONH}_4 \xrightarrow{\Delta} \text{X} \xrightarrow{\text{P}_2\text{O}_5} \text{Y}$$
- $$\xrightarrow{\text{H}_2\text{O}/\text{H}^+} \text{Z}$$
- $\text{CH}_3 - \text{CH}_2 - \text{CO} - \text{NH}_2$
 - $\text{CH}_3 - \text{CN}$
 - $(\text{CH}_3\text{CO})_2\text{O}$
 - $\text{CH}_3 - \text{COOH}$
- 30.** Correct order of first IP among following elements Be, B, C, N, O is
- $\text{B} < \text{Be} < \text{C} < \text{O} < \text{N}$
 - $\text{B} < \text{Be} < \text{C} < \text{N} < \text{O}$
 - $\text{Be} < \text{B} < \text{C} < \text{N} < \text{O}$
 - $\text{Be} < \text{B} < \text{C} < \text{O} < \text{N}$

- 31.** Select the rate law that corresponds to data shown for the following reaction
- $$A + B \longrightarrow \text{products.}$$
- | Exp. | [A] | [B] | Initial rate |
|-------------|------------|------------|---------------------|
| 1 | 0.012 | 0.035 | 0.1 |
| 2 | 0.024 | 0.070 | 0.8 |
| 3 | 0.024 | 0.035 | 0.1 |
| 4 | 0.012 | 0.070 | 0.8 |
- (a) $\text{rate} = k [B]^3$
 (b) $\text{rate} = k [B]^4$
 (c) $\text{rate} = k [A] [B]^3$
 (d) $\text{rate} = k [A]^2 [B]^2$
- 32.** The pH of 0.1 M solution of the following species increases in the order :
- (a) $\text{NaCl} < \text{NH}_4\text{Cl} < \text{NaCN} < \text{HCl}$
 (b) $\text{HCl} < \text{NH}_4\text{Cl} < \text{NaCl} < \text{NaCN}$
 (c) $\text{NaCN} < \text{NH}_4\text{Cl} < \text{NaCl} < \text{HCl}$
 (d) $\text{HCl} < \text{NaCl} < \text{NaCN} < \text{NH}_4\text{Cl}$
- 33.** Aldehydes and ketones are distinguished by which of the following test ?
- (a) Lucas test
 (b) Tollen's test
 (c) KMnO_4 solution (Baeyer's test)
 (d) None of these
- 34.** Which is not a true statement?
- (a) α -Carbon of α -amino acid is asymmetric
 (b) All proteins are found in L-form
 (c) Human body can synthesize all proteins they need
- 35.** At pH = 7 both amino and carboxylic groups exist in ionised form
- 36.** Which of the following products are obtained when Na_2CO_3 is added to a solution of copper sulphate ?
- (a) Basic copper carbonate $[\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2]$, sodium sulphate and CO_2 .
 (b) Copper hydroxide, sodium sulphate and CO_2 .
 (c) Copper carbonate, sodium sulphate and CO_2 .
 (d) Copper carbonate and sodium sulphate.
- 37.** Which of the following statement is incorrect with respect to metallic or electronic conductivity?
- (a) Metallic conductivity depends on the structure of metal and its characteristics.
 (b) Metallic conductivity depends on the number of electrons in the valence shell of atom of metal.
 (c) The electrical conductivity of metal increases with increase in temperature.
 (d) There is no change in the structure of metal during electrical conduction.
- 38.** A solid has a structure in which 'W' atoms are located at the corners of a cubic lattice, 'O' atoms at the centre of edges and Na atoms at the centre of the cube. The formula for the compound is
- (a) Na_2WO_3 (b) Na_2WO_2
 (c) NaWO_2 (d) NaWO_3

38. When phenol is treated with excess bromine water. It gives
 (a) *m*-Bromophenol
 (b) *o*-and *p*-Bromophenols
 (c) 2,4-Dibromophenol
 (d) 2,4, 6-Tribromophenol.
39. Given below, catalyst and corresponding process/reaction are matched. The one with mismatch is
 (a) $[\text{RhCl}(\text{PPh}_3)_2]$: Hydrogenation
 (b) $\text{TiCl}_4 + \text{Al}(\text{C}_2\text{H}_5)_3$: Polymerization
 (c) V_2O_5 : Haber-Bosch process
 (d) Nickel : Hydrogenation
40. The molecule which has zero dipole moment is :
 (a) CH_3Cl (b) NF_3
 (c) BF_3 (d) ClO_2^-
41. One mole of NaCl (*s*) on melting absorbed 30.5 kJ one of heat and its entropy is increased by $28.8 \text{ J K}^{-1}\text{mol}^{-1}$. The melting point of NaCl is
 (a) 1059 K (b) 30.5 K
 (c) 28.8 K (d) 28800 K
42. Which alkene on ozonolysis gives $\text{CH}_3\text{CH}_2\text{CHO}$ and CH_3CCH_3 ?

$$\begin{array}{c} \text{CH}_3 \\ || \\ \text{O} \\ \diagup \quad \diagdown \\ \text{CH}_3 \quad \text{CH}_3 \end{array}$$

 (a) $\text{CH}_3\text{CH}_2\text{CH}=\text{C}(\text{CH}_3)_2$
 (b) $\text{CH}_3\text{CH}_2\text{CH}=\text{CHCH}_2\text{CH}_3$
 (c) $\text{CH}_3\text{CH}_2\text{CH}=\text{CHCH}_3$
 (d) $\text{CH}_3 - \text{C}(\text{CH}_3) = \text{CHCH}_3$
43. On reduction of KMnO_4 by oxalic acid in acidic medium, the oxidation number of Mn changes. What is the magnitude of this change?
 (a) From 7 to 2 (b) From 6 to 2
 (c) From 5 to 2 (d) From 7 to 4
44. The half-life for radioactive decay of C-14 is 5730 years. An archaeological artifact containing wood had only 80% of the C-14 found in a living tree. The age of the sample is
 (a) 1485 years (b) 1845 years
 (c) 530 years (d) 4767 years.
45. Which one of the following complexes is an outer orbital complex ?
 (a) $[\text{Co}(\text{NH}_3)_6]^{3+}$
 (b) $[\text{Mn}(\text{CN})_6]^{4-}$
 (c) $[\text{Fe}(\text{CN})_6]^{4-}$
 (d) $[\text{Ni}(\text{NH}_3)_6]^{2+}$
 (Atomic nos. : Mn = 25; Fe = 26; Co=27, Ni=28)
- PART-II (Numerical Answer Questions)**
46. If pressure of a gas is reduced by 25%, then what should be the temperature required to make its volume twice at NTP?
47. An aromatic compound of formula $\text{C}_7\text{H}_7\text{Cl}$ has in all isomers :
48. Calculate the volume strength of 1.5 NH_2O_2 solution.
49. A metal crystallizes into a lattice containing a sequence of layers of atoms of ABABAB.....What percentage by volume of this lattice has empty space?
50. In an experiment, 4 g of M_2O_x oxide was reduced to 2.8 g of the metal. Calculate the number of O atoms in the oxide.
 (Given : Atomic mass of the metal = 56 g mol^{-1})

MATHEMATICS**PART-I (Multiple Choice Questions)**

51. If the coefficient of 4th term in the expansion of $\left(x + \frac{\alpha}{2x}\right)^n$ is 20, then

the respective values of α and n are

- (a) 2, 7 (b) 5, 8
 (c) 3, 6 (d) 2, 6

52. If the roots of the quadratic equation $x^2 + px + q = 0$ are $\tan 30^\circ$ and $\tan 15^\circ$, respectively, then the value of $2 + q - p$ is
 (a) 2 (b) 3
 (c) 0 (d) 1
53. If a^2, b^2, c^2 are in A.P. then

$\frac{1}{b+c}, \frac{1}{c+a}, \frac{1}{a+b}$ are in-

- (a) A.P.
 (b) GP.
 (c) H.P.
 (d) None of these

54. Let C be the circle with centre $(0, 0)$ and radius 3 units. The equation of the locus of the mid points of the chords of the circle C that subtend an angle of $\frac{2\pi}{3}$ at its center is

- (a) $x^2 + y^2 = \frac{3}{2}$
 (b) $x^2 + y^2 = 1$

- (c) $x^2 + y^2 = \frac{27}{4}$

- (d) $x^2 + y^2 = \frac{9}{4}$

55. If $y = \tan^{-1} \left(\frac{\log_e(e/x^2)}{\log_e(ex^2)} \right)$

$$+ \tan^{-1} \left(\frac{3 + 2 \log_e x}{1 - 6 \log_e x} \right),$$

then $\frac{d^2y}{dx^2}$ is

- (a) 2 (b) 1
 (c) 0 (d) -1

56. If $\{ \}$ denotes the fractional part of x , the range of the function

$$f(x) = \sqrt{\{x\}^2 - 2\{x\}}$$

- (a) ϕ (b) $[0, 1/2]$
 (c) $\{0, 1/2\}$ (d) $\{0\}$

57. The length of the perpendicular from the origin to a line is 7 and line makes an angle of 150° with the positive direction of y -axis, then the equation of the line is

- (a) $\sqrt{3}x + y = 7$
 (b) $\sqrt{3}x - y = 14$
 (c) $\sqrt{3}x + y + 14 = 0$
 (d) $\sqrt{3}x + y - 14 = 0$

58. $\int \frac{dx}{\cos x + \sqrt{3} \sin x}$ equals

- (a) $\log \tan \left(\frac{x}{2} + \frac{\pi}{12} \right) + C$

- (b) $\log \tan \left(\frac{x}{2} - \frac{\pi}{12} \right) + C$

- (c) $\frac{1}{2} \log \tan \left(\frac{x}{2} + \frac{\pi}{12} \right) + C$

- (d) $\frac{1}{2} \log \tan \left(\frac{x}{2} - \frac{\pi}{12} \right) + C$

59. If $\frac{\tan 3\theta - 1}{\tan 3\theta + 1} = \sqrt{3}$, then the general value of θ is

- (a) $\frac{n\pi}{3} - \frac{\pi}{12}$ (b) $n\pi + \frac{7\pi}{12}$
 (c) $\frac{n\pi}{3} + \frac{7\pi}{36}$ (d) $n\pi + \frac{\pi}{12}$

60. Three normals are drawn to the parabola $y^2 = x$ through point $(a, 0)$. Then

- (a) $a = 1/2$
 (b) $a = 1/4$
 (c) $a > 1/2$
 (d) None of these

61. If four vertices of a regular octagon are chosen at random, then the probability that the quadrilateral formed by them is a rectangle is

- (a) $\frac{1}{8}$ (b) $\frac{2}{21}$
 (c) $\frac{1}{32}$ (d) $\frac{1}{35}$

62. The function

$f(x) = x^3 - 3x^2 - 24x + 5$ is an increasing function in the interval given below

- (a) $(-\infty, -2) \cup (4, \infty)$
 (b) $(-2, \infty)$
 (c) $(-2, 4)$
 (d) $(-\infty, 4)$

63. If $y = y(x)$ and it follows the relation $x \cos y + y \cos x = \pi$ then $y''(0) =$

- (a) 1 (b) -1
 (c) π (d) $-\pi$

64. ABC is triangular park with $AB = AC = 100$ m. A clock tower is situated at the mid-point of BC. The angles of elevation of the top of the tower at A and B are $\cot^{-1} 3.2$ and $\operatorname{cosec}^{-1} 2.6$ respectively. The height of the tower is

- (a) 50 m
 (b) 25 m
 (c) 40 m
 (d) None of these

65. If the vectors $\overrightarrow{AB} = -3\hat{i} + 4\hat{k}$ and

$\overrightarrow{AC} = 5\hat{i} - 2\hat{j} + 4\hat{k}$ are the sides of a triangle ABC, then the length of the median through A is

- (a) $\sqrt{14}$ (b) $\sqrt{18}$
 (c) $\sqrt{29}$ (d) 4

66. The negation of the compound proposition $p \vee (\sim p \vee q)$ is

- (a) $(p \wedge \sim q) \wedge \sim p$
 (b) $(p \wedge \sim q) \vee \sim p$
 (c) $(p \vee \sim q) \vee \sim p$
 (d) None of these

67. Let $f(x) = \begin{cases} x^p \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$ then

$f(x)$ is continuous but not differentiable at $x = 0$ if

- (a) $0 < p \leq 1$
- (b) $1 \leq p < \infty$
- (c) $-\infty < p < 0$
- (d) $p = 0$

68. The length and foot of the perpendicular from the point $(7, 14, 5)$ to the plane $2x + 4y - z = 2$, are

- (a) $\sqrt{21}, (1, 2, 8)$
- (b) $3\sqrt{21}, (3, 2, 8)$
- (c) $21\sqrt{3}, (1, 2, 8)$
- (d) $3\sqrt{21}, (1, 2, 8)$

69. If $\Delta(x) = \begin{vmatrix} e^x & \sin x \\ \cos x & \ln(1+x^2) \end{vmatrix}$, then

the value of $\lim_{x \rightarrow 0} \frac{\Delta(x)}{x}$ is

- (a) 0
- (b) 2
- (c) -1
- (d) -2

70. The number of positive integral solutions of the equation

$$\tan^{-1} x + \cot^{-1} y = \tan^{-1} 3, \text{ is}$$

- (a) two
- (b) one
- (c) infinite
- (d) None of these

PART-II (Numerical Answer Questions)

71. A box contains two white balls, three black balls and four red balls. The number of ways such that three balls can be drawn from the box if at least one black ball is to be included in the draw is
72. Find the median from the following distribution.

Class	5–10	10–15	15–20	20–25	25–30
frequency	5	6	15	10	5

Class	30–35	35–40	40–45
frequency	4	2	2

73. If α, β are the roots of the equation $2x^2 + 3x + 5 = 0$, then the absolute value of the determinant $\begin{vmatrix} 0 & \beta & \beta \\ \alpha & 0 & \alpha \\ \beta & \alpha & 0 \end{vmatrix}$ is

74. $\int_{-3}^2 \{|x+1| + |x+2| + |x-1|\} dx$ is

75. The area bounded by the curve $y = 2x - x^2$ and the line $y = -x$ is

RESPONSE SHEET

PHYSICS

1. a b c d
2. a b c d
3. a b c d
4. a b c d
5. a b c d
6. a b c d
7. a b c d
8. a b c d
9. a b c d
10. a b c d
11. a b c d
12. a b c d
13. a b c d
14. a b c d
15. a b c d
16. a b c d
17. a b c d
18. a b c d
19. a b c d
20. a b c d

21.

22.

23.

24.

CHEMISTRY

26. a b c d
27. a b c d
28. a b c d
29. a b c d
30. a b c d
31. a b c d
32. a b c d
33. a b c d
34. a b c d
35. a b c d
36. a b c d
37. a b c d
38. a b c d
39. a b c d
40. a b c d
41. a b c d
42. a b c d
43. a b c d
44. a b c d
45. a b c d
46.
47.
48.
49.

MATHEMATICS

51. a b c d
52. a b c d
53. a b c d
54. a b c d
55. a b c d
56. a b c d
57. a b c d
58. a b c d
59. a b c d
60. a b c d
61. a b c d
62. a b c d
63. a b c d
64. a b c d
65. a b c d
66. a b c d
67. a b c d
68. a b c d
69. a b c d
70. a b c d
71.
72.
73.
74.

INSTRUCTIONS


1. This test will be a 3 hours Test.
2. This test consists of Physics, Chemistry and Mathematics questions with equal weightage of 100 marks.
3. Each question is of 4 marks.
4. There are three sections in the question paper consisting of Physics (Q.no.1 to 25), Chemistry (Q.no.26 to 50) and Mathematics (Q. no.51 to 75). Each section is divided into two parts, Part I consists of 20 multiple choice questions & Part II consists of 5 Numerical value type Questions.
5. There will be only one correct choice in the given four choices in Part I. For each question 4 marks will be awarded for correct choice, 1 mark will be deducted for incorrect choice for Part I Questions and zero mark will be awarded for not attempted question. For Part II Questions 4 marks will be awarded for correct answer and zero for unattempted and incorrect answer.
6. Any textual, printed or written material, mobile phones, calculator etc. is not allowed for the students appearing for the test.
7. All calculations / written work should be done in the rough sheet provided.

PHYSICS

PART-I (Multiple Choice Questions)

1. A bus is moving with a velocity of 10m/s on a straight road. A scooterist wishes to overtake the bus in 100 seconds. If the bus is at a distance of 1 km from the scooterist, at what velocity should the scooterist chase the bus?
(a) 50 m/sec (b) 40 m/sec
(c) 30 m/sec (d) 20 m/sec
2. The length of an elastic string is x when the tension is 5N . Its length is y when the tension is 7N . What will be its length, when the tension is 9N ?
(a) $2y+x$ (b) $2y-x$
(c) $7x-5y$ (d) $7x+5y$
3. A rod of length L is placed on x -axis between $x=0$ and $x=L$. The linear density i.e., mass per unit length denoted by ρ , of this rod, varies as, $\rho = a + bx$. What should be the dimensions of b ?
(a) $M^2L^1T^0$
(b) $M^1L^{-2}T^0$
(c) $M^{-1}L^3T^1$
(d) $M^{-1}L^2T^3$

4. A wheel is rolling on a plane road. The linear velocity of centre of mass is v . Then velocities of the points A and B on circumference of wheel relative to road will be



- (a) $v_A = v, v_B = 0$
 (b) $v_A = v_B = 0$
 (c) $v_A = 0, v_B = v$
 (d) $v_A = 0, v_B = 2v$
5. A metallic wire of density d is lying horizontal on the surface of water. The maximum length of wire so that it may not sink will be

$$(a) \sqrt{\frac{2Tg}{\pi d}} \quad (b) \sqrt{\frac{2\pi T}{dg}}$$


$$(c) \sqrt{\frac{2T}{\pi dg}} \quad (d) \text{any length}$$

6. Two points of a rod move with velocities $3v$ and v perpendicular to the rod and in the same direction, separated by a distance r . Then the angular velocity of the rod is
- (a) $3v/r$ (b) $4v/r$
 (c) $5v/r$ (d) $2v/r$
7. For hydrogen gas $C_p - C_v = a$ and for oxygen gas $C_p - C_v = b$. So, the relation between a and b is given by
- (a) $a = 16b$ (b) $16a = b$
 (c) $a = 4b$ (d) $a = b$

8. A bucket full of hot water is kept in a room and it cools from 75°C to 70°C in T_1 minutes, from 70°C to 65°C in T_2 minutes and from 65°C to 60°C in T_3 minutes. Then


- (a) $T_1 = T_2 = T_3$
 (b) $T_1 < T_2 < T_3$
 (c) $T_1 > T_2 > T_3$
 (d) $T_1 < T_3 < T_2$

9. The equivalent capacity of the network, (with all capacitors having the same capacitance C)




- (a) ∞
 (b) zero
 (c) $C[(\sqrt{3} - 1)/2]$
 (d) $C[(\sqrt{3} + 1)/2]$

10. The current I vs voltage V graphs for a given metallic wire at two different temperatures T_1 and T_2 are shown in the figure. It is concluded that




- (a) $T_1 > T_2$ (b) $T_1 < T_2$
 (c) $T_1 = T_2$ (d) $T_1 = 2T_2$

11. In the circuit shown the effective resistance between B and C is




- (a) 3Ω (b) 4Ω
 (c) $\frac{4}{3}\Omega$ (d) $\frac{3}{4}\Omega$
12. In the given circuit, the current drawn from the source is



- (a) $20A$ (b) $10A$
 (c) $5A$ (d) $5\sqrt{2} A$
13. A flat plate P of mass 'M' executes SHM in a horizontal plane by sliding over a frictionless surface with a frequency V . A block 'B' of mass 'm' rests on the plate as shown in figure. Coefficient of

friction between the surface of B and P is μ . What is the maximum amplitude of oscillation that the plate block system can have if the block B is not to slip on the plate :




- (a) $\frac{\mu g}{4\pi^2 V^2}$ (b) $\frac{\mu g}{4\pi^2 V^2}$
 (c) $\frac{\mu}{4\pi^2 V^2 g}$ (d) $\frac{\mu g}{2\pi^2 V^2}$

14. A glass slab has the left half of refractive index n_1 , and the right half of $n_2=3n_1$. The effective refractive index of the whole slab is

- (a) $\frac{n_1}{2}$ (b) $2n$
 (c) $\frac{3n_1}{2}$ (d) $\frac{2n_1}{3}$

15. In the arrangement shown L_1, L_2 are slits and S_1, S_2 two independent sources on the screen, interference fringes



- (a) will not be there
- (b) will not be there if the intensity of light reaching the screen from S_1 and S_2 are equal.
- (c) will be there under all circumstances
- (d) we will have only the central fringe


16. What is the ratio of the circumference of the first Bohr orbit for the electron in the hydrogen atom to the de Broglie wavelength of electrons having the same velocity as the electron in the first Bohr orbit of the hydrogen atom?

- (a) 1 : 1
- (b) 1 : 2
- (c) 1 : 4
- (d) 2 : 1

17. The radioactivity of a sample is R_1 at a time T_1 and R_2 at a time T_2 . If the half life of the specimen is T , the number of atoms that have disintegrated in the time $(T_2 - T_1)$ is proportional to


- (a) $(R_1 T_1 - R_2 T_2)$
- (b) $(R_1 - R_2)$
- (c) $(R_1 - R_2)/T$
- (d) $(R_1 - R_2) \times T$

18. P-V plots for two gases during adiabatic processes are shown in the figure. Plots 1 and 2 should correspond respectively to



- (a) He and Ar
- (b) He and O₂

19. Two identical thin rings, each of radius R metres, are coaxially placed at a distance R metres apart. If Q_1 coulomb and Q_2 coulomb are respectively, the charges uniformly spread on the two rings, the work done in moving a charge q from the centre of one ring to that of the other is




- (a) zero
- (b) $q(Q_1 - Q_2)$
- (c) $(\sqrt{2} - 1)/\sqrt{2} 4\pi\epsilon_0 R$
- (d) $q\sqrt{2}(Q_1 + Q_2)/4\pi\epsilon_0 R$
- (e) $q(Q_1 + Q_2)$
- (f) $(\sqrt{2} + 1)/\sqrt{2} 4\pi\epsilon_0 R$

20. The ratio of the coefficient of volume expansion of a glass container to that of a viscous liquid kept inside the container is 1 : 4. What fraction of the inner volume of the container should the liquid occupy so that the volume of the remaining vacant space will be same at all temperatures?

- (a) 2 : 5
- (b) 1 : 4
- (c) 1 : 64
- (d) 1 : 8

PART-II (Numerical Answer Questions)


21. The masses of the blocks A and B are 0.5 kg and 1 kg respectively. These are arranged as shown in the figure and are connected by a massless string. The coefficient of friction between all contact surfaces is 0.4. The force (in N) necessary to move the block B with constant velocity will be ($g = 10\text{m/s}^2$)



22. A body is thrown vertically upwards from the surface of earth in such a way that it reaches upto a height equal to $10R_e$. The velocity (in km/s) imparted to the body will be
23. A non-conducting partition divides a container into two equal compartments. One is filled with helium gas at 200 K and the other is filled with oxygen gas at 400 K. The number of molecules in each gas is the same. If the partition is removed to allow the gases to mix, the final temperature (in K) will be
24. A transformer is used to light a 140 W, 24 V bulb from a 240 V a.c. mains. The current in the main cable is 0.7 A. The efficiency (in %) of the transformer is

25. Two circular coils X and Y, having equal number of turns, carry equal currents in the same sense and subtend same solid angle at point O. If the smaller coil X is midway between O and Y, then if we represent the magnetic induction due to bigger coil Y at O as B_Y and due to smaller



coil X at O as B_X then the ratio $\frac{B_Y}{B_X}$ is



CHEMISTRY

PART-I (Multiple Choice Questions)

26. Which of the following structures does not contain any chiral C atom but represent the chirality in the structure.
- 2 – Ethyl – 3 – hexene
 - 2, 3-Pentadiene
 - 1,3 – Butadiene
 - Pent – 3 – en – 1 – yne
27. The root mean square speed of gas molecules at 25K & $1.5 \times 10^5 \text{ Nm}^{-2}$ is 100.5 ms^{-1} . If the temperature is raised to 100K & pressure to $6.0 \times 10^5 \text{ Nm}^{-2}$, the root mean square speed becomes.
- 100.5 ms^{-1}
 - 201.0 ms^{-1}
 - 402 ms^{-1}
 - 1608 ms^{-1}

- 28.** Reduction with aluminium isopropoxide in excess of isopropyl alcohol is called Meerwein Ponndorff-Verley reduction (MPV). What will be the final product when cyclohex-2-enone is selectively reduced in MPV reaction ?
- Cyclohexanol
 - Cyclohex-2-enol
 - Cyclohexanone
 - Benzene
- 29.** N_2 and O_2 are converted to mono cations N_2^+ and O_2^+ respectively, which of the following is wrong?
- In N_2^+ , the N – N bond weakens
 - In O_2^+ , the O – O bond order increases
 - In O_2^+ , paramagnetism decreases
 - N_2^+ becomes diamagnetic
- 30.** The reaction in which hydrogen peroxide acts as a reducing agent is
- $\text{PbS} + 4\text{H}_2\text{O}_2 \rightarrow \text{PbSO}_4 + 4\text{H}_2\text{O}$
 - $2\text{KI} + \text{H}_2\text{O}_2 \rightarrow 2\text{KOH} + \text{I}_2$
 - $2\text{FeSO}_4 + \text{H}_2\text{SO}_4 + \text{H}_2\text{O}_2 \rightarrow \text{Fe}_2(\text{SO}_4)_3 + 2\text{H}_2\text{O}$
 - $\text{Ag}_2\text{O} + \text{H}_2\text{O}_2 \longrightarrow 2\text{Ag} + \text{H}_2\text{O} + \text{O}_2$
- 31.** Reaction of  with RMgX leads to formation of
- RCHOHR
 - RCHOHCH_3
 - $\text{RCH}_2\text{CH}_2\text{OH}$
 - 
- 32.** Which reaction will not yield an amide?
- $\text{C}_2\text{H}_5 - \overset{\text{O}}{\underset{\parallel}{\text{C}}} - \text{Cl} + \text{NH}_3$
 - $\text{C}_2\text{H}_5 - \overset{\text{O}}{\underset{\parallel}{\text{C}}} - \text{O} - \overset{\text{O}}{\underset{\parallel}{\text{C}}} - \text{Cl} + \text{CH}_3\text{NH}_2$
 - $\text{CH}_3 - \overset{\text{O}}{\underset{\parallel}{\text{C}}} - \text{Cl} + (\text{CH}_3)_3\text{N}$
 - $\text{CH}_3 - \overset{\text{O}}{\underset{\parallel}{\text{C}}} - \text{O} - \overset{\text{O}}{\underset{\parallel}{\text{C}}} - \text{CH}_3 + \text{CH}_3\text{NH} - \text{CH}_2 - \text{CH}_3$
- 33.** How many different dipeptides can be formed by two different amino acids ?
- 4
 - 1
 - 3
 - 2
- 34.** When tert-butyl chloride is made to react with sodium methoxide, the major product is
- dimethyl ether
 - di-tert-butyl ether
 - tert-butylmethyl ether
 - isobutylene

35. If s_0 , s_1 , s_2 and s_3 are the solubilities of AgCl in water, 0.01 M CaCl_2 , 0.01 M NaCl and 0.05 M AgNO_3 solutions, respectively, then
- $s_0 > s_1 > s_2 > s_3$
 - $s_0 > s_2 > s_1 > s_3$
 - $s_0 > s_2 > s_3 > s_1$
 - $s_0 > s_1 = s_2 > s_3$
36. An organic compound is treated with NaNO_2 and dil. HCl at 0°C . The resulting solution is added to an alkaline solution of β -naphthol where by a brilliant red dye is produced. It shows the presence of
- $-\text{NO}_2$ group
 - aromatic $-\text{NH}_2$ group
 - $-\text{CONH}_2$ group
 - aliphatic $-\text{NH}_2$ group
37. Point out the incorrect statement among the following :
- The oxidation state of oxygen is +2 in OF_2 .
 - Acidic character follows the order $\text{H}_2\text{O} < \text{H}_2\text{S} < \text{H}_2\text{Se} < \text{H}_2\text{Te}$.
 - The tendency to form multiple bonds increases in moving down the group from sulphur to tellurium (towards C and N)
 - Sulphur has a strong tendency to catenate while oxygen shows this tendency to a limited extent.
38. Removal of Fe, Cu, W from Sn metal after smelting is by because
- Poling; of more affinity towards oxygen for impurities
 - Selective oxidation; of more affinity towards oxygen for impurities
 - Electrolytic refining; impurities undissolved in electrolyte
 - Liquation; Sn having low melting point compared to impurities.
39. Among KO_2 , AlO_2^- , BaO_2 and NO_2^+ , unpaired electron is present in
- NO_2^+ and BaO_2
 - KO_2 and AlO_2^-
 - KO_2 only
 - BaO_2 only
40. If a 0.1 M solution of glucose (Mol. wt 180) and 0.1 molar solution of urea (Mol. wt. 60) are placed on two sided semipermeable membrane to equal heights, then it will be correct to say that
- there will be no net movement across the membrane
 - glucose will flow across the membrane into urea solution
 - urea will flow across the membrane into glucose solution
 - water will flow from urea solution to glucose solution

41. When pink $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$ is dehydrated the colour changes to blue. The correct explanation for the change is :
- The octahedral complex becomes square planar.
 - A tetrahedral complex is formed.
 - Distorted octahedral structure is obtained.
 - Dehydration results in the formation of polymeric species.
42. Amongst the following the compound that is both paramagnetic and coloured is
- $\text{K}_2\text{Cr}_2\text{O}_7$
 - $(\text{NH}_4)_2[\text{TiCl}_6]$
 - CoSO_4
 - $\text{K}_3[\text{Cu}(\text{CN})_4]$
43. A reaction rate constant is given by $K = 1.2 \times 10^{10} e^{-2500/RT}$. It means
- $\log K$ vs T will give a straight line
 - $\log K$ vs $1/T$ gives a straight line with a slope $-2500/2.303 R$
 - half life of reaction will be more at higher temperature
 - $\log K$ vs $1/T$ gives a straight line with a slope $2500/R$
44. The correct statement among the following is :
- The alkali metals when strongly heated in oxygen form superoxides.
 - Caesium is used in photoelectric cells.
45. The e.m.f. of a Daniell cell, $\text{Zn}|\text{ZnSO}_4||\text{CuSO}_4|\text{Cu}$, at 298 K is E_1 . When the concentration of ZnSO_4 is 1.0 M and that of CuSO_4 is 0.01 M, the e.m.f. changed to E_2 . What is the relationship between E_1 and E_2 ?
- $E_1 < E_2$
 - $E_1 = E_2$
 - $E_2 = 0 \neq E_1$
 - $E_1 > E_2$

PART-II (Numerical Answer Questions)

46. The vapour pressure of benzene at a certain temperature is 640 mm of Hg. A non volatile and non electrolyte solid weighing 2.175 g is added to 39.08 g of benzene. If the vapour pressure of the solution is 600 mm of Hg, what is the molecular weight of solid substance?
47. What will be the uncertainty in the position of an electron (mass 9.1×10^{-31} g) moving with a velocity of 3.0×10^6 cm s⁻¹ accurate up to 0.011%?
48. When CO_2 dissolves in water, the following equilibrium is established
- $$\text{CO}_2 + 2\text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{HCO}_3^-;$$
- for which the equilibrium constant is 3.8×10^{-6} and pH = 6.0. What would be the ratio of concentration of bicarbonate ion to carbon dioxide?

49. The wave number of first line of Balmer series of hydrogen is 15200 cm^{-1} . What will be the wave number of first Balmer line of Li^{+2} ion?

50. A cylinder of gas supplied by Bharat Petroleum is assumed to contain 14 kg of butane. If a normal family requires 20,000 kJ of energy per day for cooking, butane gas in the cylinder last for Days.
 $(\Delta H_c \text{ of } \text{C}_4\text{H}_{10} = -2658 \text{ JK per mole})$

53. If $0 < \alpha, \beta, \gamma < \pi/2$ such that $\alpha + \beta + \gamma = \frac{\pi}{2}$ and $\cot \alpha, \cot \beta, \cot \gamma$ are in arithmetic progression, then the value of $\cot \alpha \cot \gamma$ is
(a) 1 (b) 3
(c) $\cot^2 \beta$ (d) $\cot \alpha + \cot \gamma$

54. If $\omega = \cos \frac{\pi}{n} + i \sin \frac{\pi}{n}$, then value of $1 + \omega + \omega^2 + \dots + \omega^{n-1}$ is

MATHEMATICS

PART-I (Multiple Choice Questions)

51. If a, b, c, d and p are distinct non zero real numbers such that $(a^2 + b^2 + c^2)p^2 - 2(ab + bc + cd)p + (b^2 + c^2 + d^2) \leq 0$ then a, b, c, d are in

 - A.P.
 - GP.
 - H.P.
 - satisfy $ab = cd$

52. Which of the following is correct?

 - If $a^2 + 4b^2 = 12ab$, then $\log(a + 2b) = \frac{1}{2}(\log a + \log b)$
 - If $\frac{\log x}{b-c} = \frac{\log y}{c-a} = \frac{\log z}{a-b}$, then $x^a \cdot y^b \cdot z^c = abc$
 -
 - All are correct

53. If the solution of the linear equations $x - 2y + z = 0$; $2x - y + 3z = 0$ and $\lambda x + y - z = 0$ is

 - None of these
 - The circles $x^2 + y^2 - 2x - 15 = 0$ and $x^2 + y^2 + 4y + 3 = 0$ have
 - no common tangent
 - one common tangent
 - three common tangents
 - four common tangents
 - Which of the following is correct?
 - If A and B are square matrices of order 3 such that $|A| = -1$, $|B| = 3$, then the determinant of $3AB$ is equal to 27.
 - If A is an invertible matrix, then $\det(A^{-1})$ is equal to $\det(A)$
 - If A and B are matrices of the same order, then $(A + B)^2 = A^2 + 2AB + B^2$ is possible if $AB = I$
 - None of these

trivial then the value of λ is given by

(a) $\lambda = -\frac{4}{5}$ (b) $\lambda \neq -\frac{4}{5}$

(c) $\lambda = 2$ (d) $\lambda \neq 2$

58. Let $f(x) = |x - 1|$. Then

- (a) $f(x^2) = (f(x))^2$
 (b) $f(x+y) = f(x) + f(y)$
 (c) $f(|x|) = |f(x)|$
 (d) None of these

59. If

$$\sin^{-1} \frac{2a}{1+a^2} + \sin^{-1} \frac{2b}{1+b^2} = 2 \tan^{-1} x,$$

then x is equal to

- (a) $\frac{a-b}{1+ab}$ (b) $\frac{b}{1+ab}$
 (c) $\frac{b}{1-ab}$ (d) $\frac{a+b}{1-ab}$

60. If $AB = 0$, then for the matrices

$$A = \begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix}$$

$$\text{and } B = \begin{bmatrix} \cos^2 \phi & \cos \phi \sin \phi \\ \cos \phi \sin \phi & \sin^2 \phi \end{bmatrix},$$

$\theta - \phi$ is

- (a) an odd number of $\frac{\pi}{2}$
 (b) an odd multiple of π
 (c) an even multiple of $\frac{\pi}{2}$
 (d) 0

61. The set of points where $f(x) = (x-1)^2 (x + |x-1|)$ is thrice differentiable, is

- (a) \mathbb{R} (b) $\mathbb{R} - \{0\}$
 (c) $\mathbb{R} - \{1\}$ (d) $\mathbb{R} - \{0,1\}$

62. Let $f(x) = 1/(x-1)$ and $g(x) = 1/(x^2+x-2)$. Then the set of points where $(gof)(x)$ is discontinuous, is

- (a) $\{1\}$ (b) $\{-2, 1\}$
 (c) $\{1/2, 1, 2\}$ (d) $\{1/2, 1\}$

63. $\sum_{r=0}^m {}^{n+r} C_n$ is equal to :

- (a) ${}^{n+m+1} C_{n+1}$
 (b) ${}^{n+m+2} C_n$
 (c) ${}^{n+m+3} C_{n-1}$
 (d) None of these

64. Let $f(x) = \frac{x - \{x+1\}}{x - \{x+2\}}$; where $\{x\}$ is the fractional part of x , then

$$\lim_{x \rightarrow 1/3} f(x)$$

- (a) has value 0
 (b) has value 1
 (c) has value $-\infty$
 (d) has value ∞

65. The order of the differential equation

$$\left[1 + 5 \left(\frac{dy}{dx} \right)^2 \right]^{3/2} = 11 \left(\frac{d^2 y}{dx^2} \right)^5 \text{ is}$$

66. The value of

$$\int_{-\pi/4}^{\pi/4} (x|x| + \sin^3 x + x \tan^2 x + 1) dx$$
 is

- (a) 0 (b) 1
 (c) $\pi/4$ (d) $\pi/2$

67. Let $(1-x-2x^2)^6 = 1 + a_1x + a_2x^2 + \dots + a_{12}x^{12}$. Then

$$\frac{a_2}{2^2} + \frac{a_4}{2^4} + \frac{a_6}{2^6} + \dots + \frac{a_{12}}{2^{12}}$$
 is equal to

- (a) -1 (b) -1/2
 (c) 0 (d) 1/2

68. The equation of a common tangent to $y^2 = 4x$ and the curve $x^2 + 4y^2 = 8$ can be

- (a) $x - 2y + 2 = 0$
 (b) $x + 2y + 4 = 0$
 (c) $x - 2y = 4$
 (d) $x + 2y = 4$

69. The function $f(x) = (x-3)^2$ satisfies all the conditions of mean value theorem in $\{3, 4\}$. A point on $y = (x-3)^2$, where the tangent is parallel to the chord joining $(3, 0)$ and $(4, 1)$ is

- (a) $\left(\frac{7}{2}, \frac{1}{2}\right)$ (b) $\left(\frac{7}{2}, \frac{1}{4}\right)$
 (c) $(1, 4)$ (d) $(4, 1)$

70. If $x + y - z + xyz = 0$, then $\frac{2x}{1-x^2} + \frac{2y}{1-y^2} - \frac{2z}{1-z^2}$ is equal to

(a) $\frac{xyz}{[(1-x^2)(1-y^2)(1-z^2)]}$

(b) $\frac{-xyz}{[(1-x^2)(1-y^2)(1-z^2)]}$

(c) $\frac{8xyz}{[(1-x^2)(1-y^2)(1-z^2)]}$

(d) $\frac{-8xyz}{[(1-x^2)(1-y^2)(1-z^2)]}$

PART-II (Numerical Answer Questions)

71. If one root of the equation $x^2 + px + 12 = 0$ is 4 while the equation $x^2 + px + q = 0$ has equal roots, the value of q is

72. The value of $\cos 36^\circ \cos 42^\circ / \cos 78^\circ$ is

$$\left[\text{Given : } \sin 18^\circ = \frac{\sqrt{5}-1}{4} \text{ and } \cos 36^\circ = \frac{\sqrt{5}+1}{4} \right]$$

73. If $x = 1/5$, the absolute value of $\cos(\cos^{-1}x + 2\sin^{-1}x)$ is

74. If θ_1, θ_2 are the solutions of the equation $2\tan^2\theta - 4\tan\theta + 1 = 0$, then $\tan(\theta_1 + \theta_2)$ is equal to

75. In a ΔABC , if $\begin{vmatrix} 1 & a & b \\ 1 & c & a \\ 1 & b & c \end{vmatrix} = 0$, then

$$\sin^2 A + \sin^2 B + \sin^2 C =$$

RESPONSE SHEET

PHYSICS

1. a b c d
2. a b c d
3. a b c d
4. a b c d
5. a b c d
6. a b c d
7. a b c d
8. a b c d
9. a b c d
10. a b c d
11. a b c d
12. a b c d
13. a b c d
14. a b c d
15. a b c d
16. a b c d
17. a b c d
18. a b c d
19. a b c d
20. a b c d

21.

22.

23.

24.

CHEMISTRY

26. a b c d
27. a b c d
28. a b c d
29. a b c d
30. a b c d
31. a b c d
32. a b c d
33. a b c d
34. a b c d
35. a b c d
36. a b c d
37. a b c d
38. a b c d
39. a b c d
40. a b c d
41. a b c d
42. a b c d
43. a b c d
44. a b c d
45. a b c d

46.

47.

48.

49.

MATHEMATICS

51. a b c d
52. a b c d
53. a b c d
54. a b c d
55. a b c d
56. a b c d
57. a b c d
58. a b c d
59. a b c d
60. a b c d
61. a b c d
62. a b c d
63. a b c d
64. a b c d
65. a b c d
66. a b c d
67. a b c d
68. a b c d
69. a b c d
70. a b c d

71.

72.

73.

74.

MOCK TEST

3


INSTRUCTIONS

1. This test will be a 3 hours Test.
2. This test consists of Physics, Chemistry and Mathematics questions with equal weightage of 100 marks.
3. Each question is of 4 marks.
4. There are three sections in the question paper consisting of Physics (Q.no.1 to 25), Chemistry (Q.no.26 to 50) and Mathematics (Q. no.51 to 75). Each section is divided into two parts, Part I consists of 20 multiple choice questions & Part II consists of 5 Numerical value type Questions.
5. There will be only one correct choice in the given four choices in Part I. For each question 4 marks will be awarded for correct choice, 1 mark will be deducted for incorrect choice for Part I Questions and zero mark will be awarded for not attempted question. For Part II Questions 4 marks will be awarded for correct answer and zero for unattempted and incorrect answer.
6. Any textual, printed or written material, mobile phones, calculator etc. is not allowed for the students appearing for the test.
7. All calculations / written work should be done in the rough sheet provided.

PHYSICS

PART-I (Multiple Choice Questions)

1. Astronauts look down on earth surface from a space ship parked at an altitude of 500 km. They can resolve objects of the earth of the size (It can be assumed that the pupils diameter is 5mm and wavelength of light is 500 nm)
(a) 0.5m (b) 5m
(c) 50m (d) 500m
2. The wavelength of sodium light in air is 5890 Å. The velocity of light in air is 3×10^{-8} ms⁻¹. The wavelength of light in a glass of refractive index 1.6, would be close to
3. (a) 5890 Å (b) 3680 Å
(c) 9424 Å (d) 15078 Å
4. A space craft of mass 'M', moving with velocity 'v' suddenly breaks into two pieces. After the explosion mass 'm' becomes stationary. What is the velocity of the other part of the craft ?
(a) $\frac{Mv}{M-m}$ (b) v
(c) $\frac{mv}{M}$ (d) $\frac{M-m}{m}v$
4. Using mass(M), length(L), time(T) and electric current (A) as fundamental quantities the dimensions of permittivity will be

- (a) $MLT^{-1}A^1$
 (b) $MLT^{-2}A^{-2}$
 (c) $M^{-1}L^{-3}T^{+4}A^2$
 (d) $M^2L^{-2}T^{-2}A^2$
5. A black body at a temperature of 227°C radiates heat at the rate of $20 \text{ cal m}^{-2} \text{ s}^{-1}$. When its temperature rises to 727°C the heat radiated will be
 (a) 40 units (b) 160 units
 (c) 320 units (d) 640 units
6. Two waves of wavelengths 99 cm and 100 cm both travelling with velocity 396 m/s are made to interfere. The number of beats produced by them per second are
 (a) 1 (b) 2
 (c) 4 (d) 8
7. A sphere of mass 'm' and radius 'r' is falling in the column of a viscous fluid. Terminal velocity attained by falling object is proportional to
 (a) r^2 (b) $1/r$
 (c) r (d) $-1/r^2$
8. There are two wires of the same length. The diameter of second wire is twice that of the first. On applying the same load to both the wires, the extension produced in them will be in ratio of
 (a) 1 : 4 (b) 1 : 2
 (c) 2 : 1 (d) 4 : 1
9. When a proton, anti-proton annihilate the energy released is
 (a) $1.5 \times 10^{-10} \text{ J}$
 (b) $28.8 \times 10^{-10} \text{ J}$
 (c) $6 \times 10^{-10} \text{ J}$
 (d) $9 \times 10^{-10} \text{ J}$
10. $y = 2 (\text{cm}) \sin \left[\frac{\pi t}{2} + \phi \right]$
 What is the maximum acceleration of the particle doing the SHM
 (a) $\frac{\pi}{2} \text{ cm/s}^2$ (b) $\frac{\pi^2}{2} \text{ cm/s}^2$
 (c) $\frac{\pi^2}{4} \text{ cm/s}^2$ (d) $\frac{\pi}{4} \text{ cm/s}^2$
11. What is the electric potential at the centre of the square?
- 
- (a) zero
 (b) $kq/a\sqrt{2}$
 (c) kq/a^2
 (d) None of these
12. A rectangular block of mass m and area of cross-section A floats in a liquid of density ρ . If it is given a small vertical displacement from equilibrium it undergoes oscillation with a time period T . Then
 (a) $T \propto \frac{1}{\sqrt{A}}$ (b) $T \propto \frac{1}{\rho}$
 (c) $T \propto \frac{1}{\sqrt{m}}$ (d) $T \propto \sqrt{\rho}$
13. While determining the specific resistance of a wire using a metre bridge the formula used is (where X , D , L and ρ denote unknown resistance, diameter of the wire, the length of the wire and the specific resistance of the wire)


(a) $\rho = \frac{X\pi D}{4L}$

(b) $\rho = \frac{X\pi D^2}{4L}$

(c) $\rho = \frac{X^2\pi D^2}{4L}$

(d) $\rho = \frac{X\pi D^2}{4L^2}$

14. Consider the following u-v diagram regarding the experiment to determine the focal length of a convex lens.



At the point A, the values of u and v are equal. The focal length of the lens is

- (a) 40 cm (b) 20 cm
(c) 10 cm (d) 15 cm


15. Two metallic plates A and B, each of area $5 \times 10^{-4} \text{ m}^2$, are placed parallel to each other at a separation of 1 cm. Plate B carries a positive charge of $33.7 \times 10^{-12} \text{ C}$. A mono-chromatic beam of light, with photons of energy 5 eV each, starts falling on plate A at $t = 0$ so that 10^{16} photons fall on it per square meter per second. Assume that one photoelectron is emitted for every 10^6 incident photons. Also assume

that all the emitted photoelectrons are collected by plate B and the work function of plate A remains constant at the value 2 eV.

No. of photoelectrons emitted up to 10 sec

- (a) 5×10^7 (b) 2×10^6
(c) 5×10^6 (d) 2×10^7

16. The circuit shown here has two batteries of 8.0 V and 16.0 V and three resistors 3Ω , 9Ω and 9Ω and a capacitor of $5.0 \mu\text{F}$.



How much is the current I in the circuit in steady state?

- (a) 1.6 A (b) 0.67 A
(c) 2.5 A (d) 0.25 A

17. An electromagnetic wave of frequency 1×10^{14} hertz is propagating along z-axis. The amplitude of electric field is 4 V/m. If $\epsilon_0 = 8.8 \times 10^{-12} \text{ C}^2/\text{N}\cdot\text{m}^2$, then average energy density of electric field will be:

- (a) $35.2 \times 10^{-10} \text{ J/m}^3$
(b) $35.2 \times 10^{-11} \text{ J/m}^3$
(c) $35.2 \times 10^{-12} \text{ J/m}^3$
(d) $35.2 \times 10^{-13} \text{ J/m}^3$

18. A block is placed on a frictionless horizontal table. The mass of the block is m and springs are attached on either side with force constants K_1 and K_2 . If the block is displaced a little and left to oscillate, then the angular frequency of oscillation will be

$$(a) \quad \left(\frac{K_1 + K_2}{m} \right)^{\frac{1}{2}}$$

$$(b) \left[\frac{K_1 K_2}{m(K_1 + K_2)} \right]^{\frac{1}{2}}$$

$$(c) \left[\frac{K_1 K_2}{(K_1 - K_2)m} \right]^{\frac{1}{2}}$$

$$(d) \quad \left[\frac{K_1^2 + K_2^2}{(K_1 + K_2)m} \right]^{\frac{1}{2}}$$

19. A sphere is placed in front of a convex lens of focal length f . The radius of the sphere is much smaller compared to f . The image of the sphere would look spherical if the object distance is

(a) f (b) $\frac{3f}{2}$
 (c) $2f$ (d) $\frac{f}{2}$

20. Which of the following expressions corresponds to simple harmonic motion along a straight line, where x is the displacement and a, b, c are positive constants?

(a) $a + bx - cx^2$
 (b) bx^2
 (c) $a - bx + cx^2$
 (d) $-bx$

PART-II (Numerical Answer Questions)

21. The source of sound generating a frequency of 3 kHz reaches an observer with a speed of 0.5 times in air. The frequency (in kHz) heard by the observer is

22. The temperature of reservoir of Carnot's engine operating with an efficiency of 70% is 1000 kelvin. The temperature (in kelvin) of its sink is

23. The escape velocity for a body of mass 1 kg from the earth surface is 11.2 kms^{-1} . The escape velocity (in kms^{-1}) for a body of mass 100 kg would be

24. At the centre of a circular coil of radius 5 cm carrying current, magnetic field due to earth is $0.5 \times 10^{-5} \text{ W/m}^2$. What should be the current (in A) flowing through the coil so that it annuls the earth's magnetic field

25. A beam of light of intensity 12 watt/cm² is incident on a totally reflecting plane mirror of area 1.5 cm², then the force (in newton) acting on the mirror will be

CHEMISTRY

PART-I (Multiple Choice Questions)

26. Ethylene dichloride and ethylidine chloride are isomeric compounds. The false statement about these isomers is that they :

 - react with alcoholic potash and give the same product
 - are position isomers
 - contain the same percentage of chlorine
 - are both hydrolysed to the same product

27. An aqueous solution of sodium carbonate has a pH greater than 7 because :

 - it contains more carbonate ions than H_2O molecules
 - contains more sodium ions than carbonate ions
 - Na^+ ions react with water
 - carbonate ions react with H_2O

- 28.** By what ratio the average velocity of the molecule in gas changes when the temperature is raised from 50 to 200°C ?
- (a) $\frac{1.21}{1}$ (b) $\frac{1.46}{1}$
 (c) $\frac{2}{1}$ (d) $\frac{4}{1}$
- 29.** How many H-atoms are present in 0.046 g of ethanol ?
- (a) 6×10^{20} (b) 1.2×10^{21}
 (c) 3×10^{21} (d) 3.6×10^{21}
- 30.** A metal M reacts with N₂ to give a compound 'A' (M₃N). 'A' on heating at high temperature gives back 'M' and 'A' on reacting with H₂O gives a gas 'B'. 'B' turns CuSO₄ solution blue on passing through it. M and B can be :
- (a) Al & NH₃ (b) Li & NH₃
 (c) Na & NH₃ (d) Mg & NH₃
- 31.** When ethanal reacts with CH₃MgBr and C₂H₅OH/dry HCl, the product formed are :
- (a) ethyl alcohol and 2-propanol
 (b) ethane and hemi acetal
 (c) 2-propanol and acetal
 (d) propane and methyl acetate
- 32.** If the solutions of NaCl and NaNO₃ are mixed in one beaker and the temperature adjusted to 383°K, the contents of the beaker will most likely:
- (a) freeze
 (b) boil
 (c) exhibit precipitation of NaNO₃
 (d) exhibit a marked color change
- 33.** Given the molecular formula of the hexa-coordinated complexes (i) CoCl₃.6NH₃, (ii) CoCl₃.5NH₃, (iii) CoCl₃.4NH₃. If the number of co-ordinated NH₃ molecules in i, ii and iii respectively
- are 6, 5, 4, the primary valencies in (i), (ii) and (iii) are :
 (a) 6, 5, 4 (b) 3, 2, 1
 (c) 0, 1, 2 (d) 3, 3, 3
- 34.** Polyethylene is
- (a) Random copolymer
 (b) Homopolymer
 (c) Alternate copolymer
 (d) Crosslinked copolymer
- 35.** Which of the following is used in our body as a fuel for muscles and nerves and to build and repair body tissues?
- (a) Cane sugar (b) Fructose
 (c) Proteins (d) Glucose
- 36.** With a change in hybridization of the carbon bearing the charge, the stability of a carbanion decreases in the order :
- (a) sp < sp² < sp³
 (b) sp < sp³ < sp²
 (c) sp³ < sp² < sp
 (d) sp² < sp < sp³
- 37.** In O₂⁻, O₂ and O₂²⁻ molecular species, the total number of antibonding electrons respectively are :
- (a) 7, 6, 8 (b) 1, 0, 2
 (c) 6, 6, 6 (d) 8, 6, 8
- 38.** Which of the given sets of temperature and pressure will cause a gas to exhibit the greatest deviation from ideal gas behaviour?
- (a) 100 °C & 4 atm
 (b) 100 °C & 2 atm
 (c) -100 °C & 4 atm
 (d) 0 °C & 2 atm
- 39.** Which of the following pairs has heat of neutralisation equal to 13.7 kcal's ?
- (a) HCl, NH₄OH
 (b) HNO₃, KOH
 (c) NaOH, CH₃COOH
 (d) H₂SO₄, NH₄OH

- 40.** The relative abundance of two isotopes of atomic weight 85 and 87 is 75% and 25% respectively. The average atomic weight of element is
 (a) 75.5 (b) 85.5
 (c) 40.0 (d) 86.0
- 41.** In Kjeldahl's method, nitrogen present in the organic compound is quantitatively converted into
 (a) ammonium nitrite
 (b) ammonium sulphate
 (c) ammonium phosphate
 (d) ammonium nitrate
- 42.** An organic amino compound reacts with aqueous nitrous acid at low temperature to produce an oily nitrosoamine. The compound is:
 (a) CH_3NH_2
 (b) $\text{CH}_3\text{CH}_2\text{NH}_2$
 (c) $\text{CH}_3\text{CH}_2\text{NH}.\text{CH}_2\text{CH}_3$
 (d) $(\text{CH}_3\text{CH}_2)_3\text{N}$
- 43.** Which reaction characteristics are changed by the addition of a catalyst to a reaction at constant temperature ?
 (i) Activation energy
 (ii) Equilibrium constant
 (iii) Reaction enthalpy
 (a) (i) only
 (b) (iii) only
 (c) (i) and (ii) only
 (d) all of these
- 44.** Which is not true for beryllium ?
 (a) Beryllium is amphoteric
 (b) It forms unusual carbide, Be_2C
 (c) $\text{Be}(\text{OH})_2$ is basic
 (d) Beryllium halides are electron deficient
- 45.** A fire of lithium, sodium and potassium can be extinguished by
 (a) H_2O
 (b) Nitrogen
 (c) CO_2
 (d) Asbestos blanket

PART-II (Numerical Answer Questions)

- 46.** What is the molarity of H_2SO_4 solution if 25ml is exactly neutralized with 32.63 ml of 0.164 M, NaOH ?
- 47.** 3.92 g of ferrous ammonium sulphate react completely with 50 ml $\frac{\text{N}}{10}$ KMnO_4 solution. What will be the percentage purity of the sample?
- 48.** An 'fcc' in a unit cell of aluminium contains the equivalent of how many atoms ?
- 49.** If K_{sp} of Ag_2CO_3 is 8, the molar solubility of Ag_2CO_3 in 0.1 M AgNO_3 is :
- 50.** When 5 litres of a gas mixture of methane and propane is perfectly combusted at 0 °C and 1 atmosphere, 16 litres of oxygen at the same temperature and pressure is consumed. The amount of heat released from this combustion in kJ
 $[\Delta H_{\text{comb.}}(\text{CH}_4) = 890 \text{ kJ mol}^{-1}$,
 $\Delta H_{\text{comb.}}(\text{C}_3\text{H}_8) = 2220 \text{ kJ mol}^{-1}]$ is

MATHEMATICS

PART-I (Multiple Choice Questions)

51. The function $f(x) = \tan^{-1}(\sin x + \cos x)$ is an increasing function in
- (a) $\left(0, \frac{\pi}{2}\right)$ (b) $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
(c) $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$ (d) $\left(-\frac{\pi}{2}, \frac{\pi}{4}\right)$
52. The degree of differential equation satisfying the relation $\sqrt{1+x^2} + \sqrt{1+y^2} = \lambda(x\sqrt{1+y^2} - y\sqrt{1+x^2})$ is
- (a) 1 (b) 2
(c) 3 (d) 4
53. $\int_0^2 [x^2] dx$, where $[x]$ is the greatest integer $\leq x$ is
- (a) $5 + \sqrt{2} + \sqrt{3}$
(b) $-5 + \sqrt{2} - \sqrt{3}$
(c) $5 - \sqrt{2} - \sqrt{3}$
(d) $-4 + \sqrt{3} - \sqrt{2}$
54. α, β be the roots of $x^2 - 3x + a = 0$ and γ, δ be the roots of $x^2 - 12x + b = 0$ and numbers $\alpha, \beta, \gamma, \delta$ (in order) form an increasing G.P. then
- (a) $a = 3, b = 12$
(b) $a = 12, b = 3$
(c) $a = 2, b = 32$
(d) $a = 4, b = 16$
55. Assume R and S are (non-empty) relations in a set A. Which of the following relation given below is false
- (a) If R and S are transitive, then $R \cup S$ is transitive.
(b) If R and S are transitive, then $R \cap S$ is transitive.
(c) If R and S are symmetric, then $R \cup S$ is symmetric.
(d) If R and S are reflexive, Then $R \cap S$ is reflexive.
56. $\int \log 2x dx$ is
- (a) $x \log 2x - \frac{x^2}{2}$
(b) $x \log 2x - \frac{x}{2}$
(c) $x^2 \log 2x - \frac{x}{2}$
(d) $x \log 2x - x + c$
57. For a given integer k, in the interval $\left[2\pi k + \frac{\pi}{2}, 2\pi k - \frac{\pi}{2}\right]$ the graph of $\sin x$ is
- (a) increasing from -1 to 1
(b) decreasing from -1 to 0
(c) decreasing from 0 to 1
(d) None of these
58. $\frac{dy}{dx} + y = 2e^{2x}$ then y is
- (a) $ce^{-x} + \frac{2}{3}e^{2x}$
(b) $(1+x)e^{-x} + \frac{2}{3}e^{2x} + c$
(c) $ce^{-x} + \frac{2}{3}e^{2x} + c$
(d) $e^{-x} + \frac{2}{3}e^{2x} + c$

59. If $\begin{vmatrix} a+x & a-x & a-x \\ a-x & a+x & a-x \\ a-x & a-x & a+x \end{vmatrix} = 0$ then
 x is
 (a) $0, 2a$
 (b) $a, 2a$
 (c) $0, 3a$
 (d) None of these
60. The equation of a circle with origin as centre and passing through the vertices of an equilateral triangle whose median is of length $3a$ is
 (a) $x^2 + y^2 = 9a^2$
 (b) $x^2 + y^2 = 16a^2$
 (c) $x^2 + y^2 = 4a^2$
 (d) $x^2 + y^2 = a^2$
61. If a positive integer n is divisible by 9, then the sum of the digits of n is divisible by 9. So which statement is its contrapositive.
 (a) (sum of digits of n is divisible by 9)
 $\Rightarrow (n \text{ is divisible by 9})$
 (b) (sum of digits of n is not divisible by 9)
 $\Rightarrow (n \text{ is not divisible by 9})$
 (c) (sum of digits of n is divisible by 9)
 $\Rightarrow (n \text{ is divisible by 9})$
 (d) none of these
62. Fifteen coupons are numbered 1, 2 15, respectively. Seven coupons are selected at random one at a time with replacement. The probability that the largest number appearing on a selected coupon is 9, is
 (a) $\left(\frac{9}{16}\right)^6$
 (b) $\left(\frac{8}{15}\right)^7$
 (c) $\left(\frac{3}{5}\right)^7$
 (d) None of these
63. If $a \leq 0$ then roots of $x^2 - 2a|x-a| - 3a^2 = 0$ is
 (a) $(-1+\sqrt{6})a$
 (b) $(\sqrt{6}-1)a$
 (c) a
 (d) None of these
64. If X and Y are two sets, then $X \cap (X \cup Y)^c$ equals.
 (a) X
 (b) Y
 (c) \emptyset
 (d) None of these
65. If $y = \log_2 \{\log_2(x)\}$, then $\frac{dy}{dx}$ is
 (a) $\frac{\log_2 e}{x \ln x}$
 (b) $\frac{2.3026}{x \ln x \ln 2}$
 (c) $\frac{1}{\ln(2x)^x}$
 (d) None of these

66. $f(x) = \sin|x|$. $f(x)$ is not differentiable at
 (a) $x = 0$ only
 (b) all x
 (c) multiples of π
 (d) multiples of $\frac{\pi}{2}$

67. $\int_0^{\pi/3} \frac{\cos x + \sin x}{\sqrt{1 + \sin 2x}} dx$ is
 (a) $\frac{4\pi}{3}$ (b) $\frac{2\pi}{3}$
 (c) π (d) $\frac{\pi}{3}$

68. The angle between the pair of tangents drawn to the ellipse $3x^2 + 2y^2 = 5$ from the point $(1,2)$ is
 (a) $\tan^{-1}\left(\frac{12}{5}\right)$
 (b) $\tan^{-1}(6\sqrt{5})$
 (c) $\tan^{-1}\left(\frac{12}{\sqrt{5}}\right)$
 (d) $\tan^{-1}(12\sqrt{5})$

69. Let α and β be the roots of the equation $x^2 + x + 1 = 0$, the equation whose roots are α^{19}, β^7 is
 (a) $x^2 - x - 1 = 0$
 (b) $x^2 - x + 1 = 0$
 (c) $x^2 + x - 1 = 0$
 (d) $x^2 + x + 1 = 0$

70. The x satisfying $\sin^{-1} x + \sin^{-1}(1-x) = \cos^{-1} x$ are
 (a) 1, 0
 (b) 1, -1
 (c) $0, \frac{1}{2}$
 (d) None of these

PART-II (Numerical Answer Questions)

71. Area between curves $y = x^2$, $x = y^2$ is
 72. The probability of A = Probability of
 $B = \text{Probability of } C = \frac{1}{4}$

$$P(A) \cap P(B) \cap P(C) = 0, P(B \cap C) = 0$$

$$\text{and } P(A \cap C) = \frac{1}{8}, P(A \cap B) = 0$$

- the probability that atleast one of the events A, B, C exists is
 73. Coefficient of x^6 in the expansion

$$\left(x + \frac{1}{x^2}\right)^6$$

74. $f(x) = \frac{\sin 3x}{\sin x}$, when $x \neq 0$
 $= k$, when $x = 0$

- for the function to be continuous k should be
 75. A line passes through $(2,2)$ and is perpendicular to the line $3x + y = 3$ its y intercept is

RESPONSE SHEET

PHYSICS

1. a b c d
2. a b c d
3. a b c d
4. a b c d
5. a b c d
6. a b c d
7. a b c d
8. a b c d
9. a b c d
10. a b c d
11. a b c d
12. a b c d
13. a b c d
14. a b c d
15. a b c d
16. a b c d
17. a b c d
18. a b c d
19. a b c d
20. a b c d

21.

22.

23.

24.

CHEMISTRY

26. a b c d
27. a b c d
28. a b c d
29. a b c d
30. a b c d
31. a b c d
32. a b c d
33. a b c d
34. a b c d
35. a b c d
36. a b c d
37. a b c d
38. a b c d
39. a b c d
40. a b c d
41. a b c d
42. a b c d
43. a b c d
44. a b c d
45. a b c d

46.

47.

48.

49.

MATHEMATICS

51. a b c d
52. a b c d
53. a b c d
54. a b c d
55. a b c d
56. a b c d
57. a b c d
58. a b c d
59. a b c d
60. a b c d
61. a b c d
62. a b c d
63. a b c d
64. a b c d
65. a b c d
66. a b c d
67. a b c d
68. a b c d
69. a b c d
70. a b c d

71.

72.

73.

74.

INSTRUCTIONS

1. This test will be a 3 hours Test.
2. This test consists of Physics, Chemistry and Mathematics questions with equal weightage of 100 marks.
3. Each question is of 4 marks.
4. There are three sections in the question paper consisting of Physics (Q.no.1 to 25), Chemistry (Q.no.26 to 50) and Mathematics (Q. no.51 to 75). Each section is divided into two parts, Part I consists of 20 multiple choice questions & Part II consists of 5 Numerical value type Questions.
5. There will be only one correct choice in the given four choices in Part I. For each question 4 marks will be awarded for correct choice, 1 mark will be deducted for incorrect choice for Part I Questions and zero mark will be awarded for not attempted question. For Part II Questions 4 marks will be awarded for correct answer and zero for unattempted and incorrect answer.
6. Any textual, printed or written material, mobile phones, calculator etc. is not allowed for the students appearing for the test.
7. All calculations / written work should be done in the rough sheet provided.

PHYSICS

PART-I (Multiple Choice Questions)

1. A solid cylinder rolls down an inclined plane of height 3 m and reaches the bottom of plane with angular velocity of $2\sqrt{2}$ rad.s $^{-1}$. The radius of cylinder must be (Take $g=10\text{ ms}^{-2}$)
(a) 5 cm (b) 0.5 cm
(c) $\sqrt{10}\text{ cm}$ (d) $\sqrt{5}\text{ m}$
 2. In the figure shown, a particle of mass m is released from the position A on a smooth track. When the particle reaches at B, then normal reaction on it by the track is
-
- (a) mg (b) $2mg$
(c) $\frac{2}{3}mg$ (d) $\frac{m^2g}{h}$
3. The density ρ of water of bulk modulus B at a depth y in the ocean is related to the density at surface ρ_0 by the relation
(a) $\rho = \rho_0 \left[1 - \frac{\rho_0 gy}{B} \right]$

$$(b) \rho = \rho_0 \left[1 + \frac{\rho_0 g y}{B} \right]$$

$$(c) \rho = \rho_0 \left[1 + \frac{B}{\rho_0 h g y} \right]$$

$$(d) \rho = \rho_0 \left[1 - \frac{B}{\rho_0 h g y} \right]$$

4. The electric field in a certain region is given by $\vec{E} = (5\hat{i} - 3\hat{j}) kV/m$.

The potential difference $V_B - V_A$ between points A and B, having coordinates (4, 0, 3)m and (10, 3, 0)m respectively, is equal to

- (a) 21 kV (b) -21 kV
(c) 39 kV (d) -39 kV

5. Two electric bulbs marked 25W – 220 V and 100W – 220V are connected in series to a 440 V supply. Which of the bulbs will fuse?

- (a) Both (b) 100 W
(c) 25 W (d) Neither

6. Two long parallel wires P and Q are held perpendicular to the plane of the paper at a separation of 5 m. If P and Q carry currents of 2.5 A and 5 A respectively in the same direction, then the magnetic field at a point midway between P and Q is

(a) $\frac{\mu_0}{\pi}$ (b) $\sqrt{3} \frac{\mu_0}{\pi}$

(c) $\frac{\mu_0}{2\pi}$ (d) $\frac{3\mu_0}{2\pi}$

7. Two seconds after projection a projectile is travelling in a direction

inclined at 30° to the horizontal. After one more second, it is travelling horizontally. The magnitude and direction of its initial velocity are-


(a) $2\sqrt{20}$ m/s 60°

(b) $20\sqrt{3}$ m/s 60°

(c) $6\sqrt{40}$ m/s 30°

(d) $40\sqrt{6}$ m/s 30°

8. A 40 kg slab rests on a frictionless floor as shown in the figure. A 10 kg block rests on the top of the slab. The static coefficient of friction between the block and slab is 0.60 while the coefficient of kinetic friction is 0.40. The 10 kg block is acted upon by a horizontal force 100 N. If $g = 9.8$ m/s 2 , the resulting acceleration of the slab will be



- (a) 0.98 m/s^2 (b) 1.47 m/s^2
(c) 1.52 m/s^2 (d) 6.1 m/s^2

9. Two cars P and Q start from a point at the same time in a straight line and their positions are represented by $x_P(t) = at + bt^2$ and $x_Q(t) = ft - t^2$. At what time do the cars have the same velocity

(a) $\frac{f-a}{2(1+b)}$ (b) $\frac{a-1}{1+b}$

(c) $\frac{a+1}{2(b-1)}$ (d) $\frac{a+f}{2(1+b)}$

10. Ultraviolet light of wavelength 300 nm and intensity 1.0 watt/m^2 falls on the surface of a photosensitive material. If 1% of the incident photons produce photoelectrons, then find the number of photoelectrons emitted from an area of 1.0 cm^2 of the surface.

- (a) 9.61×10^{14} per sec
- (b) 4.12×10^{13} per sec
- (c) 1.51×10^{12} per sec
- (d) 2.13×10^{11} per sec





11. If the wavelength of the first line of the Balmer series of hydrogen is 6561 \AA , find the wavelength of the second line of the series.

- (a) 13122 \AA
- (b) 3280 \AA
- (c) 4860 \AA
- (d) 2187 \AA

12. The concentration of hole - electron pairs in pure silicon at $T = 300 \text{ K}$ is 7×10^{15} per cubic meter. Antimony is doped into silicon in a proportion of 1 atom in 10^7 Si atoms. Assuming that half of the impurity atoms contribute electron in the conduction band, calculate the factor by which the number of charge carriers increases due to doping. The number of silicon atoms per cubic meter is 5×10^{28}

- (a) 2.8×10^5
- (b) 3.1×10^2
- (c) 4.2×10^5
- (d) 1.8×10^5

13. Shown below are the black body radiation curves at temperatures T_1 and T_2 ($T_2 > T_1$). Which one of the following plots is correct?



14. A forced oscillator is acted upon by a force $F = F_0 \sin \omega t$. The amplitude of oscillation is given by

$$\frac{55}{\sqrt{2\omega^2 - 36\omega + 9}}.$$

The resonant angular frequency is


- (a) 2 units
- (b) 9 units
- (c) 18 units
- (d) 36 units

15. Three closed vessels A , B and C are at the same temperature T and contain gases which obey the Maxwellian distribution of velocities. Vessel A contains only O_2 , B only N_2 and C a mixture of

equal quantities of O_2 and N_2 . If the average speed of the O_2 molecules in vessel A is V_1 , that of the N_2 molecules in vessel B is V_2 , the average speed of the O_2 molecules in vessel C is


- (a) $(V_1 + V_2)/2$
- (b) V_1
- (c) $(V_1 V_2)^{1/2}$
- (d) $\sqrt{3kT/M}$

16. In the figure shown a source of sound of frequency 510 Hz moves with constant velocity $v_s = 20 \text{ m/s}$ in the direction shown. The wind is blowing at a constant velocity $v_w = 20 \text{ m/s}$ towards an observer who is at rest at point B. Corresponding to the sound emitted by the source at initial position A, the frequency detected by the observer is equal to (speed of sound relative to air = 330 m/s)



- (a) 510 Hz
- (b) 500 Hz
- (c) 525 Hz
- (d) 550 Hz

17. In fig, CODF is a semicircular loop of a conducting wire of resistance R and radius r. It is placed in a uniform magnetic field B, which is directed into the page (perpendicular to the plane of the loop).



The loop is rotated with a constant angular speed ω about an axis passing through the centre O, and perpendicular to the page. Then the induced current in the wire loop is

- (a) zero
- (b) $Br^2\omega/R$
- (c) $Br^2\omega/2R$
- (d) $B\pi r^2\omega/R$

18. If $E = 100 \sin(100t)$ volt and $I = 100$

$$\sin\left(100t + \frac{\pi}{3}\right) mA$$

are the instantaneous values of voltage and current, then the r.m.s. values of voltage and current are respectively

- (a) 70.7V, 70.7mA
- (b) 70.7V, 70.7A
- (c) 141.4V, 141.4mA
- (d) 141.4V, 141.4A

19. A plane electromagnetic wave is incident on a plane surface of area A, normally and is perfectly reflected. If energy E strikes the surface in time t then average pressure exerted on the surface is ($c = \text{speed of light}$)

- (a) zero
- (b) E/Atc
- (c) $2E/Atc$
- (d) E/c

- 20.** A 2.0 cm tall object is placed 15 cm in front of a concave mirror of focal length 10 cm. What is the size and nature of the image
- 4 cm, real
 - 4 cm, virtual
 - 1.0 cm, real
 - None of these

PART-II (Numerical Answer Questions)


- 21.** An inclined plane making an angle of 30° with the horizontal is placed in a uniform electric field of intensity 100 V/m. A particle of mass 1 kg and charge 0.01 C is allowed to slide down from rest on the plane from a height of 1 m. If the coefficient of friction is 0.2, then find the time taken (in second) by the particle to reach the bottom.

- 22.** A satellite is to be placed in equatorial geostationary orbit around earth for communication. The height (in metre) of such a satellite is

$$[M_E = 6 \times 10^{24} \text{ kg}, R_E = 6400 \text{ km}, T = 24 \text{ h}, G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}]$$

- 23.** A simple electric motor has an armature resistance of 1Ω and runs from a dc source of 12 volt. When running unloaded it draws a current of 2 amp. When a certain load is connected, its speed becomes one-half of its unloaded value. What is the new value of current drawn (in ampere)?

- 24.** A gas can be taken from A to B via two different processes ACB and ADB.



- When path ACB is used 60 J of heat flows into the system and 30J of work is done by the system. If path ADB is used work done by the system is 10 J. The heat flow (in joule) into the system in path ADB is :
- 25.** If 200 MeV energy is released per fission of U^{235} nuclei. Find the mass of U^{235} consumed (in mg) per day in a reactor of power 1MW assuming its efficiency is 80%.

CHEMISTRY

PART-I (Multiple Choice Questions)

- 26.** The reason for almost doubling the rate of reaction on increasing the temperature of the reaction system by 10°C is
- The value of threshold energy increases
 - Collision frequency increases
 - The fraction of the molecule having energy equal to threshold energy or more increases
 - Activation energy decreases

27. Which of the following factors may be regarded as the main cause of lanthanoid contraction?
- Greater shielding of $5d$ electrons by $4f$ electrons.
 - Poorer shielding of $5d$ electrons by $4f$ electrons.
 - Effective shielding of one of $4f$ electrons by another in the subshell.
 - Poor shielding of one of $4f$ electron by another in the subshell.
28. Isobutyl magnesium bromide with dry ether and ethyl alcohol gives :
- $\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{OH}$ & $\text{CH}_3\text{CH}_2\text{MgBr}$
 - $\text{CH}_3\text{CH}(\text{CH}_3)_2$ & $\text{MgBr}(\text{OC}_2\text{H}_5)_2$
 - $\text{CH}_3\text{CH}(\text{CH}_3)=\text{CH}_2$ & $\text{Mg}(\text{OH})\text{Br}$
 - $\text{CH}_3\text{CH}(\text{CH}_3)_2$ & $\text{CH}_3\text{CH}_2\text{OMgBr}$
29. The K_p/K_c ratio will be highest in case of
- $\text{CO}(g) + \frac{1}{2}\text{O}_2(g) \rightleftharpoons \text{CO}_2(g)$
 - $\text{H}_2(g) + \text{I}_2(g) \rightleftharpoons 2\text{HI}(g)$
 - $\text{PCl}_5(g) \rightleftharpoons \text{PCl}_3(g) + \text{Cl}_2(g)$
 - $7\text{H}_2(g) + 2\text{NO}_2(g) \rightleftharpoons 2\text{NH}_3(g) + 4\text{H}_2\text{O}(g)$
30. Which of the following substances has the greatest ionic character ?
- Cl_2O
 - NCl_3
 - PbCl_2
 - BaCl_2
31. An organic compound contains 49.3% carbon, 6.84% hydrogen and its vapour density is 73. Molecular formula of the compound is :
- $\text{C}_3\text{H}_5\text{O}_2$
 - $\text{C}_4\text{H}_{10}\text{O}_2$
 - $\text{C}_6\text{H}_{10}\text{O}_4$
 - $\text{C}_3\text{H}_{10}\text{O}_2$
32. Consider the following transformations :
- $$\text{CH}_3\text{COOH} \xrightarrow{\text{CaCO}_3} \text{A} \xrightarrow{\text{heat}} \text{B} \xrightarrow[\text{NaOH}]{\text{I}_2} \text{C}$$
- The molecular formula of C is
- $\text{CH}_3-\overset{\text{OH}}{\underset{\text{I}}{|}\text{C}-\text{CH}_3}$
 - $\text{ICH}_2-\text{COCH}_3$
 - CHI_3
 - CH_3I
33. The values of ΔH and ΔS for the reaction,
- $$\text{C(graphite)} + \text{CO}_2(g) \rightarrow 2\text{CO}(g)$$
- are 170 kJ and 170 JK^{-1} , respectively. This reaction will be spontaneous at
- 910 K
 - 1110 K
 - 510 K
 - 710 K
34. Containers A and B have same gases. Pressure, volume and temperature of A are all twice that





of B, then the ratio of number of molecules of A and B are

- (a) 1:2 (b) 2:1
(c) 1:4 (d) 4:1

35. The stability of +1 oxidation state among Al, Ga, In and Tl increases in the sequence :

- (a) Ga < In < Al < Tl
(b) Al < Ga < In < Tl
(c) Tl < In < Ga < Al
(d) In < Tl < Ga < Al

36. Among the following four structures I to IV,



it is true that

- (a) only I and II are chiral compounds
(b) only III is a chiral compound
(c) only II and IV are chiral compounds
(d) all four are chiral compounds

37. Which is a dangerous radiological pollutant?

- (a) C^{14} (b) S^{35}
(c) Sr^{90} (d) P^{32}

38. A compound of formula A_2B_3 has the *hcp* lattice. Which atom forms the *hcp* lattice and what fraction of tetrahedral voids is occupied by the other atoms:

- (a) *hcp* lattice – A, $\frac{2}{3}$
Tetrahedral voids – B

- (b) *hcp* lattice – A, $\frac{1}{3}$
Tetrahedral voids – B

- (c) *hcp* lattice – B, $\frac{2}{3}$
Tetrahedral voids – A

- (d) *hcp* lattice – B, $\frac{1}{3}$
Tetrahedral voids – A

39. Of the four isomeric hexanes, the isomer which can give two monochlorinated compounds is

- (a) 2-methylpentane
(b) 2, 2-dimethylbutane
(c) 2, 3-dimethylbutane
(d) n-hexane


40. The solubility product of PbCl_2 is 1.7×10^{-5} . The solubility in moles per litre would be :

- (a) 1.62×10^{-4} (b) 1.62×10^{-8}
(c) 1.62×10^{-2} (d) 1.62×10^{-6}

41. Crystal field stabilization energy for high spin d^4 octahedral complex is:

- (a) $-1.8 \Delta_0$ (b) $-1.6 \Delta_0 + P$
(c) $-1.2 \Delta_0$ (d) $-0.6 \Delta_0$

42. In the reaction sequence



the product 'C' is:

- (a) benzonitrile
- (b) benzaldehyde
- (c) benzoic acid
- (d) benzylamine

43. Nylon threads are made of

- (a) polyester polymer
- (b) polyamide polymer
- (c) polyethylene polymer
- (d) polyvinyl polymer

44. In the reaction of oxalate with permanganate in acidic medium, the number of electrons involved in producing one molecule of CO_2 is:


- (a) 1
- (b) 10
- (c) 2
- (d) 5

45. Momentum of radiations of wavelength 0.33 nm is :

- (a) $2.01 \times 10^{-21} \text{ kg m sec}^{-1}$
- (b) $2.01 \times 10^{-24} \text{ g m sec}^{-1}$
- (c) $2.01 \times 10^{-21} \text{ g m sec}^{-1}$
- (d) $2.01 \times 10^{-24} \text{ kg m sec}^{-1}$

PART-II (Numerical Answer Questions)

46. Calculate the difference in the heat of formation (in calories) of carbon monoxide at constant pressure and at constant volume at 27°C .
47. Calculate number of molecules of Grignard reagent consumed by 1 molecule of following compound.



48. In a metal oxide, there is 20% oxygen by weight. What is its equivalent weight?

49. Find the total number of possible isomers for the complex compound $[\text{Cu}^{\text{II}}(\text{NH}_3)_4][\text{Pt}^{\text{II}}\text{Cl}_4]$

50. Calculate the strength in % of labelled 10 volume H_2O_2 solution.

MATHEMATICS

PART-I (Multiple Choice Questions)

51. If $(7 - 4\sqrt{3})^{x^2 - 4x + 3} +$

$$(7 + 4\sqrt{3})^{x^2 - 4x + 3} = 14,$$

then the value of x is given by

- (a) $2, 2 \pm \sqrt{2}$
- (b) $2 \pm \sqrt{3}, 3$
- (c) $3 \pm \sqrt{2}, 2$
- (d) None of these

52. The minimum value of the function

$$f(x) = x^{3/2} + x^{-3/2} - 4\left(x + \frac{1}{x}\right) \text{ for}$$

all permissible real x , is

- (a) -10
- (b) -6
- (c) -7
- (d) -8

53. In the expansion of $\left(\frac{x}{2} - \frac{3}{x^2}\right)^{10}$,
the coefficient of x^4 is
(a) $\frac{405}{256}$ (b) $\frac{504}{259}$
(c) $\frac{450}{263}$
(d) None of these
54. If the plane $3x + y + 2z + 6 = 0$ is parallel to the line
 $\frac{3x-1}{2b} = 3-y = \frac{z-1}{a}$, then the value of $3a+3b$ is
(a) $\frac{1}{2}$ (b) $\frac{3}{2}$
(c) 3 (d) 4
55. The domain of definition of the function
 $f(x) = \sqrt{1 + \log_e(1-x)}$ is
(a) $-\infty < x \leq 0$
(b) $-\infty < x \leq \frac{e-1}{e}$
(c) $-\infty < x \leq 1$
(d) $x \geq 1-e$
56. The function $f(x) = [x]^2 - [x^2]$ (where $[y]$ is the greatest integer less than or equal to y), is discontinuous at
(a) All integers
(b) All integers except 0 and 1
(c) All integers except 0
(d) All integers except 1
57. The line $y = mx$ bisects the area enclosed by lines $x=0$, $y=0$ and $x=3/2$ and the curve $y=1+4x-x^2$. Then the value of m is
(a) $\frac{3}{5}\log 2$ (b) $\frac{2}{5}\log 2$
(c) $-\frac{3}{2}\log 2$
(d) None of these
58. The sum of the series
 $3+33+333+\dots+n$ terms is
(a) $\frac{1}{27}(10^{n+1}+9n-28)$
(b) $\frac{1}{27}(10^{n+1}-9n-10)$
(c) $\frac{1}{27}(10^{n+1}+10n-9)$
(d) None of these
59. If $\int \frac{1}{1+\sin x} dx = \tan\left(\frac{x}{2} + a\right) + b$ then
(a) $a = -\frac{\pi}{4}$, $b \in \mathbf{R}$
(b) $a = \frac{\pi}{4}$, $b \in \mathbf{R}$
(c) $a = \frac{5\pi}{4}$, $b \in \mathbf{R}$
(d) None of these
60. If $y = \tan^{-1}\left(\frac{2^x}{1+2^{2x+1}}\right)$, then
 $\frac{dy}{dx}$ at $x=0$ is
(a) $\frac{3}{5}\log 2$ (b) $\frac{2}{5}\log 2$
(c) $-\frac{3}{2}\log 2$
(d) None of these

61. The value of

$$\cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{6\pi}{7}$$
 is

- (a) 0 (b) 1
(c) $\frac{1}{2}$ (d) $-\frac{1}{2}$

62. An integrating factor of the differential equation

$$\frac{dy}{dx} = y \tan x - y^2 \sec x$$
 is equal to:

- (a) $\tan x$ (b) $\sec x$
(c) cosec x (d) $\cot x$

63. If $y = 2x$ is a chord of the circle $x^2 + y^2 = 10x$, then the equation of the circle whose diameter is this chord, is -

- (a) $x^2 + y^2 + 2x + 4y = 0$
(b) $x^2 + y^2 + 2x - 4y = 0$
(c) $x^2 + y^2 - 2x - 4y = 0$
(d) None of these

64. Magnitudes of vectors $\vec{a}, \vec{b}, \vec{c}$ are 3, 4, 5 respectively. If \vec{a} and $\vec{b} + \vec{c}$, \vec{b} and $\vec{c} + \vec{a}$, \vec{c} and $\vec{a} + \vec{b}$ are mutually perpendicular, then

magnitude of $\vec{a} + \vec{b} + \vec{c}$ is

- (a) $4\sqrt{2}$ (b) $3\sqrt{2}$
(c) $5\sqrt{2}$ (d) $3\sqrt{3}$

65. If $a + b + c = 0$, then the solution of the equation

$$\begin{vmatrix} a-x & c & b \\ c & b-x & a \\ b & a & c-x \end{vmatrix} = 0$$
 is

(a) 0

(b) $\pm \frac{3}{2}(a^2 + b^2 + c^2)$

(c) $0, \pm \sqrt{\frac{3}{2}(a^2 + b^2 + c^2)}$

(d) $0, \pm \sqrt{(a^2 + b^2 + c^2)}$

66. If $I_1 = \int_0^1 2x^2 dx$, $I_2 = \int_0^1 2x^3 dx$,

$$I_3 = \int_1^2 2x^2 dx \text{ and } I_4 = \int_1^2 2x^3 dx$$

then

- (a) $I_2 > I_1$ (b) $I_1 > I_2$
(c) $I_3 = I_4$ (d) $I_3 > I_4$

67. If $f : R \rightarrow R$ and $g : R \rightarrow R$ are defined by $f(x) = |x|$ and $g(x) = [x - 3]$ for $x \in R$, then

$\left\{ g(f(x)) : -\frac{8}{5} < x < \frac{8}{5} \right\}$ is equal to

- (a) $\{0, 1\}$ (b) $\{1, 2\}$
(c) $\{-3, -2\}$ (d) $\{2, 3\}$

68. If A and B are two events such that

$$P(A) = \frac{1}{2} \text{ and } P(B) = \frac{2}{3}, \text{ then}$$

(a) $P(A \cup B) \geq \frac{2}{3}$

(b) $\frac{1}{6} \leq P(A \cap B) \leq \frac{1}{2}$

(c) $\frac{1}{6} \leq P(A' \cap B) \leq \frac{1}{2}$

- (d) All of these

69. If PQ is a double ordinate of

hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ such that

OPQ is an equilateral triangle, O being the centre of the hyperbola. Then the eccentricity e of the hyperbola satisfies

- (a) $1 < e < 2/\sqrt{3}$ (b) $e = 2/\sqrt{3}$
(c) $e = \sqrt{3}/2$ (d) $e > 2/\sqrt{3}$

70. The equation of the lines on which the perpendiculars from the origin make 30° angle with x -axis and

which form a triangle of area $\frac{50}{\sqrt{3}}$

with axes, are

- (a) $x + \sqrt{3}y \pm 10 = 0$
(b) $\sqrt{3}x + y \pm 10 = 0$
(c) $x \pm \sqrt{3}y - 10 = 0$
(d) None of these

PART-II (Numerical Answer Questions)

71. The number of pairs (x, y) satisfying the equations $\sin x + \sin y = \sin(x+y)$ and $|x| + |y| = 1$ is

72. The value of

$$\lim_{x \rightarrow 0} \left\{ \frac{\sin x - x + \frac{x^3}{6}}{x^5} \right\} \text{ is } \frac{1}{k},$$

then k is

73. An edge of a variable cube is increasing at the rate cm/sec. Then, the state of increase in volume of the cube when the edge is 5 cm long, is

74. If $2x = -1 + \sqrt{3}i$, then the value of $(1 - x^2 + x)^6 - (1 - x + x^2)^6$ is

75. Sum of all three digit numbers (no digit being zero) having the property that all digits are perfect squares, is

RESPONSE SHEET

PHYSICS

1. (a) (b) (c) (d)
2. (a) (b) (c) (d)
3. (a) (b) (c) (d)
4. (a) (b) (c) (d)
5. (a) (b) (c) (d)
6. (a) (b) (c) (d)
7. (a) (b) (c) (d)
8. (a) (b) (c) (d)
9. (a) (b) (c) (d)
10. (a) (b) (c) (d)
11. (a) (b) (c) (d)
12. (a) (b) (c) (d)
13. (a) (b) (c) (d)
14. (a) (b) (c) (d)
15. (a) (b) (c) (d)
16. (a) (b) (c) (d)
17. (a) (b) (c) (d)
18. (a) (b) (c) (d)
19. (a) (b) (c) (d)
20. (a) (b) (c) (d)

21.

22.

23.

24.

CHEMISTRY

26. (a) (b) (c) (d)
27. (a) (b) (c) (d)
28. (a) (b) (c) (d)
29. (a) (b) (c) (d)
30. (a) (b) (c) (d)
31. (a) (b) (c) (d)
32. (a) (b) (c) (d)
33. (a) (b) (c) (d)
34. (a) (b) (c) (d)
35. (a) (b) (c) (d)
36. (a) (b) (c) (d)
37. (a) (b) (c) (d)
38. (a) (b) (c) (d)
39. (a) (b) (c) (d)
40. (a) (b) (c) (d)
41. (a) (b) (c) (d)
42. (a) (b) (c) (d)
43. (a) (b) (c) (d)
44. (a) (b) (c) (d)
45. (a) (b) (c) (d)

46.

47.

48.

49.

MATHEMATICS

51. (a) (b) (c) (d)
52. (a) (b) (c) (d)
53. (a) (b) (c) (d)
54. (a) (b) (c) (d)
55. (a) (b) (c) (d)
56. (a) (b) (c) (d)
57. (a) (b) (c) (d)
58. (a) (b) (c) (d)
59. (a) (b) (c) (d)
60. (a) (b) (c) (d)
61. (a) (b) (c) (d)
62. (a) (b) (c) (d)
63. (a) (b) (c) (d)
64. (a) (b) (c) (d)
65. (a) (b) (c) (d)
66. (a) (b) (c) (d)
67. (a) (b) (c) (d)
68. (a) (b) (c) (d)
69. (a) (b) (c) (d)
70. (a) (b) (c) (d)

71.

72.

73.

74.

MOCK TEST

5

INSTRUCTIONS

1. This test will be a 3 hours Test.
2. This test consists of Physics, Chemistry and Mathematics questions with equal weightage of 100 marks.
3. Each question is of 4 marks.
4. There are three sections in the question paper consisting of Physics (Q.no.1 to 25), Chemistry (Q.no.26 to 50) and Mathematics (Q. no.51 to 75). Each section is divided into two parts, Part I consists of 20 multiple choice questions & Part II consists of 5 Numerical value type Questions.
5. There will be only one correct choice in the given four choices in Part I. For each question 4 marks will be awarded for correct choice, 1 mark will be deducted for incorrect choice for Part I Questions and zero mark will be awarded for not attempted question. For Part II Questions 4 marks will be awarded for correct answer and zero for unattempted and incorrect answer.
6. Any textual, printed or written material, mobile phones, calculator etc. is not allowed for the students appearing for the test.
7. All calculations / written work should be done in the rough sheet provided.

PHYSICS

PART-I (Multiple Choice Questions)

1. A spherical ball A of mass 4 kg , moving along a straight line strikes another spherical ball B of mass 1 kg at rest. After the collision, A and B move with velocities $v_1 \text{ ms}^{-1}$ and $v_2 \text{ ms}^{-1}$ respectively making angles of 30° and 60° with respect to the original direction of motion of A.


The ratio $\frac{v_1}{v_2}$ will be

(a) $\sqrt{3}/4$ (b) $4/\sqrt{3}$

(c) $1/\sqrt{3}$ (d) $\sqrt{3}$

2. A thin wire of length L and uniform linear mass density ρ is bent into a circular loop with centre at O as shown. The moment of inertia of the loop about the axis XX' is

X ————— XX'







(a) $\frac{\rho L^3}{8\pi^2}$

(b) $\frac{\rho L^3}{16\pi^2}$

(c) $\frac{5\rho L^3}{16\pi^2}$






(d) $\frac{3\rho L^3}{8\pi^2}$

3. The diagram showing the variation of gravitational potential of earth with distance from the centre of earth is




4. Two thin dielectric slabs of dielectric constants K_1 and K_2 ($K_1 < K_2$) are inserted between plates of a parallel plate capacitor, as shown in the figure. The variation

of electric field ‘ E ’ between the plates with distance ‘ d ’ as measured from plate P is correctly shown by :



5. Two identical capacitors having plate separation d_0 are connected parallel to each other across points A and B as shown in figure. A charge Q is imparted to the





system by connecting a battery across A and B and battery is removed. Now first plate of first capacitor and second plate of second capacitor starts moving with constant velocity u_0 towards left. Find the magnitude of current flowing in the loop during the process.



(a) $\frac{Q}{2d_0}u_0$ (b) $\frac{Q}{d_0}u_0$

(c) $\frac{2Q}{d_0}u_0$ (d) $\frac{Q}{3d_0}u_0$

6. Two long parallel wires are at a distance $2d$ apart. They carry steady equal currents flowing out of the plane of the paper as shown. The variation of the magnetic field B along the line XX' is given by



7. If dimensions of critical velocity v_c of a liquid flowing through a tube are expressed as $[\eta^\alpha \rho^\beta r^\gamma]$, where η , ρ and r are the coefficient of viscosity of liquid, density of liquid and radius of the tube respectively, then the values of x , y and z are given by :

(a) -1, -1, 1 (b) -1, -1, -1
 (c) 1, 1, 1 (d) 1, -1, -1

8. A car accelerates from rest at a constant rate α for some time, after which it decelerates at a constant rate β and comes to rest. If the total time elapsed is t , then the maximum velocity acquired by the car is

(a) $\left(\frac{\alpha^2 + \beta^2}{\alpha\beta} \right)t$

(b) $\left(\frac{\alpha^2 - \beta^2}{\alpha\beta} \right)t$

(c) $\frac{(\alpha + \beta)t}{\alpha\beta}$

(d) $\frac{\alpha\beta t}{\alpha + \beta}$

9. The speed of a projectile at its maximum height is $\frac{\sqrt{3}}{2}$ times its initial speed. If the range of the projectile is 'P' times the maximum height attained by it. P is-




- (a) $\frac{4}{3}$ (b) $2\sqrt{3}$
 (c) $4\sqrt{3}$ (d) $\frac{3}{4}$

10. All electrons ejected from a surface by incident light of wavelength 200nm can be stopped before travelling 1m in the direction of uniform electric field of 4N/C. The work function of the surface is
 (a) 4eV (b) 6.2eV
 (c) 2eV (d) 2.2eV





11. Find the ratio of longest wavelength and the shortest wavelength observed in the five spectral series of emission spectrum of hydrogen.

- (a) $\frac{4}{3}$ (b) $\frac{525}{376}$
 (c) 25 (d) $\frac{900}{11}$


12. The energy spectrum of β -particles [Number N(E) as a function of β -energy E] emitted from a radioactive source is




13. If a piece of metal is heated to temperature θ and then allowed to cool in a room which is at temperature θ_0 , the graph between the temperature T of the metal and time t will be closest to



14. 'n' moles of an ideal gas undergoes a process A \rightarrow B as shown in the figure. The maximum temperature of the gas during the process will be :




- (a) $\frac{9P_0 V_0}{2nR}$ (b) $\frac{9P_0 V_0}{nR}$
 (c) $\frac{9P_0 V_0}{4nR}$ (d) $\frac{3P_0 V_0}{2nR}$
15. In the given (V-T) diagram, what is the relation between pressure P_1 and P_2 ?



- (a) $P_2 > P_1$
 (b) $P_2 < P_1$
 (c) Cannot be predicted
 (d) $P_2 = P_1$
16. A particle executes simple harmonic motion with a time period of 16s. At time $t = 2\text{s}$, the particle crosses the mean position while at $t = 4\text{s}$, its velocity is 4 m/s^{-1} . The amplitude of motion in metre is


- (a) $\sqrt{2}\pi$ (b) $16\sqrt{2}\pi$
 (c) $24\sqrt{2}\pi$ (d) $\frac{32\sqrt{2}}{\pi}$

17. A metallic rod of length ' ℓ ' is tied to a string of length 2ℓ and made to rotate with angular speed ω on a horizontal table with one end of the string fixed. If there is a vertical magnetic field 'B' in the region, the e.m.f. induced across the ends of the rod is



- (a) $\frac{2B\omega\ell^2}{2}$ (b) $\frac{3B\omega\ell^2}{2}$
 (c) $\frac{4B\omega\ell^2}{2}$ (d) $\frac{5B\omega\ell^2}{2}$


18. In an LCR circuit as shown below both switches S_1 and S_2 are open initially. Now switch S_1 is closed, S_2 kept open. (q is charge on the capacitor and $\tau = RC$ is capacitive time constant). Which of the following statements is correct ?



- (a) Work done by the battery is half of the energy dissipated in the resistor

- (b) At $t = \tau$, $q = CV/2$
- (c) At $t = 2\tau$, $q = CV(1 - e^{-2})$
- (d) At $t = \frac{\tau}{2}$, $q = CV(1 - e^{-1})$


19. An electromagnetic wave in vacuum has the electric and magnetic field \vec{E} and \vec{B} , which are always perpendicular to each other. The direction of polarization is given by \vec{X} and that of wave propagation by \vec{k} . Then
- (a) $\vec{X} \parallel \vec{B}$ and $\vec{k} \parallel \vec{B} \times \vec{E}$
 - (b) $\vec{X} \parallel \vec{E}$ and $\vec{k} \parallel \vec{E} \times \vec{B}$
 - (c) $\vec{X} \parallel \vec{B}$ and $\vec{k} \parallel \vec{E} \times \vec{B}$
 - (d) $\vec{X} \parallel \vec{E}$ and $\vec{k} \parallel \vec{B} \times \vec{E}$
20. Two blocks each of mass m lie on a smooth table. They are attached to two other masses as shown in the figure. The pulleys and strings are light. An object O is kept at rest on the table. The sides AB and CD of the two blocks are made reflecting. The acceleration of two images formed in these two reflecting surfaces w.r.t. each other is $17g/A$ then find the value of



- (a) 2
- (b) 6
- (c) 4
- (d) 5

PART-II (Numerical Answer Questions)


21. Combination of two identical capacitors, a resistor R and a DC voltage source of voltage 6 V is used in an experiment on C-R circuit. It is found that for a parallel combination of the capacitor the time in which the voltage of the fully charged combination reduces to half its original voltage is 10 s. For series combination the time (in sec) needed for reducing the voltage of the fully charged series combination by half is
22. Escape velocity for earth surface is 11 km/s. If the radius of any planet is two times the radius of the earth but average density is same as that of earth. Then the escape velocity (in km/s) at the planet will be
23. Two identical glass rods S_1 and S_2 (refractive index = 1.5) have one convex end of radius of curvature 10 cm. They are placed with the curved surfaces at a distance d as shown in the figure, with their axes (shown by the dashed line) aligned. When a point source of light P is placed inside rod S_1 on its axis at a distance of 50 cm from the curved face, the light rays emanating from it are found to be parallel to the axis inside S_2 . The distance d (in cm) is




24. The displacement of a particle executing SHM is given by
 $y = 5 \sin\left(4t + \frac{\pi}{3}\right)$. If T is the time period and mass of the particle is 2g, the kinetic energy (in joule) of the particle when $t = \frac{T}{4}$ is given by
25. A zener diode of voltage $V_Z (= 6V)$ is used to maintain a constant voltage across a load resistance $R_L (= 1000 \Omega)$ by using a series resistance $R_s (= 100\Omega)$. If the e.m.f. of source is E ($= 9 V$), what is the power (in watt) being dissipated in Zener diode ?

CHEMISTRY

PART-I (Multiple Choice Questions)

26. The compound that does not produce nitrogen gas by the thermal decomposition is :
 (a) $\text{Ba}(\text{N}_3)_2$ (b) $(\text{NH}_4)_2\text{Cr}_2\text{O}_7$
 (c) NH_4NO_2 (d) $(\text{NH}_4)_2\text{SO}_4$
27. The following compound is used as
- 
- (a) an anti-inflammatory compound
 (b) analgesic
 (c) hypnotic
 (d) antiseptic
28. In a reversible reaction the energy of activation of the forward reaction is 50 kcal. The energy of activation for the reverse reaction will be
 (a) < 50 kcal
 (b) either greater than or less than 50 kcal
 (c) 50 kcal
 (d) > 50 kcal
29. The method not used in metallurgy to refine the impure metal is
 (a) Mond's process
 (b) van-Arkel process
 (c) Amalgamation process
 (d) Liquation
30. When PbO_2 reacts with conc. HNO_3 the gas evolved is
 (a) NO_2 (b) O_2
 (c) N_2 (d) N_2O
31. Soap helps in cleaning clothes, because
 (a) chemical of soap change
 (b) it increases the surface tension of the solution
 (c) it absorbs the dirt
 (d) it lowers the surface tension of the solution
32. Orthoboric acid –
 (a) donate proton to form H_2BO_3^-
 (b) accept proton of form H_4BO_3^+
 (c) donate OH^- to form H_2BO_2^+
 (d) accept OH^- to form $[\text{B}(\text{OH})_4]^-$

33. Consider the reaction



The equilibrium constant of the above reaction is K_p . If pure ammonia is left to dissociate, the partial pressure of ammonia at equilibrium is given by (Assume that $P_{\text{NH}_3} \ll P_{\text{total}}$ at equilibrium)

(a) $\frac{3^{3/2} K_p^{1/2} P^2}{16}$

(b) $\frac{K_p^{1/2} P^2}{16}$

(c) $\frac{K_p^{1/2} P^2}{4}$

(d) $\frac{3^{3/2} K_p^{1/2} P^2}{4}$

34. The property which distinguishes formic acid from acetic acid is

- (a) only ammonium salt of formic acid on heating gives amide.
(b) when heated with alcohol/ H_2SO_4 only acetic acid forms ester.
(c) only acetic acid forms salts with alkali.
(d) only formic acid reduces Fehling's solution.

35. The standard emf of a cell, involving one electron change is found to be 0.591 V at 25°C. The equilibrium constant of the reaction is ($F = 96500 \text{ C mol}^{-1}$)

- (a) 1.0×10^1 (b) 1.0×10^5

- (c) 1.0×10^{10} (d) 1.0×10^{30}

36. Calomel (Hg_2Cl_2) on reaction with ammonium hydroxide gives

- (a) HgO

- (b) Hg_2O

- (c) $\text{NH}_2 - \text{Hg} - \text{Hg} - \text{Cl}$

- (d) $\text{Hg}_2\text{NH}_2\text{Cl}$

37. If one strand of DNA has the sequence ATGCTTGA, the sequence in the complimentary strand would be

- (a) TCCGAACT

- (b) TACGTAGT

- (c) TACGAACT

- (d) TAGCTAGT

38. Bromination of toluene gives

- (a) only *m*-substituted product

- (b) only *p*-substituted product

- (c) mixture of *o*-and *p*-substituted products

- (d) mixture of *o*-and *m*-substituted products

39. In sodium fusion test of organic compounds, the nitrogen of the organic compound is converted into


- (a) sodamide

- (b) sodium cyanide

- (c) sodium nitrite

- (d) sodium nitrate

40. What happens when magnesium is burnt in air and the products X and Y are treated with water?



- | | X | Y | P | Q |
|-----|----------|--------------------------------|---------------------|-----------------|
| (a) | MgO | Mg(OH) ₂ | Mg(OH) ₂ | N ₂ |
| (b) | MgO | Mg ₃ N ₂ | Mg(OH) ₂ | NH ₃ |
| (c) | MgO | Mg ₃ N ₂ | Mg(OH) ₂ | N ₂ |
| (d) | MgO | MgCO ₃ | Mg(OH) ₂ | CO ₂ |

41. When a small quantity of FeCl₃ solution is added to the fresh precipitate of Fe(OH)₃, a colloidal sol is obtained.

The process through which this sol is formed is known as

- (a) exchange of solvent
- (b) chemical double decomposition
- (c) peptization
- (d) electrophoresis


42. The molal elevation constant of water = 0.52 °C kg mol⁻¹. The boiling point of 1.0 molal aqueous KCl solution (assuming complete dissociation of KCl), therefore should be

- (a) 100.52 °C
- (b) 101.04 °C
- (c) 99.48 °C
- (d) 98.96 °C


43. The oxidation state of Cr in [Cr(NH₃)₄Cl₂]⁺ is

- (a) 0
- (b) +1
- (c) +2
- (d) +3


44. Standard reduction potentials of the half reactions are given below :



$$E^\circ = +2.85 \text{ V}$$



$$E^\circ = +1.36 \text{ V}$$




$$E^\circ = +1.06 \text{ V}$$



The strongest oxidising and reducing agents respectively are :

- (a) F₂ and I⁻
- (b) Br₂ and Cl⁻
- (c) Cl₂ and Br⁻
- (d) Cl₂ and I₂

45. Cyclopropane rearranges to form propene



This follows first order kinetics. The rate constant is $2.714 \times 10^{-3} \text{ s}^{-1}$. The initial concentration of cyclopropane is 0.29 M. What will be the concentration of cyclopropane after 100 s?


- (a) 0.035 M
- (b) 0.22 M
- (c) 0.145 M
- (d) 0.0018 M

PART-II (Numerical Answer Questions)

46. What is the order of reaction of the formation of gas at the surface of tungsten due to adsorption?

47. 1.0 g of metal nitrate gave 0.86 g of metal sulphate. Calculate equivalent wt. of metal in grams.

48. How many isomeric naphthylamines are expected in the following reaction?



49. At infinite dilution, the molar conductance of Ba^{2+} and Cl^- are 127 and 76 $\text{S cm}^2 \text{ mol}^{-1}$. What is the molar conductivity of BaCl_2 at indefinite dilution?
50. The enthalpy of hydrogenation of cyclohexene is $-119.5 \text{ kJ mol}^{-1}$. If resonance energy of benzene is $-150.4 \text{ kJ mol}^{-1}$, calculate its enthalpy of hydrogenation in kJ.

MATHEMATICS

PART-I (Multiple Choice Questions)

51. If one root is square of the other root of the equation $x^2 + px + q = 0$, then the relation between p and q is

- (a) $p^3 - (3p - 1)q + q^2 = 0$
- (b) $p^3 - q(3p + 1) + q^2 = 0$
- (c) $p^3 + q(3p - 1) + q^2 = 0$
- (d) $p^3 + q(3p + 1) + q^2 = 0$

52. A chord AB drawn from the point $A(0, 3)$ on circle $x^2 + 4x + (y - 3)^2 = 0$ meets to M in such a way that $AM = 2AB$, then the locus of point M will be

- (a) Straight line
- (b) Circle
- (c) Parabola
- (d) None of these

53. Let $f(x) = \begin{cases} (x-1)\sin \frac{1}{x-1} & \text{if } x \neq 1 \\ 0 & \text{if } x = 1 \end{cases}$

Then which one of the following is true?

- (a) f is differentiable at $x = 0$ and $x = 1$
- (b) f is differentiable at $x = 0$ but not at $x = 1$
- (c) f is differentiable at $x = 1$ but not at $x = 0$
- (d) f is neither differentiable at $x = 0$ nor at $x = 1$

54. In a town of 10,000 families it was found that 40% family buy newspaper A, 20% buy newspaper B and 10% families buy newspaper C, 5% families buy A and B, 3% buy B and C and 4% buy A and C. If 2% families buy all the three newspapers, then number of families which buy A only is
- (a) 3100
 - (b) 3300
 - (c) 2900
 - (d) 1400

55. The numbers P, Q and R for which the function

$f(x) = \text{Pe}^{2x} + \text{Qe}^x + \text{Rx}$ satisfies the conditions

$$f(0) = -1, f'(\log 2) = 31 \text{ and}$$

$$\int_0^{\log 4} [f(x) - Rx] dx = \frac{39}{2}$$

are given by

- (a) P=2, Q=-3, R=4
 (b) P=-5, Q=2, R=3
 (c) P=5, Q=-2, R=3
 (d) P=5, Q=-6, R=3

- 56.** The value of $\lim_{x \rightarrow 0^+} x^m (\log x)^n$, m,

n ∈ N is

- (a) 0
 (b) $\frac{m}{n}$
 (c) mn
 (d) None of these

- 57.** The value of a in order that $f(x) = \sin x - \cos x - ax + b$ decreases for all real values is given by

- (a) $a \geq \sqrt{2}$ (b) $a < \sqrt{2}$
 (c) $a \geq 1$ (d) $a < 1$

- 58.** If in ΔABC , $2b^2 = a^2 + c^2$, then

$$\frac{\sin 3B}{\sin B} =$$

(a) $\frac{c^2 - a^2}{2ca}$

(b) $\frac{c^2 - a^2}{ca}$

(c) $\left(\frac{c^2 - a^2}{ca} \right)^2$

(d) $\left(\frac{c^2 - a^2}{2ca} \right)^2$

- 59.** The equation of the normal to the curve

$$y = (1+x)^y + \sin^{-1}(\sin^2 x) \text{ at}$$

x=0 is

- (a) $x+y=1$
 (b) $x+y+1=0$
 (c) $2x-y+1=0$
 (d) $x+2y+2=0$

- 60.** If a circles $x^2 + y^2 = a^2$ and the rectangular hyperbola $xy = c^2$

intersect in four points, $\left(ct_r, \frac{c}{t_r} \right)$, $r=1, 2, 3, 4$ then $t_1 t_2 t_3 t_4$ is equal to

- (a) -1 (b) 1
 (c) c^4 (d) $-c^4$

- 61.** $\int (32x^3 (\log x)^2 dx)$ is equal to :

- (a) $8x^4 (\log x)^2 + C$
 (b) $x^4 \{8(\log x)^2 - 4(\log x) + 1\} + C$
 (c) $x^4 \{8(\log x)^2 - 4(\log x)\} + C$
 (d) $x^3 \{(\log x)^2 - 2\log x\} + C$

- 62.** Differential coefficient of

$$\tan^{-1} \frac{2x}{1-x^2} \text{ with respect to } \sin^{-1} \frac{2x}{1+x^2} \text{ will be}$$

- (a) 1 (b) -1
 (c) $-1/2$ (d) x

- 63.** The area of the plane region bounded by the curves $x+2y^2=0$ and $x+3y^2=1$ is equal to
 (a) $\frac{1}{3}$ (b) $\frac{2}{3}$
 (c) $\frac{4}{3}$ (d) $\frac{5}{3}$
- 64.** The inverse of the statement $(p \wedge \sim q) \rightarrow r$ is
 (a) $\sim(p \vee \sim q) \rightarrow \sim r$
 (b) $(\sim p \wedge q) \rightarrow \sim r$
 (c) $(\sim p \vee q) \rightarrow \sim r$
 (d) None of these
- 65.** The coefficient of the term independent of x in the expansion of $(1+x+2x^3)\left(\frac{3}{2}x^2 - \frac{1}{3x}\right)^9$ is
 (a) $\frac{1}{3}$ (b) $\frac{19}{54}$
 (c) $\frac{17}{54}$ (d) $\frac{1}{4}$
- 66.** The solution to the differential equation $\frac{dy}{dx} = \frac{yf'(x)-y^2}{f(x)}$ where $f(x)$ is a given function is
 (a) $f(x)=y(x+c)$
 (b) $f(x)=cxy$
 (c) $f(x)=c(x+y)$
 (d) $yf(x)=cx$
- 67.** Two fixed points are $A(a, 0)$ and $B(-a, 0)$. If $\angle A - \angle B = \theta$, then the locus of point C of triangle ABC will be
 (a) $x^2 + y^2 + 2xy \tan \theta = a^2$
 (b) $x^2 - y^2 + 2xy \tan \theta = a^2$
 (c) $x^2 + y^2 + 2xy \cot \theta = a^2$
 (d) $x^2 - y^2 + 2xy \cot \theta = a^2$
- 68.** The equation of the planes passing through the line of intersection of the planes $3x-y-4z=0$ and $x+3y+6=0$ whose distance from the origin is 1, are
 (a) $x-2y-2z-3=0$,
 $2x+y-2z+3=0$
 (b) $x-2y+2z-3=0$,
 $2x+y+2z+3=0$
 (c) $x+2y-2z-3=0$,
 $2x-y-2z+3=0$
 (d) None of these
- 69.** In a triangle the length of the two larger sides are 10 and 9, respectively. If the angles are in A.P., then the length of the third side can be :
 (a) $\sqrt{91}$
 (b) $3\sqrt{3}$
 (c) 5
 (d) None of these

70. If $\vec{a} = (1, -1, 2)$, $\vec{b} = (-2, 3, 5)$,
 $\vec{c} = (2, -2, 4)$ and \hat{i} is the unit vector in the x -direction, then
 $(\vec{a} - 2\vec{b} + 3\vec{c})\hat{i} =$

(a) 11	(b) 15
(c) 18	(d) 36

PART-II (Numerical Answer Questions)

71. Find the greatest angle of a triangle whose sides are $a, b, \sqrt{a^2 + b^2 + ab}$.

72. How many 3×3 matrices M with entries from $\{0, 1, 2\}$ are there, for

which the sum of the diagonal entries of $M^T M$ is 5

73. For all complex numbers z_1 , z_2 satisfying $|z_1|=12$ and $|z_2-3-4i|=5$, the minimum value of $|z_1-z_2|$ is

74. The number of positive integral solution of the equation $x_1 x_2 x_3 x_4 x_5 = 1050$ is

75. Two numbers are selected at random from 1, 2, 3..... 100 and are multiplied, then the probability correct to two places of decimals that the product thus obtained is divisible by 3, is

RESPONSE SHEET

PHYSICS

1. a b c d
2. a b c d
3. a b c d
4. a b c d
5. a b c d
6. a b c d
7. a b c d
8. a b c d
9. a b c d
10. a b c d
11. a b c d
12. a b c d
13. a b c d
14. a b c d
15. a b c d
16. a b c d
17. a b c d
18. a b c d
19. a b c d
20. a b c d

21.

22.

23.

24.

CHEMISTRY

26. a b c d
27. a b c d
28. a b c d
29. a b c d
30. a b c d
31. a b c d
32. a b c d
33. a b c d
34. a b c d
35. a b c d
36. a b c d
37. a b c d
38. a b c d
39. a b c d
40. a b c d
41. a b c d
42. a b c d
43. a b c d
44. a b c d
45. a b c d

46.

47.

48.

49.

MATHEMATICS

51. a b c d
52. a b c d
53. a b c d
54. a b c d
55. a b c d
56. a b c d
57. a b c d
58. a b c d
59. a b c d
60. a b c d
61. a b c d
62. a b c d
63. a b c d
64. a b c d
65. a b c d
66. a b c d
67. a b c d
68. a b c d
69. a b c d
70. a b c d

71.

72.

73.

74.

MOCK TEST-1

PHYSICS

1. (b) Since the speeds of the stars are negligible when they are at a distance r , hence the initial kinetic energy of the system is zero. Therefore, the initial total energy of the system is

$$E_i = KE + PE = 0 + \left(-\frac{GMM}{r} \right) = -\frac{GM^2}{r}$$

where M represents the mass of each star and r is initial separation between them.

When two stars collide their centres will be at a distance twice the radius of a star i.e. $2R$.

Let v be the speed with which two stars collide. Then total energy of the system at the instant of their collision is given by

$$E_f = 2 \times \left(\frac{1}{2} Mv^2 \right) + \left(-\frac{GMM}{2R} \right) = Mv^2 - \frac{GM^2}{2R}$$


According to law of conservation of mechanical energy,

$$E_f = E_i$$

$$Mv^2 - \frac{GM^2}{2R} = -\frac{GM^2}{r} \text{ or } v^2 = GM \left(\frac{1}{2R} - \frac{1}{r} \right)$$

$$\text{or } v = \sqrt{GM \left(\frac{1}{2R} - \frac{1}{r} \right)}$$

2. (b)




Work done by normal reaction

3. (c) Rise in temperature, $\Delta\theta = \frac{3T}{J S d} \left(\frac{1}{r} - \frac{1}{R} \right)$

$$\therefore \Delta\theta = \frac{3T}{J} \left(\frac{1}{r} - \frac{1}{R} \right) \quad (\text{For water } S = 1 \text{ and } d = 1)$$

4. (c) $|\vec{F}_B| = |\vec{F}_C| = k \cdot \frac{Q^2}{a^2}$




Hence force experienced by the charge at A in the direction normal to BC is zero.

5. (a) The magnitude of magnetic field at P $\left(\frac{R}{2}, y, \frac{R}{2} \right)$ is


$$B = \frac{\mu_0 J r}{2} = \frac{\mu_0 i}{2\pi R^2} \times \frac{R}{\sqrt{2}} = \frac{\mu_0 i}{2\sqrt{2}\pi R}$$

(independent on y-coordinate)




$$\hat{B} = \frac{\hat{i} - \hat{k}}{\sqrt{2}} \text{ (shown by dotted lines)}$$

$$\therefore \vec{B} = B\hat{B} = \frac{\mu_0 i}{4\pi R} (\hat{i} - \hat{k})$$



$$\text{From figure } B_{net} = \sqrt{B_a^2 + B_e^2}$$

$$\begin{aligned} &= \sqrt{\left(\frac{\mu_0}{4\pi} \cdot \frac{2M}{d^3}\right)^2 + \left(\frac{\mu_0}{4\pi} \cdot \frac{M}{d^3}\right)^2} \\ &= \sqrt{5} \cdot \frac{\mu_0}{4\pi} \cdot \frac{M}{d^3} = \sqrt{5} \times 10^{-7} \times \frac{10}{(0.1)^3} = \sqrt{5} \times 10^{-3} \text{ tesla} \end{aligned}$$



Velocity of A relative to B is given by

$$v_{A/B} = v_A - v_B = v - v_1 \quad \dots (1)$$

By taking x-components of equation (1), we get

$$0 = v \sin \theta - v_1 \Rightarrow \sin \theta = \frac{v_1}{v} \quad \dots (2)$$

Time taken by boy at A to catch the boy at B is given by

$$\begin{aligned} t &= \frac{\text{Relative displacement along Y - axis}}{\text{Relative velocity along Y - axis}} \\ &= \frac{a}{v \cos \theta} = \frac{a}{v \cdot \sqrt{1 - \sin^2 \theta}} = \frac{a}{v \cdot \sqrt{1 - \left(\frac{v_1}{v}\right)^2}} \quad [\text{From equation (1)}] \\ &= \frac{a}{v \cdot \sqrt{\frac{v^2 - v_1^2}{v^2}}} = \frac{a}{\sqrt{v^2 - v_1^2}} = \sqrt{\frac{a^2}{v^2 - v_1^2}} \end{aligned}$$

8. (c)

9. (b) Joule is a unit of energy.

SI	New system
$n_1 = 5$	$n_2 = ?$
$M_1 = 1 \text{ kg}$	$M^2 = \alpha \text{ kg}$
$L_1 = 1 \text{ m}$	$L^2 = \beta \text{ m}$
$T_1 = 1 \text{ s}$	$T^2 = \gamma \text{ s}$

Dimensional formula of energy is comparing with, $[M^a L^b T^c]$, we get
 $a = 1, b = 2, c = -2$

$$\begin{aligned} \text{As } n_2 &= n_1 \left(\frac{M^1}{M^2} \right)^a \left(\frac{L^1}{L^2} \right)^b \left(\frac{T_1}{T_2} \right)^c \\ &= 5 \left(\frac{1 \text{ kg}}{\alpha \text{ kg}} \right)^1 \left(\frac{1 \text{ m}}{\beta \text{ m}} \right)^2 \left(\frac{1 \text{ s}}{\gamma \text{ s}} \right)^{-2} = \frac{5\gamma^2}{\alpha\beta^2} = 5\alpha^{-1}\beta^{-2}\gamma^2 \end{aligned}$$

10. (d)

$$11. (a) I_m = \frac{V_m}{R_f + R_L} = \frac{25}{(10 + 1000)} = 24.75 \text{ mA}$$

$$I_{dc} = \frac{I_m}{\pi} = \frac{24.75}{3.14} = 7.87 \text{ mA}$$

$$I_{rms} = \frac{I_m}{2} = \frac{24.75}{2} = 12.37 \text{ mA}$$

$$P_{dc} = I_{dc}^2 \times R_L = (7.87 \times 10^{-3})^2 \times 10^3 = 61.9 \text{ mW}$$

Rectifier efficiency

$$\eta = \frac{P_{dc}}{P_{ac}} \times 100 = \frac{61.9}{154.54} \times 100 = 40.05\%$$

12. (c) $K_{max} = E - W_0$

$$\therefore T_A = 4.25 - (W_0)_A$$

$$T_B = (T_A - 1.5) = 4.70 - (W_0)_B$$

Equation (i) and (ii) gives $(W_0)_B - (W_0)_A = 1.95 \text{ eV}$

$$\text{De Broglie wave length } \lambda = \frac{h}{\sqrt{2mK}} \Rightarrow \lambda \propto \frac{1}{\sqrt{K}}$$

$$\Rightarrow \frac{\lambda_B}{\lambda_A} = \sqrt{\frac{K_A}{K_B}} \Rightarrow 2 = \sqrt{\frac{T_A}{T_B - 1.5}} \Rightarrow T_A = 2eV$$

From equation (i) and (ii)

$$W_A = 2.25 \text{ eV} \text{ and } W_B = 4.20 \text{ eV.}$$

13. (c) $PV = \mu RT = \frac{m}{M} RT,$

where m = mass of the gas

and $\frac{m}{M} = \mu$ = number of moles.

$$\frac{PV}{T} = \mu R = \text{a constant for all values of } P.$$

That is why, ideally it is a straight line.

$$\therefore \frac{PV}{T} = \frac{1g}{32g \text{ mol}^{-1}} \times 8.31 \text{ J mol}^{-1} \text{ K}^{-1} = 0.259 \text{ JK}^{-1}$$

Also, $T_1 > T_2$


14. (c) $f_{\text{apparent}} = \left(\frac{u + u/5}{u} \right) f = \frac{6}{5} f = 1.2f$

Wavelength remains constant (unchanged) in this case.

15. (d) Consider a shell of thickness (dr) and of radii (r) and the temperature of inner and outer surfaces of this shell be T , $(T - dT)$

$$\frac{dQ}{dt} = \text{rate of flow of heat through it}$$

$$= -4\pi K r^2 \frac{dT}{dr} \quad (\because A = 4\pi r^2)$$




To measure the radial rate of heat flow, integration technique is used, since the area of the surface through which heat will flow is not constant.

$$\text{Then, } \left(\frac{dQ}{dt} \right) = \int_{r_1}^{r_2} \frac{1}{r^2} dr = -4\pi K \int_{T_1}^{T_2} dT$$


$$\frac{dQ}{dt} \left[\frac{1}{r_1} - \frac{1}{r_2} \right] = -4\pi K [T_2 - T_1]$$

$$\text{or } \frac{dQ}{dt} = \frac{-4\pi K r_1 r_2 (T_2 - T_1)}{(r_2 - r_1)} \quad \therefore \quad \frac{dQ}{dt} \propto \frac{r_1 r_2}{(r_2 - r_1)}$$

- 16. (b)** $W_{\text{ext}} =$ negative of area with volume-axis
 $W(\text{adiabatic}) > W(\text{isothermal})$



- 17. (a)**



Given $AQ = AR$ and $\angle A = 60^\circ$

$$\therefore \angle AQR = \angle ARQ = 60^\circ$$

$$\therefore r_1 = r_2 = 30^\circ$$

Applying Snell's law on face AB.

$$1. \sin i_1 = \mu \sin r_1$$

$$\Rightarrow \sin i_1 = \sqrt{3} \sin 30^\circ = \sqrt{3} \times \frac{1}{2} = \frac{\sqrt{3}}{2}$$

$$\therefore i_1 = 60^\circ$$

Similarly, $i_2 = 60^\circ$

In a prism, deviation


$$\delta = i_1 + i_2 - A = 60^\circ + 60^\circ - 60^\circ = 60^\circ$$

18. (a) Both magnetic and electric fields have zero average value in a plane e.m. wave.

19. (a) At $t = 0$, current will flow only in 12Ω resistance

$$\therefore I_{\min} = \frac{5}{12}$$

At $t \rightarrow \infty$ both L_1 and L_2 behave as conductign wires



$$\therefore R_{\text{eff}} = \frac{3}{2}, I_{\max} = \frac{10}{3}$$

$$\frac{I_{\max}}{I_{\min}} = 8$$


20. (d) For the first minima,

$$\theta = \frac{\eta\lambda}{a} \Rightarrow \sin 30^\circ = \frac{\lambda}{a} = \frac{1}{2}$$

First secondary maxima will be at

21. (11.75) Let E_1 and E_2 be potential drops across R and X so

$$\frac{E_2}{E_1} = \frac{IX}{IR} = \frac{X}{R} \quad \text{or} \quad X = \frac{E_2}{E_1} R$$



$$\text{But } \frac{E_2}{E_1} = \frac{\ell_2}{\ell_1}$$

$$\text{so } X = \frac{\ell_2}{\ell_1} R = \frac{68.5}{58.3} \times 10 = 11.75 \Omega$$

22. (6.66) Given : Speed $V = 54 \text{ kmh}^{-1} = 15 \text{ ms}^{-1}$

Moment of inertia, $I = 3 \text{ kgm}^2$

Time $t = 15 \text{ s}$

$$\omega_i = \frac{V}{r} = \frac{15}{0.45} = \frac{100}{3} \quad \omega_f = 0$$

$$\omega_f = \omega_i + \alpha t$$

$$0 = \frac{100}{3} + (-\alpha)(15) \quad \Rightarrow \quad \alpha = \frac{100}{45}$$

Average torque transmitted by brakes to the wheel

$$\tau = (I)(\alpha) = 3 \times \frac{100}{45} = 6.66 \text{ kgm}^2 \text{s}^{-2}$$

23. (0.05) Given : $A = 4 \text{ m}^2$, $e = 0.32 \text{ V}$, $dt = 0.5 \text{ sec}$.

B_1 is the initial magnetic induction and when it is reduced to 20%

$$B_2 = 0.2 B_1$$

$$e = \frac{d\phi}{dt} = \frac{A(B_1 - B_2)}{\Delta t} \quad \text{or} \quad 0.32 = \frac{4(B_1 - 0.2 B_1)}{0.5}$$


24. (0.94) Time period of a physical pendulum is

$$T = 2\pi \sqrt{\frac{I}{mgh}}$$

where I is the moment of inertia
of the pendulum about an axis

through the pivot, m is the mass of the pendulum and h is the distance
from the pivot to the centre of mass.

In this case, a solid disc of R oscillates as a physical pendulum about an
axis perpendicular to the plane of the disc at a distance r from its centre.



$$\begin{aligned} \therefore I &= \frac{mR^2}{2} + mr^2 = \frac{mR^2}{2} + m\left(\frac{R}{4}\right)^2 = \frac{mR^2}{2} + \frac{mR^2}{16} \\ &= \frac{9mR^2}{16} \quad \left(\because r = \frac{R}{4} \right) \end{aligned}$$

$$\text{Here, } R = 10 \text{ cm} = 0.1 \text{ m}, h = \frac{R}{4}$$

$$\therefore T = 2\pi \sqrt{\frac{\frac{9mR^2}{16}}{mgR}} = 2\pi \sqrt{\frac{9R}{4g}}$$

$$= 2\pi \sqrt{\frac{9 \times 0.1}{4 \times 10}} = 2\pi \times \frac{3}{2} \times \frac{1}{10} = 0.94 \text{ s}$$

25. (488.9) $\frac{1}{\lambda_1} = R \left(\frac{1}{2^2} - \frac{1}{3^2} \right) = \frac{5R}{36}$

$$\frac{1}{\lambda_2} = R \left(\frac{1}{2^2} - \frac{1}{4^2} \right) = \frac{3R}{16}$$

$$\therefore \frac{\lambda_2}{\lambda_1} = \frac{80}{108}$$

$$\lambda_2 = \frac{80}{108} \lambda_1 = \frac{80}{108} \times 660 = 488.9 \text{ nm.}$$

27. (b) CuF₂ is both paramagnetic and coloured.

28. (c) *s*-character \propto bond angle

For 25% *s* character (as in *sp*³ hybrid orbital), bond angle is 109.5°, for 33.3% *s* character (as in *sp*² hybrid orbital), bond angle is 120° and for 50% *s* character (as in *sp* hybrid orbital), bond angle is 180°.

Similarly, when the bond angle decreases below 109.5°, the *s*-character will decrease accordingly.

$$\text{Decrease in angle} = 120^\circ - 109.5^\circ = 10.5^\circ$$


$$\text{Decrease in } s\text{-character} = 33.3 - 25 = 8.3$$

$$\text{Actual decrease in bond angle} = 109.5^\circ - 105^\circ = 4.5^\circ$$

Expected decrease in *s*-character

$$= \frac{8.3}{10.5} \times 4.5 = 3.56\%$$


Thus, the *s*-character should decrease by about 3.56% i.e., *s*-character = 25 - 3.56 = 21.44%




30. (a) Be – 1s²2s²; B – 1s²2s²2p¹; C – 1s²2s²2p²; N – 1s²2s²2p³; O – 1s²2s²2p⁴. IP increases along the period. But IP of Be > B. Further IP of O < N because atoms with fully or partly filled orbitals are most stable and hence have high ionisation energy.

31. (a) From data 1 and 3, it is clear that keeping (B) const, [A] is doubled, rate remains unaffected. Hence rate is independent of [A]. From 1 and 4, keeping [A] constant, [B] is doubled, rate become 8 times. Hence rate $\propto [B]^3$.

32. (b) (i) HCl is a strong acid. Hence its pH is lowest among the others.
(ii) NaCl is a salt of strong acid and strong base so it is not hydrolysed and hence its pH is 7.



∴ The solution is acidic and pH is less than that of 0.1 M HCl.



∴ The solution is basic and pH is more than that of 0.1 M HCl.

∴ Correct order for increase in pH is



34. (b) All proteins are not found in L-form but they may be present in form of D or L



36. (c) For metal, as temperature increases, resistance increases and hence conductivity decreases.

37. (d) In a unit cell, W atoms at the corner = $\frac{1}{8} \times 8 = 1$


$$\text{O-atoms at the centre of edges} = \frac{1}{4} \times 12 = 3$$

Na-atoms at the centre of the cube = 1

W : O : Na = 1 : 3 : 1


Hence, formula = NaWO_3

38. (d) With Br_2 water, phenol gives 2, 4, 6-tribromophenol.



39. (c) V_2O_5 is used as catalyst in contact process of manufacturing H_2SO_4

40. (c) The dipole moment of symmetrical molecule is zero.



Triangular planar (symmetrical molecule)


41. (a) $\text{NaCl(s)} \rightarrow \text{NaCl(l)}$

Given that : $\Delta H = 30.5 \text{ kJ mol}^{-1}$

$$\Delta S = 28.8 \text{ JK}^{-1} \text{mol}^{-1} = 28.8 \times 10^{-3} \text{ kJ K}^{-1} \text{mol}^{-1}$$

$$\text{By using } \Delta S = \frac{\Delta H}{T} = \frac{30.5}{28.8 \times 10^{-3}} = 1059 \text{ K.}$$

42. (a)



44. (b) Radioactive decay follows first order kinetics. therefore,



$$\text{Decay constant } (\lambda) = \frac{0.693}{t_{1/2}} = \frac{0.693}{5730}$$

$$\text{Given, } R_0 = 100 \quad \therefore \quad R = 80$$

$$\text{and } t = \frac{2.303}{\lambda} \log \frac{[R]_0}{[R]} = \frac{2.303}{\left(\frac{0.693}{5730}\right)} \log \frac{100}{80}$$

$$= \frac{2.303 \times 5730}{0.693} \times 0.0969 = 1845 \text{ years}$$

45. (d) Hybridisation



Hence $[\text{Ni}(\text{NH}_3)_6]^{2+}$ is outer orbital complex.

46. (409.5) According to combined gas equation,

$$\frac{PV}{T} = \frac{P_1V_1}{T_1}$$





$$P = 1 \text{ atm}, P_1 = \frac{3}{4} \text{ atm (on reducing by 25%)}$$

$$V = v, V_1 = 2v, T = 273K, T_1 = ?$$

$$\frac{1 \times v}{273} = \frac{3 \times 2v}{4 \times T_1}$$

$$T_1 = \frac{3 \times 2 \times 273}{4} = 409.5 \text{ K}$$

47. (4) $\text{C}_7\text{H}_7\text{Cl}$ has 4 isomers



48. (8.4) Normality of $\text{H}_2\text{O}_2 = \frac{\text{vol. strength}}{5.6}$

Volume of (1N) H_2O_2 solution = 5.6 volumes.

$$\therefore \text{Volume strength of } 1.5 \text{ N } \text{H}_2\text{O}_2 \\ = 1.5 \times 5.6 = 8.4 \text{ volumes.}$$

49. (26) ABAB... is hexagonal close packing (*hcp*) in which space occupied = 74 % and, empty space = 26%.



Eq. of M_2O_x = eq. of Metal

$$\frac{\text{Wt. of } \text{M}_2\text{O}_x}{\text{Eq. wt. of } \text{M}_2\text{O}_x} = \frac{\text{Wt. of Metal}}{\text{Eq. wt. of Metal}}$$

$$\frac{4}{2 \times 56 + x \times 16} = \frac{2.8}{56}$$

$$\frac{2x}{x} = \frac{56}{2.8}$$

On solving we get,

$$\Rightarrow \frac{4}{56 + 8x} = \frac{2.8}{56} \Rightarrow \frac{1}{14 + 2x} = \frac{1}{20} \Rightarrow 2x = 6 \Rightarrow x = 3$$

Hence, the oxide is M_2O_3 .

MATHEMATICS

51. (d) $T_4 = {}^nC_3 x^{n-3} \left(\frac{\alpha}{2x}\right)^3 \Rightarrow {}^nC_3 x^{n-6} \left(\frac{\alpha}{2}\right)^3 = 20$

$$\text{If } n = 6, \text{ then } {}^6C_3 \left(\frac{\alpha}{2}\right)^3 = 20 \Rightarrow \alpha = 2$$

52. (b) Given equation is $x^2 + px + q = 0$

Sum of roots = $\tan 30^\circ + \tan 15^\circ = -p$

Product of roots = $\tan 30^\circ \cdot \tan 15^\circ = q$

$$\tan 45^\circ = \frac{\tan 30^\circ + \tan 15^\circ}{1 - \tan 30^\circ \cdot \tan 15^\circ} = \frac{-p}{1-q} = 1$$

$$\Rightarrow -p = 1 - q \Rightarrow q - p = 1$$

$$\therefore 2 + q - p = 2$$

or $(a+c)(a+b), (b+c)(a+b), (c+a)(b+c)$.. are also in

A.P. $\Rightarrow \frac{1}{b+c}, \frac{1}{c+a}, \frac{1}{a+b}$ are in A.P.

[dividing by $(a+b)(b+c)(c+a)$]

- 54. (d)** Let $M(h, k)$ be the mid-point

of chord AB where

$$\angle ACB = \frac{2\pi}{3}$$

$$\therefore \angle ACM = \frac{\pi}{3}$$

$$\text{Also } CM = 3 \cos \frac{\pi}{3} = \frac{3}{2}$$

$$\Rightarrow \sqrt{h^2 + k^2} = \frac{3}{2} \Rightarrow h^2 + k^2 = \frac{9}{4}$$

$$\therefore \text{Locus of } (h, k) \text{ is } x^2 + y^2 = \frac{9}{4}$$

$$\begin{aligned} \text{55. (c)} \quad y &= \tan^{-1} \left(\frac{\log_e(e/x^2)}{\log_e(ex^2)} \right) + \tan^{-1} \left(\frac{3+2\log_e x}{1-6\log_e x} \right) \\ &= \tan^{-1} \left(\frac{1-2\log_e x}{1+2\log_e x} \right) + \tan^{-1} \left(\frac{3+2\log_e x}{1-3.2\log_e x} \right) \\ &= \tan^{-1}(a) - \tan^{-1}(2\log_e x) \\ &\quad + \tan^{-1}(c) + \tan^{-1}(2\log_e x) \\ &= \tan^{-1}(a) + \tan^{-1}(c) \end{aligned}$$

$$\therefore \frac{dy}{dx} = 0$$

- 56. (d)** $\{x^2\} - 2\{x\} \geq 0$

$$\Rightarrow \{x\}(\{x\} - 2) \geq 0$$


$$\Rightarrow \{x\} \leq 0 \text{ or } \{x\} \geq 2$$


Second case is not possible.

Hence $\{x\} = 0$, as $\{x\} \leq [0, 1)$

Hence range of $f(x)$ contains only one element 0.

- 57. (d)** Here $p = 7$ and $\alpha = 30^\circ$





$$\text{or } \sqrt{3}x + y = 14$$

58. (c) $I = \int \frac{dx}{\cos x + \sqrt{3} \sin x}$

$$I = \int \frac{dx}{2 \left[\frac{1}{2} \cos x + \frac{\sqrt{3}}{2} \sin x \right]}$$

$$= \frac{1}{2} \int \frac{dx}{\left[\sin \frac{\pi}{6} \cos x + \cos \frac{\pi}{6} \sin x \right]} = \frac{1}{2} \int \frac{dx}{\sin \left(x + \frac{\pi}{6} \right)}$$

$$\Rightarrow I = \frac{1}{2} \int \operatorname{cosec} \left(x + \frac{\pi}{6} \right) dx$$

$$\therefore \int \operatorname{cosec} x dx = \log |(\tan x / 2)| + C$$

$$\therefore I = \frac{1}{2} \log \tan \left(\frac{x}{2} + \frac{\pi}{12} \right) + C$$

59. (c) Given $\frac{\tan 3\theta - 1}{\tan 3\theta + 1} = \sqrt{3}$

$$\Rightarrow \sqrt{3} (\tan 3\theta + 1) = \tan 3\theta - 1$$

$$\Rightarrow \sqrt{3} \tan 3\theta + \sqrt{3} = \tan 3\theta - 1$$

$$\Rightarrow \tan 3\theta (\sqrt{3} - 1) = -(1 + \sqrt{3})$$

$$\Rightarrow \tan 3\theta = \frac{-(\sqrt{3} + 1)}{(\sqrt{3} - 1)} = \frac{-(1 + \sqrt{3})}{-(1 - \sqrt{3})} = \frac{1 + \sqrt{3}}{1 - \sqrt{3}}$$

$$\Rightarrow \tan 3\theta = \tan 105^\circ = \tan \frac{7\pi}{12}$$

[Note: $\tan \theta = \tan \alpha \Rightarrow \theta = n\pi + \alpha$]

$$\therefore 3\theta = n\pi + \frac{7\pi}{12} \Rightarrow \theta = \frac{n\pi}{3} + \frac{7\pi}{36}$$

60. (c) Equation of normal in slope form on $y^2 = 4Ax$ is

$$y = mx - 2Am - Am^3 \quad \dots(i)$$

$$= mx - 2\left(\frac{1}{4}\right)m - \left(\frac{1}{4}\right)m^3 \quad \begin{bmatrix} \because y^2 = x \\ \therefore A = \frac{1}{4} \end{bmatrix}$$

$$\Rightarrow 4mx - 4y - m^3 - 2m = 0$$

$\because (a, 0)$ lies on the normal. Then, $4m \times a - 4 \times 0 - m^3 - 2m = 0$

$$\Rightarrow m(m^2 + 2 - 4a) = 0$$

$$\Rightarrow m = 0 \text{ or } m^2 + 2 - 4a = 0$$

If $m = 0$, then from (i),

$y = 0$ i.e., x -axis is one normal.

$$\text{If } m^2 + 2 - 4a = 0 \Rightarrow m^2 = 4a - 2 \quad [\because m^2 > 0]$$


$$\Rightarrow 4a - 2 > 0 \Rightarrow a > \frac{1}{2}.$$

61. (d) By the diagram only 2 rectangles are formed $ADEH, GFCH$.

\therefore number of favourable cases = 2

Total number of cases = 8C_4

$$\therefore \text{required probability} = \frac{2}{{}^8C_4} = \frac{1}{35}$$



62. (a) $f(x) = x^3 - 3x^2 - 24x + 5$

For increasing, $f'(x) > 0, 3x^2 - 6x - 24 > 0$

$$\Rightarrow x^2 - 2x - 8 > 0 \Rightarrow x^2 - 4x + 2x - 8 > 0$$

$$\Rightarrow (x+2)(x-4) > 0.$$

Now, by the sign scheme for $3x^2 - 6x - 24$,



$$\Rightarrow x \in (-\infty, -2) \cup (4, \infty)$$

63. (c) Given that $y = y(x)$ and $x \cos y + y \cos x = \pi$... (i)

For $x = 0$ in (i) we get $y = \pi$

Differentiating (i) with respect to x , we get,

$$-x \sin y \cdot y' + \cos y + y' \cos x - y \sin x = 0$$

$$\Rightarrow y' = \frac{y \sin x - \cos y}{\cos x - x \sin y} \quad \dots \text{(ii)}$$


$$\Rightarrow y'(0) = 1 \text{ (Using } y(0) = \pi)$$

Differentiating (ii) with respect to x , we get,

$$y'' = \frac{(y' \sin x + y \cos x + \sin y \cdot y')(\cos x - x \sin y) - (-\sin x - \sin y - x \cos y \cdot y')(\cos x - x \sin y)}{(\cos x - x \sin y)^2}$$

$$\Rightarrow y''(0) = \frac{\pi(1) - 0}{1} = \pi.$$

64. (b) DP is a clock tower standing at the middle point D of BC .




$$\therefore \cot \beta = \sqrt{\operatorname{cosec}^2 \beta - 1} = \sqrt{(5.76)} = 2.4$$

In the triangles PAD and PBD ,

$$AD = h \cot \alpha = 3.2 \text{ } h \text{ and } BD = h \cot \beta = 2.4 \text{ } h$$

$$\text{In the right angled } \Delta ABD, AB^2 = AD^2 + BD^2 \\ \Rightarrow 100^2 = [(3.2)^2 + (2.4)^2] h^2 \Rightarrow h^2 = 16 \text{ } h^2 \Rightarrow h = 25 \text{ m.}$$

65. (b)



$$\overrightarrow{AD} = \frac{(-3+5)\hat{i} + (0-2)\hat{j} + (4+4)\hat{k}}{2}$$

$$= \frac{2\hat{i} - 2\hat{j} + 8\hat{k}}{2} = \hat{i} - \hat{j} + 4\hat{k}$$

\therefore length of median

$$= |\overrightarrow{AD}| = \sqrt{(1)^2 + (-1)^2 + (4)^2} = \sqrt{18}$$

$$\begin{aligned} \text{66. (a)} \quad & \sim [p \vee (\sim p \vee q)] \equiv \sim p \wedge \sim (\sim p \vee q) \\ & \equiv \sim p \wedge (\sim (\sim p) \wedge \sim q) \\ & \equiv \sim p \wedge (p \wedge \sim q). \end{aligned}$$

$$\text{67. (a)} \quad f(x) = x^p \sin \frac{1}{x}, x \neq 0 \text{ and } f(x) = 0, x = 0$$

Since at $x = 0$, $f(x)$ is a continuous function

$$\therefore \lim_{x \rightarrow 0} f(x) = f(0) = 0$$

$$\Rightarrow \lim_{x \rightarrow 0} x^p \sin \frac{1}{x} = 0 \Rightarrow p > 0$$

$f(x)$ is differentiable at $x = 0$, if $\lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0}$ exists


$$x^p \sin \frac{1}{x} - 0$$

$$\Rightarrow \lim_{x \rightarrow 0} x^{p-1} \sin \frac{1}{x} \text{ exists}$$

$$\Rightarrow p-1 > 0 \text{ or } p > 1$$

\therefore for $0 < p \leq 1$, $f(x)$ is a continuous function at $x = 0$ but not differentiable.

68. (d) Let M be the foot of perpendicular from $(7, 14, 5)$ to the given plane, then PM is normal to the plane. So, its d.r.'s are $2, 4, -1$. Since PM passes through $P(7, 14, 5)$ and has d.r.'s $2, 4, -1$.



$$\text{Therefore, its equation is } \frac{x-7}{2} = \frac{y-14}{4} = \frac{z-5}{-1} = r,$$

$$\Rightarrow x = 2r + 7, \quad y = 4r + 14, \quad z = -r + 5$$

$$\text{Co-ordinates of } M \text{ be } (2r+7, 4r+14, -r+5)$$

$$\text{Since } M \text{ lies on the plane } 2x + 4y - z = 2, \text{ therefore}$$

$$2(2r+7) + 4(4r+14) - (-r+5) = 2 \Rightarrow r = -3$$

$$\text{Co-ordinates of foot of perpendicular are } M(-1, 2, 8).$$

$$PM = \text{Length of perpendicular from } P$$

$$= \sqrt{(1-7)^2 + (2-14)^2 + (8-5)^2} = 3\sqrt{21}.$$

$$69. (c) \quad \Delta(x) = \begin{vmatrix} e^x & \sin x \\ \cos x & \ln(1+x^2) \end{vmatrix}$$

$$= e^x \ln(1+x^2) - \sin x \cos x$$

$$\text{So, } \lim_{x \rightarrow 0} \frac{\Delta(x)}{x} = \lim_{x \rightarrow 0} \frac{e^x \ln(1+x^2) - \sin x \cos x}{x}$$

70. (a) $\tan^{-1} x + \tan^{-1} \frac{1}{y} = \tan^{-1} 3$

$$\Rightarrow \tan^{-1} \frac{x + \frac{1}{y}}{1 - \frac{x}{y}} = \tan^{-1} 3 \Rightarrow \frac{xy + 1}{y - x} = 3$$

$$\Rightarrow y = \frac{1+3x}{3-x} > 0 \quad [\because x \text{ and } y \text{ are positive}]$$

$$\Rightarrow x - 3 < 0 \Rightarrow x < 3 \text{ or } x = 1, 2$$

$$\therefore y = 2, 7$$

solution set is $(x, y) \in \{(1, 2), (2, 7)\}$

71. (64) A selection of 3 balls so as to include at least one black ball, can be made in the following 3 mutually exclusive ways

$$(i) 1 \text{ black ball and 2 others} = {}^3C_1 \times {}^6C_2 = 3 \times 15 = 45$$

$$(ii) 2 \text{ black balls and one other} = {}^3C_2 \times {}^6C_1 = 3 \times 6 = 18$$

$$(iii) 3 \text{ black balls and no other} = {}^3C_3 \times {}^6C_0 = 1$$

$$\therefore \text{Total number of ways} = 45 + 18 + 1 = 64.$$

72. (19.5)

Class	Frequency	Cumulative Frequency
5–10	5	5
10–15	6	11
15–20	15	26
20–25	10	36
25–30	5	41
30–35	4	45
35–40	2	47
40–45	2	49

$$N = 49$$

$$\text{Here } N = 49. \quad \therefore \frac{N}{2} = \frac{49}{2} = 24.5$$

The cumulative frequency just greater than $N/2$ is 26 and corresponding class is 15–20. Thus 15–20 is the median class such that $\ell = 15, f = 15, F = 11, h = 5$

$$\therefore \text{median} = \ell + \frac{\frac{N}{2} - F}{f} \times h$$

73. (3.75) α, β are roots of the equation $2x^2 + 3x + 5 = 0$

$$\text{Therefore sum of roots } (\alpha + \beta) = -\frac{3}{2}$$

$$\text{And product of roots } (\alpha \cdot \beta) = \frac{5}{2}.$$

$$\begin{aligned}\text{Now, } & \left| \begin{array}{ccc} 0 & \beta & \beta \\ \alpha & 0 & \alpha \\ \beta & \alpha & 0 \end{array} \right| \\ &= 0 | 0 - \alpha^2 | - \beta | 0 - \alpha\beta | + \beta | \alpha^2 - 0 | \\ &= \alpha\beta^2 + \beta\alpha^2 = \alpha\beta(\alpha + \beta) \\ &= \frac{5}{2} \left(\frac{-3}{2} \right) = \frac{-15}{4} = -3.75\end{aligned}$$

$$\textbf{74. (21.5)} \text{ Let } I = \int_{-3}^2 \{ |x+1| + |x+2| + |x-1| \} dx$$

Breaking points are

$$x+1=0 \Rightarrow x=-1$$

$$x+2=0 \Rightarrow x=-2$$

$$x-1=0 \Rightarrow x=1$$

$$\therefore I = \int_{-3}^{-2} f(x) dx + \int_{-2}^{-1} f(x) dx + \int_{-1}^1 f(x) dx + \int_1^2 f(x) dx$$

$$\text{where } f(x) = |x+1| + |x+2| + |x-1|$$

$$\text{Now, } I_1 = \int_{-3}^{-2} [-(x+1) - (x+2) - (x-1)] dx$$

$$= - \left[\frac{x^2}{2} + x + \frac{x^2}{2} + 2x + \frac{x^2}{2} + x \right]_{-3}^{-2} = \frac{7}{2}$$

$$I_2 = \int_{-2}^{-1} [-(x+1) + (x+2) - (x-1)] dx$$

$$= - \left[\frac{x^2}{2} - x + \frac{x^2}{2} + 2x - \frac{x^2}{2} - x \right]_{-2}^{-1} = \frac{7}{2}$$

$$= \frac{-x^2}{2} + 2x \Big|_{-2}^{-1} = \left(\frac{-1}{2} - 2 \right) - (-2 - 4)$$

$$= -\frac{5}{2} + 6 = \frac{7}{2}$$

$$I_3 = \int_{-1}^1 [(x+1) + (x+2) - (x-1)] dx$$

$$= \int_{-1}^1 (x+4) dx = \frac{x^2}{2} + 4x \Big|_{-1}^1 = 8$$

$$I_4 = \int_1^2 [(x+1) + (x+2) + (x-1)] dx$$

$$= \int_1^2 (3x+2) dx = \frac{3x^2}{2} + 2x \Big|_1^2 = \frac{13}{2}$$

$$\therefore I = I_1 + I_2 + I_3 + I_4$$

$$= \frac{7}{2} + \frac{7}{2} + 8 + \frac{13}{2} = \frac{43}{2}$$

$$= \frac{27+16}{2} = \frac{43}{2} = 21.5$$

75. (4.5) Given curves are, $y = 2x - x^2$

.....(i)

and $y = -x$

.....(ii)

Putting the value of y in (i),


$$-x = 2x - x^2$$

$$\Rightarrow x(x-3) = 0 \Rightarrow x = 0, 3$$

\therefore area under the curve

$$= \int_0^3 [(2x - x^2) - (-x)] dx$$

$$= \int_0^3 (3x - x^2) dx = \left[\frac{3x^2}{2} - \frac{x^3}{3} \right]_0^3$$



MOCK TEST-2

PHYSICS

1. (d) Let the scooterist velocity be v . Then

$$1000 + (10 \times 100) = v \times 100$$

$$\Rightarrow 100v = 2000 \Rightarrow v = \frac{2000}{100} = 20 \text{ m/s}$$

2. (b) We have, $F = kx$

where, F , x and k are force, length and constant respectively.

$$\therefore 5 = kx \quad \dots\dots(1)$$

$$\text{and } 7 = ky \quad \dots\dots(2)$$

Multiplying eq. (2) by 2

$$14 = 2ky \quad \dots\dots(3)$$

Subtracting eq. (1) from (3),

$$14 - 5 = 2ky - kx \text{ or } 9 = k(2y - x)$$

Hence, required length = $2y - x$

3. (b) In the given equation $[\rho] = [b][x]$;

$\therefore [b] = [\rho]/[x]$. But ρ is mass per unit length and x is distance, therefore $[b] = ML^{-1}/L = ML^{-2}T^0$

4. (d) Point A is at rest w.r.t. motion, hence, v at A = 0. At point B there are two horizontal velocities. Hence, $v_B = 2v$.

5. (d) $mg = 2TL \Rightarrow \pi r^2 L dg = 2TL \Rightarrow \pi r^2 dg = 2T$.

This relation is independent of L.

6. (d) $\omega_{\text{rod}} = \omega_{\text{point}} = \left(\frac{v_{\text{rel}}}{r} \right)$,

v_{rel} represents the velocity of one point w.r.t. other.

$$= \frac{3v - v}{r} \text{ and 'r' being the distance between them.}$$

$$= \frac{2v}{r}$$

7. (d) Both are diatomic gases and $C_p - C_V = R$ for all gases.

8. (b) More the initial temperature more is the rate of cooling.

Hence, $T_3 > T_2 > T_1$

or

The rate of cooling decreases with decrease in temperature difference between body and surrounding.



9. (c) $\frac{1}{C_\infty} = \frac{1}{C_\infty + C} + \frac{2}{C} = \frac{3C + 2C_\alpha}{C_\alpha(C_\alpha + C)}$

10. (b) R increases with increasing temp:

$$V=IR$$

Slope of graph $= \frac{I}{V} = \frac{1}{R}$; Slope of T_1 is more i.e. $\frac{1}{R_1}$ is more, hence R_1 is less. This concludes that T_1 will be less than T_2 as R_1 is less than R_2 .

11. (c) $R_{eq} = \frac{R_1 R_2}{R_1 + R_2}$ (Parallel) $= \frac{4 \times 2}{4 + 2} = \frac{8}{6} = \frac{4}{3} \Omega$



12. (d) $i_R = \frac{V_0}{R} = \frac{100}{20} = 5$, $i_L = \frac{V_0}{X_L} = \frac{100}{10} = 10$ and $i_C = \frac{V_0}{X_C} = \frac{100}{20} = 5$

Current, $i = \sqrt{i_R^2 + (i_C - i_L)^2} = \sqrt{5^2 + 5^2} = 5\sqrt{2}$ amp.

13. (a) $f_{max} = \mu mg$, $a_{max} = \mu g$.

If A is the amplitude $a_{max} = A\omega^2 = 4\pi^2 AV^2 = \mu g$.

Therefore, $A = \frac{\mu g}{4\pi^2 V^2}$.

14. (c) Total time taken to travel distance d is :

$$\frac{d}{2n_1} + \frac{d}{2n_2} = d \left(\frac{n_1 + n_2}{2n_1 n_2} \right) = \frac{d}{n_{eff}}; \quad n_2 = 3n_1 \Rightarrow n_{eff} = \frac{3}{2} n_1$$

However, if A_{12} and A_{34} have equal magnitude because of random phase of A_{12} and A_{34} , no fringes will be seen.

16. (a) $mvr = \frac{nh}{2\pi}, \lambda = \frac{h}{mv};$

Using the two concept we get, $mvr = \frac{nh}{2\pi}$ (where $n = 1$)

$$2\pi r = \frac{1 \times h}{mv} \quad \dots(1)$$

$$\lambda = \frac{h}{mv} \quad \dots(2)$$

$$\text{Divide (2) by (1), } \frac{2\pi r}{\lambda} = \frac{h \times mv}{mv \times h} = \frac{1}{1} = 1 : 1$$

17. (d) 1. $\lambda = \frac{0.693}{t^{1/2}}$ 2. $R = \lambda N_t$

Radioactivity at T_1 is $R_1 = \lambda N_1$,

Radioactivity at T_2 is $R_2 = \lambda N_2$

\therefore Number of atoms decayed in time

$$(T_1 - T_2) = (N_1 - N_2) \text{ or } \frac{R_1 - R_2}{\lambda} = \frac{(R_1 - R_2)T}{0.693}$$

i.e., $\alpha (R_1 - R_2)T$

18. (d) In the graph given, slope of curve 2 is greater than the slope of curve 1.


$$\left(\frac{\gamma P}{V}\right)_2 > \left(\frac{\gamma P}{V}\right)_1 \Rightarrow \gamma_2 > \gamma_1$$

$$\gamma_{\text{He}} > \gamma_{\text{O}_2}$$

Since, $\gamma_{\text{monoatomic}} > \gamma_{\text{diatomic}}$

Hence, curve 2 corresponds to helium and curve 1 corresponds to oxygen.

19. (b) $V_A = \frac{1}{4\pi \epsilon_0} \left[\frac{Q_1}{R} + \frac{Q_2}{\sqrt{2}R} \right]; \quad V_B = \frac{1}{4\pi \epsilon_0} \left[\frac{Q_2}{R} + \frac{Q_1}{\sqrt{2}R} \right]$



$$V_A - V_B = \frac{1}{4\pi \epsilon_0 R} \left[Q_1 + \frac{Q_2}{\sqrt{2}} - Q_2 - \frac{Q_1}{\sqrt{2}} \right]$$

$$\text{Work done} = Q \times V = q \times (V_A - V_B)$$

$$= \frac{q}{4\pi \epsilon_0 R} \left[Q_1 + \frac{Q_2}{\sqrt{2}} - Q_2 - \frac{Q_1}{\sqrt{2}} \right]$$


$$= \frac{q}{4\pi \epsilon_0 R} \times \frac{1}{\sqrt{2}} \left[\sqrt{2}Q_1 + Q_2 - \sqrt{2}Q_2 - Q_1 \right]$$

$$= \frac{q(Q_1 - Q_2)(\sqrt{2} - 1)}{(\sqrt{2} 4\pi \epsilon_0 R)}$$

- 20. (b)** When there is no change in liquid level in vessel then $\gamma'_{\text{real}} = \gamma'_{\text{vessel}}$
 Change in volume in liquid relative to vessel

$$\Delta V_{\text{app}} = V\gamma'_{\text{app}} \quad \Delta\theta = V(\gamma'_{\text{real}} - \gamma'_{\text{vessel}})$$

- 21. (10)** Here, $m_A = 0.5\text{kg}$; $m_B = 1\text{kg}$



Force on block A

$$T = \mu m_A g \quad \dots\dots(1)$$

Force acting on block B

$$F = T + \mu m_A g + \mu(m_A + m_B)g \quad \dots\dots(2)$$

From (1) & (2),

$$F = \mu m_A g + \mu m_A g + \mu m_A g + \mu m_B g$$

$$\begin{aligned} F &= 3\mu m_A g + \mu m_B g = \mu g(3m_A + m_B) \\ &= 0.4 \times 10 \times (3 \times 0.5 + 1) = 10\text{N} \end{aligned}$$

- 22. (10.6)**

Work done in going from a distance r_1 to a distance r_2 away from centre of the earth, by a body of mass m , is,

$$W = GMm(1/r_1 - 1/r_2),$$

For our case we should have

$$1/2 mv^2 = GMm [(1/R_e) - (1/10R_e)]$$

23. (300)

Here the number of molecules is same. Hence,

$$T_{\text{final}} = \frac{T_1 + T_2}{2} = \frac{200 + 400}{2} = 300 \text{ K}$$

24. (83.3) Power of source = $EI = 240 \times 0.7 = 166$

$$\Rightarrow \text{Efficiency} = \frac{140}{166} \Rightarrow \eta = 83.3\%$$

25. (0.5) Magnetic induction at O due to coil Y is given by,

$$B_Y = \frac{\mu_0}{4\pi} \times \frac{2\pi I(2r)^2}{[(2r)^2 + d^2]^{3/2}} \quad \dots\dots(1)$$

Similarly, the magnetic induction at O due to coil X is given by

$$B_X = \frac{\mu_0}{4\pi} \times \frac{2\pi Ir^2}{[r^2 + (d/2)^2]^{3/2}} \quad \dots\dots(2)$$

From eq. (1) & (2) $\frac{B_Y}{B_X} = \frac{1}{2}$

CHEMISTRY



26. (b) The molecule 2,3 - pentadiene does not have any chiral C but at the same time it does not have any mirror plane which makes the molecule chiral.

27. (a) In such a case there is no change in velocity

$$u = \sqrt{(3RT/M)} = \sqrt{(3PV/M)}$$


The increase in temperature = 4 times &

also the increase in pressure = 4 times. Both of these reinforce each other




29. (d) In N_2^+ , there is one unpaired electron hence it is paramagnetic.

31. (c) We know that




32. (c) Tertiary amine do not have hydrogen atom to replace Cl. Hence it will not form an amide with acid chloride.
33. (a) No. of dipeptides = 2^n ; n= no. of amino acids i.e., $2^2 = 4$ dipeptides can be formed.
34. (d) Tertiary halides on treatment with base, such as sodium methoxide, readily undergo elimination resulting in the formation of alkenes. (Williamson's Synthesis)



35. (b) $s_0 = \sqrt{K_{\text{sp}}}; s_1 = K_{\text{sp}} / 0.02 \text{ M}; s_2 = K_{\text{sp}} / 0.01 \text{ M}; s_3 = K_{\text{sp}} / 0.05 \text{ M}$


Obviously $s_0 > s_2 > s_1 > s_3$

36. (b) $\text{C}_6\text{H}_5\text{NH}_2 + \text{NaNO}_2 + \text{HCl} \xrightarrow[0^\circ\text{C}]{\quad} \text{C}_6\text{H}_5\text{N}=\text{NCl}$



37. (c) Multiple bonds formation tendency with carbon and nitrogen decreases from sulphur to tellurium.
 CS_2 ($\text{S}=\text{C}=\text{S}$) is moderately stable,
 CSe_2 ($\text{Se}=\text{C}=\text{Se}$) decomposes readily whereas,
 CTe_2 ($\text{Te}=\text{C}=\text{Te}$) does not exist
38. (d) Liquation is the principle based on difference in melting points.
39. (c) In NO_2^+ odd (unpaired) electron is removed. In peroxides (O_2^{2-}) no unpaired electrons are present as the antibonding pi M.O.'s acquired one more electron each for pairing. AlO_2^- containing Al^{3+} ($2s^2 p^6$)

40. (a) The two solutions are isotonic hence there will be no movement of H_2O .
 41. (b) Hydrated $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ is pink coloured and contains octahedral $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$ ions. If this is partially dehydrated by heating, then blue coloured tetrahedral ions $[\text{Co}(\text{H}_2\text{O})_4]^{2+}$ are formed.



42. (c) In $(\text{NH}_4)_2[(\text{TiCl}_6)]$, $\text{Ti}^{4+}(3\text{d}^0 4\text{s}^0)$ has no unpaired electrons.
 In $\text{K}_2\text{Cr}_2\text{O}_7$, $\text{Cr}^{6+}(3\text{p}^6 \text{d}^0)$ has no unpaired electrons.
 In CoSO_4 , $\text{Co}^{2+}(\text{d}^7)$ has unpaired electrons in *d*-orbitals, so it is both paramagnetic and coloured.
 In $\text{K}_3[\text{Cu}(\text{CN})_4]$, $\text{Cu}^+(3\text{d}^{10})$, has no unpaired electron.

43. (b) $\log K = \log A - \frac{E_a}{2.303R} \frac{1}{T}$ (Arrhenius equation)

Plot of $\log K$ Vs $1/T$ gives a straight line with slope $-E_a/2.303R$

44. (b) • Li does not form peroxide or superoxide due to its small size.
 • Solubility of carbonates and bicarbonates increases on moving down the group.
 • The increasing order of size of hydrated ions of alkali metals is $\text{Li}^+ > \text{Na}^+ > \text{K}^+ > \text{Rb}^+ > \text{Cs}^+$
 • Cesium used in photoelectric cells due to its low I.E. Hence statement (b) is the only correct choice.

45. (d) Cell reaction $\text{Zn} + \text{Cu}^{++} \longrightarrow \text{Zn}^{++} + \text{Cu}$

$$E_1 = E_{\text{cell}}^\circ - \frac{0.059}{2} \log \frac{0.01}{1.0} \quad \therefore E_1 = (E_{\text{cell}}^\circ + 0.059) \text{ V}$$

$$E_2 = E_{\text{cell}}^\circ - \frac{0.059}{2} \log \frac{1.0}{0.01}$$

$$\therefore E_2 = (E_{\text{cell}}^\circ - 0.059) \text{ V} . \text{ Thus, } E_1 > E_2.$$

46. (69.60) $\frac{P^\circ - P_S}{P^\circ} = \frac{w/m}{W/M}; (640 - 600)/640 = wM/mW$

$$40/640 = 2.175 \times 78/\text{m} \times 39.08$$

$$m = 2.175 \times 78 \times 640 / 39.08 \times 40 = 69.458 \cong 69.60$$

47. (0.17) $\Delta x = (h/4\pi) \times m \times \Delta v$

48. (3.8) $K = \frac{[\text{H}_3\text{O}^+][\text{HCO}_3^-]}{[\text{CO}_2][\text{H}_2\text{O}]^2}$ As pH = 6.0 $[\text{H}_3\text{O}^+] = 10^{-6}$

$K = \frac{[\text{H}_3\text{O}^+][\text{HCO}_3^-]}{[\text{CO}_2][\text{H}_2\text{O}]^2}$ (H_2O is in excess, therefore its conc. remains constant)

$$\frac{[\text{HCO}_3^-]}{[\text{CO}_2]} = \frac{K}{[\text{H}_3\text{O}^+]} = \frac{3.8 \times 10^{-6}}{10^{-6}} = 3.8$$

49. (136800) v for hydrogen like species

$$= v_{\text{H}} \times Z^2 = 15200 \times 3^2 = 15200 \times 9 = 136800 \text{ cm}^{-1}$$

50. (32.06) Calorific value of butane = $\frac{\Delta H_c}{\text{mol. wt.}} = \frac{2658}{58} = 45.8 \text{ kJ/g}$

Cylinder consist 14 Kg of butane means 14000 g of butane

$$\therefore 1\text{g gives} = 45.8 \text{ kJ/g}$$

$$\therefore 14000 \text{ g gives} = 14000 \times 45.8 = 641200 \text{ kJ}$$

Family need 20,000 kJ/day

So gas full fill the requirement for $\frac{641200}{20,000} = 32.06 \text{ days}$

MATHEMATICS

51. (b) $(a^2 + b^2 + c^2)p^2 - 2(ab + bc + cd)p + b^2 + c^2 + d^2 \leq 0$

$$\Rightarrow (a^2 p^2 - 2abp + b^2) + (b^2 p^2 - 2bcp + c^2) + (c^2 p^2 - 2cdp + d^2) \leq 0$$

$$\Rightarrow (ap - b)^2 + (bp - c)^2 + (cp - d)^2 \leq 0$$

$$\Rightarrow ap - b = 0, bp - c = 0 \text{ & } cp - d = 0$$

52. (c)

$$(a) \log(a+2b) = \frac{1}{2} \log(a+2b)^2$$

$$= \frac{1}{2} \log(a^2 + 4b^2 + 4ab)$$

$$= \frac{1}{2} \log(12ab + 4ab)$$

$$= \frac{1}{2} \log(2^4 \cdot ab)$$

$$= \frac{1}{2}(4 \log 2 + \log a + \log b)$$

$$(b) \text{ Let } \frac{\log x}{b-c} = \frac{\log y}{c-a} = \frac{\log z}{a-b} = k$$

$$\Rightarrow \log x = k(b-c), \log y = k(c-a),$$

$$\log z = k(a-b)$$

$$\therefore x^a \cdot y^b \cdot z^c = p^{k[a(b-c)] + b(c-a) + c(a-b)]}$$

$$= p^{k(0)} = 1$$

where p is any arbitrary base of the log.

(c) Given expression

$$= \log_{xyz} xy + \log_{xyz} yz + \log_{xyz} zx$$

$$= \log_{xyz}(xy \cdot yz \cdot zx) = \log_{xyz}(x^2 \cdot y^2 \cdot z^2)$$

$$= 2 \log_{xyz}(xyz) = 2 \times 1 = 2$$

53. (b) $\alpha + \beta + \gamma = \frac{\pi}{2} \Rightarrow \alpha + \gamma = \frac{\pi}{2} - \beta.$

so that $\cot(\alpha + \gamma) = \cot\left(\frac{\pi}{2} - \beta\right)$

$$\Rightarrow \frac{\cot \alpha \cot \gamma - 1}{\cot \alpha + \cot \gamma} = \frac{1}{\cot \beta}$$

54. (c) We know, $1 + \omega + \omega^2 + \dots + \omega^{n-1} = \frac{1 - \omega^n}{1 - \omega}$

$$\text{But } \omega^n = \cos\left(\frac{n\pi}{n}\right) + i \sin\left(\frac{n\pi}{n}\right)$$

$$= \cos \pi + i \sin \pi = -1$$

$$\text{and } 1 - \omega = 2 \sin^2 \frac{\pi}{2n} - 2i \sin \frac{\pi}{2n} \cos \frac{\pi}{2n}$$

$$= -2i \sin\left(\frac{\pi}{2n}\right) \left[\cos \frac{\pi}{2n} + i \sin \frac{\pi}{2n} \right]$$

$$\text{Thus, } 1 + \omega + \omega^2 + \dots + \omega^{n-1}$$

$$= \frac{2}{-2i \sin\left(\frac{\pi}{2n}\right) \left[\cos \frac{\pi}{2n} + i \sin \frac{\pi}{2n} \right]} = \frac{i \left(\frac{\cos \pi}{2n} - i \sin \frac{\pi}{2n} \right)}{\sin \frac{\pi}{2n}}$$

$$= \frac{i \cos \frac{\pi}{2n}}{\sin \frac{\pi}{2n}} - i^2 \frac{\sin \frac{\pi}{2n}}{\sin \frac{\pi}{2n}} = 1 + i \cot (\pi / 2n)$$

55. (a) $C_1(1, 0); C_2(0, -2)$

$$r_1 = \sqrt{1+15} = 4, \quad r_2 = \sqrt{4-3} = 1$$

$$C_1C_2 = \sqrt{1+4} = \sqrt{5}$$

$$r_1 - r_2 = 3 \Rightarrow C_1C_2 < r_1 - r_2$$

Hence, C_2 lies inside C_1 .

56. (d)

(a) We have $|AB| = |A||B|$

Also for a square matrix of order 3, $|kA| = k^3 |A|$ because each element of the matrix A is multiplied by k and hence in this case we will have k^3 common

$$\therefore |3AB| = 3^3 |A||B| = 27(-1)(3) = -81$$

(b) Since A is invertible, therefore A^{-1} exists and

$$\Rightarrow \det(A)\det(A^{-1}) = 1$$

$$\Rightarrow \det(A^{-1}) = \frac{1}{\det(A)}$$

$$\begin{aligned}(c) \quad (A+B)^2 &= (A+B)(A+B) \\&= A^2 + AB + BA + B^2 \\&= A^2 + 2AB + B^2 \text{ if } AB = BA.\end{aligned}$$

57. (b) For trivial solution,

$$\begin{vmatrix} 1 & -2 & 1 \\ 2 & -1 & 3 \\ \lambda & 1 & -1 \end{vmatrix} \neq 0 \Rightarrow -5\lambda - 4 \neq 0 \text{ or } \lambda \neq -\frac{4}{5}$$

58. (d) $f(x) = |x-1| = \begin{cases} -x+1, & x < 1 \\ x-1, & x \geq 1 \end{cases}$

Consider $f(x^2) = (f(x))^2$

If it is true it should be $\forall x$

$$\therefore \text{Put } x = 2$$

$$\text{LHS} = f(2^2) = |4-1| = 3$$

$$\text{RHS} = (f(2))^2 = 1$$

\therefore (a) is not correct

Consider $f(x+y) = f(x) + f(y)$

Put $x = 2, y = 5$ we get

$$f(7) = 6; f(2) + f(5) = 1 + 4 = 5$$

\therefore (b) is not correct

Consider $f(|x|) = |f(x)|$

Put $x = -5$ then $f(|-5|) = f(5) = 4$

$$|f(-5)| = |-5-1| = 6$$

\therefore (c) is not correct.

Hence (d) is the correct alternative.

59. (d) Let $a = \tan \theta$ and $b = \tan \phi$

$$\therefore \sin^{-1} \left[\frac{2a}{1+a^2} \right] = \sin^{-1} \left[\frac{2 \tan \theta}{1+\tan^2 \theta} \right]$$

$$\text{and } \sin^{-1} \left[\frac{2b}{1+b^2} \right] = \sin^{-1} \left[\frac{2 \tan \phi}{1+\tan^2 \phi} \right]$$

$$= \sin^{-1} [\sin 2\phi] = 2\phi = 2 \tan^{-1} b$$

$$\text{Thus, } \sin^{-1} \left[\frac{2a}{1+a^2} \right] = 2 \tan^{-1} a \text{ and}$$

$$\sin^{-1} \left[\frac{2b}{1+b^2} \right] = 2 \tan^{-1} b$$

$$\therefore 2 \tan^{-1} x = \sin^{-1} \left[\frac{2a}{1+a^2} \right] + \sin^{-1} \left[\frac{2b}{1+b^2} \right]$$

$$= 2 \tan^{-1} a + 2 \tan^{-1} b$$

$$\Rightarrow \tan^{-1} x = \tan^{-1} a + \tan^{-1} b$$

$$\tan^{-1} x = \tan^{-1} \frac{a+b}{1-ab}$$

$$\therefore x = \frac{a+b}{1-ab}$$

60. (a) We have,


$$AB = \begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix} \begin{bmatrix} \cos^2 \phi & \cos \phi \sin \phi \\ \cos \phi \sin \phi & \sin^2 \phi \end{bmatrix}$$

$$\begin{aligned} &= \begin{bmatrix} \cos^2 \theta \cos^2 \phi + \cos \theta \cos \phi \sin \theta \sin \phi \\ \cos \theta \sin \theta \cos^2 \phi + \sin^2 \theta \cos \phi \sin \phi \end{bmatrix} \\ &\quad \begin{bmatrix} \cos^2 \theta \cos \phi \sin \phi + \cos \theta \sin \theta \sin^2 \phi \\ \cos \theta \cos \phi \sin \theta \sin \phi + \sin^2 \theta \sin^2 \phi \end{bmatrix} \end{aligned}$$

$$= \cos(\theta - \phi) \begin{bmatrix} \cos \theta \cos \phi & \cos \theta \sin \phi \\ \sin \theta \cos \phi & \sin \theta \sin \phi \end{bmatrix}$$

Since $AB = 0$, $\therefore \cos(\theta - \phi) = 0$

61. (c) Since $|x|$ is not diff. at $x=0$



$\Rightarrow |x - 1|$ is not diff at $x = 1$.

$x^n |x|$ in n times diff. at $x = 0$

$\Rightarrow (x-1)^2 |x-1|$ is twice diff. at $x = 1$

but not thrice diff. at $x = 1$

62. (c) $f(x) = \frac{1}{x-1}$ is discontinuous at $x=1$.

$(gof)(x) = g(f(x)) = -\frac{(x-1)^2}{(2x-1)(x-2)}$, which is not defined at $x = 1/2, 2$.

Hence the set of points where $(gof)(x)$ is discontinuous is $\{1/2, 1, 2\}$

$$63. (a) \sum_{r=0}^m {}^{n+r} C_n = \sum_{r=0}^m {}^{n+r} C_r \quad (\because {}^{n+r} C_n = {}^{n+r} C_{n+r-n})$$

$$= {}^n C_0 + {}^{n+1} C_1 + {}^{n+2} C_2 + {}^{n+3} C_3 + \dots + {}^{n+m} C_m$$

$$\text{Using, } {}^n C_0 = 1 = {}^{n+1} C_1$$

$$= ({}^{n+1} C_0 + {}^{n+1} C_1) + {}^{n+2} C_2 + {}^{n+3} C_3 + \dots + {}^{n+m} C_m$$

$$\text{Using, } {}^n C_r + {}^n C_{r+1} = {}^{n+1} C_{r+1}$$

$$= ({}^{n+2} C_1 + {}^{n+2} C_2) + {}^{n+3} C_3 + \dots + {}^{n+m} C_m$$

Using this again and again, we are left with

$$= {}^{n+m} C_{m-1} + {}^{n+m} C_m$$

$$= {}^{n+m+1} C_m = {}^{n+m+1} C_{n+1}$$

64. (b) As $x \rightarrow \frac{1}{3}$; $\{x+1\} \rightarrow \{1+1/3\} \rightarrow 1/3$

Similarly $\{x+2\} \rightarrow \frac{1}{3}$ as $x \rightarrow \frac{1}{3}$

65. (b) Squaring both sides we get highest order as 2.

66. (d) $I = \int_{\pi/4}^{\pi/4} (x|x| + \sin^3 x + x \tan^2 x + 1) dx$
 $\downarrow \text{odd f} \quad \downarrow \text{odd f} \quad \downarrow \text{odd f}$

$$I = \int_{-\pi/4}^{\pi/4} dx = \frac{\pi}{2} \quad [\because \int_{-a}^a f(x) dx = 0, \text{ as } f(x) \text{ is an odd function}]$$

67. (b) $(1-x-2x^2)^6 = (1+x)^6(1-2x)^6$

$$= 1 + a_1 x + a_2 x^2 + \dots + a_{12} x^{12}$$

$$\text{Putting } x = 1/2, \text{ we have } 0 = 1 + a_1/2 + a_2/2^2 + a_3/2^3 + a_4/2^4 + \dots + a_{12}/2^{12} \quad \dots(1)$$

Putting $x = -1/2$, we have

$$1 = 1 - a_1/2 + a_2/2^2 - a_3/2^3 + a_4/2^4 - \dots + a_{12}/2^{12} \dots(2)$$

Adding (1) and (2), we have

$$1 = 2(1 + a_2/2^2 + a_4/2^4 + \dots + a_{12}/2^{12})$$

$$\Rightarrow a_2/2^2 + a_4/2^4 + a_6/2^6 + \dots + a_{12}/2^{12} = -1/2$$

68. (b) $y^2 = 4x \text{ & } \frac{x^2}{8} + \frac{y^2}{2} = 1$

Equation of tangent to above curves are respectively.

$$y^2 = mx + \frac{1}{m} \text{ and } y = mx + \sqrt{8m^2 + 2}$$

$$\text{Comparing } \frac{1}{m} = \sqrt{8m^2 + 2}$$

$$\Rightarrow m^2(8m^2 + 2) = 1$$

seeing the options

$$m = \pm \frac{1}{2} \text{ satisfy the equation}$$

$$\Rightarrow y = \pm \frac{1}{2}x \pm 2 \Rightarrow 2y = \pm x \pm 4$$

$$\text{i.e. } 2y = x + 4 \text{ & } x + 2y + 4 = 0$$

69. (b) Let the point be (x_1, y_1) .

$2(x_1 - 3)$, but it is equal to 1.

$$\text{Therefore, } 2(x_1 - 3) = 1 \Rightarrow x_1 = \frac{7}{2}$$

$$y_1 = \left(\frac{7}{2} - 3\right)^2 = \frac{1}{4}.$$

Hence the point is $\left(\frac{7}{2}, \frac{1}{4}\right)$.

70. (c) $x = \tan A$, $y = \tan B$, $-z = \tan C$. Then $(x + y - z) = -xyz$.

$$\Rightarrow \tan A + \tan B + \tan C = \tan A \tan B \tan C$$

$$\Rightarrow A + B + C = \pi \Rightarrow 2A + 2B = 2\pi - 2C$$

$$\Rightarrow \tan(2A + 2B) = \tan(2\pi - 2C) = -\tan 2C$$

$$\Rightarrow \tan 2A + \tan 2B + \tan 2C = \tan 2A \cdot \tan 2B \cdot \tan 2C$$

$$\Rightarrow \frac{2 \tan A}{1 - \tan^2 A} + \frac{2 \tan B}{1 - \tan^2 B} + \frac{2 \tan C}{1 - \tan^2 C}$$

$$= \frac{2 \tan A}{1 - \tan^2 A} \cdot \frac{2 \tan B}{1 - \tan^2 B} \cdot \frac{2 \tan C}{1 - \tan^2 C}$$

Put the value of $\tan A$, $\tan B$, $\tan C$, we get

$$\Rightarrow \frac{2x}{1-x^2} + \frac{2y}{1-y^2} - \frac{2z}{1-z^2}$$

$$= -\frac{8xyz}{(1-x^2)(1-y^2)(1-z^2)}$$

71. (12.25) $4p^2 + 4p + 12 = 0 \Rightarrow p = -7$

The equation $x^2 + px + q = 0$ has equal roots then $D = 0$.

$$\text{or } p^2 = 4q \Rightarrow q = \frac{49}{4}$$

72. (0.125)

$$\cos 36^\circ \cos 42^\circ \cos 78^\circ$$

$$= \cos 36^\circ \cos(60^\circ - 18^\circ) \cos(60^\circ + 18^\circ)$$

$$= \frac{\sqrt{5} + 1}{4} \left(\cos^2 60^\circ - \sin^2 18^\circ \right)$$

$$\begin{aligned}
&= \left(\frac{\sqrt{5}+1}{4} \right) \frac{1}{4} - \left(\frac{\sqrt{5}+1}{4} \right) \left(\frac{5+1-2\sqrt{5}}{16} \right) \\
&= \left(\frac{\sqrt{5}+1}{16} \right) - \frac{(\sqrt{5}+1)(\sqrt{5}-1)^2}{64} = \frac{\sqrt{5}+1}{16} \left[1 - \frac{(\sqrt{5}-1)^2}{4} \right] \\
&= \frac{\sqrt{5}+1}{16} \left[\frac{4-6+2\sqrt{5}}{4} \right] = \frac{1}{8}
\end{aligned}$$

73. (0.2) The given expression is equal to

$$\begin{aligned}
&\cos(\cos^{-1} x + \sin^{-1} x + \sin^{-1} x) \\
&= \cos\left(\frac{\pi}{2} + \sin^{-1} x\right) = -\sin(\sin^{-1} x) = -x = -\frac{1}{5} \\
&[\text{Using } \cos^{-1} x + \sin^{-1} x = \frac{\pi}{2}]
\end{aligned}$$

74. (4) Let $\tan\theta_1, \tan\theta_2$ be the roots of the equation $2\tan^2\theta - 4\tan\theta + 1 = 0$. Thus $\tan\theta_1 + \tan\theta_2 = 4/2 = 2$; $\tan\theta_1 \tan\theta_2 = 1/2$.

$$\begin{aligned}
&\text{Now } \tan(\theta_1 + \theta_2) = [(\tan\theta_1 + \tan\theta_2)/(1 - \tan\theta_1 \tan\theta_2)] \\
&= 2/[1 - (1/2)] = 4.
\end{aligned}$$

$$\begin{aligned}
75. (2.25) \quad &\text{Given, in } \Delta ABC \begin{vmatrix} 1 & a & b \\ 1 & c & a \\ 1 & b & c \end{vmatrix} = 0
\end{aligned}$$

$$\begin{aligned}
&\Rightarrow 1(c^2 - ab) - a(c - a) + b(b - c) = 0 \\
&\Rightarrow a^2 + b^2 + c^2 - ab - bc - ca = 0 \\
&\Rightarrow 2a^2 + 2b^2 + 2c^2 - 2ab - 2bc - 2ca = 0 \\
&\Rightarrow (a^2 + b^2 - 2ab) + (b^2 + c^2 - 2bc) + (c^2 + a^2 - 2ca) = 0 \\
&\Rightarrow (a - b)^2 + (b - c)^2 + (c - a)^2 = 0
\end{aligned}$$

Here, sum of squares of three members can be zero if and only if $a = b = c$

$\Rightarrow \Delta ABC$ is equilateral.

$$\Rightarrow \angle A = \angle B = \angle C = 60^\circ$$

$$\therefore \sin^2 A + \sin^2 B + \sin^2 C$$

$$-(\sin^2 60^\circ + \sin^2 60^\circ + \sin^2 60^\circ)$$

MOCK TEST-3

PHYSICS

1. (c) Resolving power of eye = λ / a

$$= \frac{500 \times 10^{-9}}{5 \times 10^{-3}} = 10^{-4} \text{ radians}$$

Now, arc = angle \times radius

$$= 10^{-4} \times (500 \times 10^3) \text{ m} = 50 \text{ m}$$

2. (b) Frequency doesn't depend on medium

$$\frac{\mu_1}{\mu_2} = \frac{V_2}{V_1} = \frac{\lambda_2 f}{\lambda_1 f}, \text{ or } \mu_2 \lambda_2 = \lambda_1 \mu_1$$

$$\mu_2 = 3680 \text{ \AA}$$

3. (a) Total momentum will be conserved.

Initial momentum = Final momentum

$$M.v = m \times 0 + (M - m)v'$$

$$\therefore v' = \frac{Mv}{M - m}$$

4. (c) Force, $F = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^2}$

$$\Rightarrow \epsilon_0 = \frac{q_1 q_2}{4\pi F r^2}$$

So dimension of ϵ_0

$$= \frac{[AT]^2}{[MLT^{-2}][L^2]} = [M^{-1}L^{-3}T^4A^2]$$

5. (c) Heat radiated by black body

$$E = \sigma A T^4 \Rightarrow E \propto T^4$$

$$\text{or } \frac{E_1}{E_2} = \frac{T_1^4}{T_2^4}$$

$$\text{or } \frac{20}{E_2} = \left(\frac{500}{1000} \right)^4 = \left(\frac{1}{2} \right)^4 = \frac{1}{16}$$

$$\Rightarrow E_2 = 16 \times 20 \text{ cal m}^{-2} \text{ s}^{-1}$$
$$= 320 \text{ cal m}^{-2} \text{ s}^{-1}$$

6. (c) Velocity of wave $v = n\lambda$

$$n_2 = \frac{v_2}{\lambda_2} = \frac{396}{100 \times 10^{-2}} = 396 \text{ Hz}$$

$$\text{no. of beats} = n_1 - n_2 = 4$$

7. (a) Terminal velocity attained by falling object

$$V_t = \frac{3r^2(d - \rho)g}{a\eta}$$

$$\text{thus, } V_t \propto r^2$$

8. (d) Young's modulus, $Y = \frac{\text{Stress}}{\text{Strain}}$

$$\text{or stress} = Y \cdot \text{strain}$$

$$\text{or strain} = \text{Stress} / Y$$

$$\text{or } \Delta l = \frac{Fl}{YA}; \frac{\Delta l_1}{\Delta l_2} = \frac{F_1 l_1}{A_1 Y_1} \cdot \frac{A_2 Y_2}{F_1 l_1}$$

$$l_1 = l_2 \text{ & } Y_1 = Y_2, F_1 = F_2$$

$$\Rightarrow \frac{\Delta l_1}{\Delta l_2} = \frac{\pi r_2^2}{\pi r_1^2} = \frac{4r^2}{r^2} = 4$$

9. (b) $E = mc^2 = (2 \times 1.6 \times 10^{-27}) \times (3 \times 10^8)^2$

$$= 28.8 \times 10^{-27} \times 10^{16} \text{ J} = 28.8 \times 10^{-11} \text{ J}$$

$$= 28.8 \times 10^{-10} \text{ J}$$


10. (b) $y = 2 \sin\left(\frac{\pi t}{2} + \phi\right)$

$$\text{velocity of particle } \frac{dy}{dt} = 2 \times \frac{\pi}{2} \cos\left(\frac{\pi t}{2} + \phi\right)$$

$$\text{acceleration } \frac{d^2y}{dt^2} = -\frac{\pi^2}{2} \sin\left(\frac{\pi t}{2} + \phi\right)$$

$$\text{Thus, } a_{\max} = \frac{\pi^2}{2}$$

11. (a)



Electric field inside

- 12. (a)** Let the body be depressed by distance x from its equilibrium position. The extra upthrust created is $x \rho A g$ which applies to whole body. If a be acceleration created then,

$$x \rho A g = m g a \Rightarrow a = \frac{\rho A}{m} x$$

Since, acceleration $\propto x$. So it is equation of S.H.M.

$$\text{So, } \omega^2 = \frac{\rho A}{m} \Rightarrow T = 2\pi \sqrt{\frac{m}{\rho A}}$$

$$T \propto \frac{1}{\sqrt{A}}$$

- 13. (b)** The specific resistance (ρ) is determined by the formula


$$\rho = \frac{\pi D^2}{4L}$$

where symbols have their usual meaning.

- 14. (b)** Clearly the co-ordinates of A are $(2f, 2f)$

$$\therefore f = \frac{40}{2} = 20 \text{ cm.}$$


- 15. (a)** Number of electrons falling on the metal plate A
 $= 10^{16} \times (5 \times 10^{-4})$



\therefore Number of photoelectrons emitted from metal plate A upto 10 seconds is

$$n_e = \frac{(5 \times 10^{-4}) \times 10^{16}}{10^6} \times 10 = 5 \times 10^7$$

- 16. (b)** line 1 line 2 line 3



In steady state capacitor is fully charged hence no current will flow through line 2.

By simplifying the circuit



Hence resultant potential difference across resistances will be 8.0 V.

$$\text{Thus current } I = \frac{V}{R} = \frac{8.0}{3+9} = \frac{8}{12}$$

$$\text{or, } I = \frac{2}{3} = 0.67 \text{ A}$$

- 17. (c)** Given: Amplitude of electric field,

$$E_0 = 4 \text{ v/m}$$

Absolute permittivity,

$$\epsilon_0 = 8.8 \times 10^{-12} \text{ C}^2/\text{N}\cdot\text{m}^2$$

Average energy density $u_E = ?$

Applying formula,

$$\text{Average energy density } u_E = \frac{1}{4} \epsilon_0 E^2$$

$$\Rightarrow u_E = \frac{1}{4} \times 8.8 \times 10^{-12} \times (4)^2 \\ = 35.2 \times 10^{-12} \text{ J/m}^3$$

- 18. (a)** When springs are in parallel, then

$$T = 2\pi \sqrt{\frac{m}{K_1 + K_2}} \Rightarrow \frac{2\pi}{T} = \omega = \sqrt{\frac{K_1 + K_2}{m}}$$

- 19. (c)** Lateral magnitude = v/u ;

$$\text{Mag. along axis} = \left| \frac{dv}{du} \right| = \frac{u^2}{v^2} = 1 \text{ if } v = u, \therefore u = 2$$

- 20. (d)** In linear S.H.M., the restoring force acting on particle should always be proportional to the displacement of the particle and directed towards the equilibrium position.

21. (6) According to Doppler's effect

$$f = \left(\frac{v \pm v_0}{v \pm v_s} \right) f$$

here $v_0 = 0$ and $v_s = 0.5v$

$$\therefore f = \left(\frac{v}{v - .5v} \right) 3 = 6 \text{ kHz}$$

22. (300) $\eta = \left(1 - \frac{T_c}{T_H} \right) \times 100$

$$\Rightarrow 70 = \left(1 - \frac{T_c}{1000} \right) \times 100$$

$$0.7 = 1 - \frac{T_c}{1000}$$

$$\therefore \frac{T_c}{1000} = 0.3 \text{ or } T_c = 300\text{K.}$$

23. (11.2) Escape velocity $v_e = \sqrt{2gR}$

thus, it doesn't depend on mass.

24. (0.4)r=5 cm. = $5 \times 10^{-2}\text{m}$

$$B_E = 0.5 \times 10^{-5} \text{ W/m}^2$$


we know that field due to coil at

$$\text{centre } B = \frac{\mu_0 I}{2r}$$

it annuals the earth's magnetic field

$$\text{So, } \frac{\mu_0 I}{2r} = 0.5 \times 10^{-5}$$

$$I = \frac{2R \times 0.5 \times 10^{-5}}{\mu} = \frac{5}{4\pi} A = 0.4A$$



25. (1.2×10^{-7}) Pressure of light on totally reflecting surface



$$P = \frac{2I}{C} \quad (C = \text{velocity of light})$$

$$P = \frac{F}{A} = \frac{2I}{C}$$

$$\Rightarrow F = \frac{2IA}{C} = \frac{2 \times 12 \times 1.5 \times 10^{-4}}{4} \text{ N}$$

CHEMISTRY

26. (d) (a) $\begin{array}{c} \text{CH}_2\text{Cl} \\ | \\ \text{CH}_2\text{Cl} \end{array} \xrightarrow[\text{alcohol}]{\text{KOH}} \begin{array}{c} \text{CH} \\ || \\ \text{CH} \end{array};$
- $\begin{array}{c} \text{CH}_3\text{Cl} \\ | \\ \text{CH} \end{array} \xrightarrow[\text{alcohol}]{\text{KOH}} \begin{array}{c} \text{CH} \\ || \\ \text{CH} \end{array}; \text{ hence true}$
- (b) Both are position isomers
 (c) Since, they are isomers, percentage of C, H and Cl in both will be same.



Hence, statement (d) is wrong.

27. (d) Sodium carbonate is the salt of a weak acid (H_2CO_3) with a strong base (NaOH). In solution, it is completely ionised as Na^+ and CO_3^{2-} ions. The CO_3^{2-} ion being the conjugate base of the weak acid H_2CO_3 undergoes hydrolysis in solution according to the equilibrium of hydrolysis:




The OH^- ion produced being a strong base makes the solution basic, hence the pH of the solution will be greater than 7.

28. (a) According to the kinetic theory of gases, the average velocity of the molecules in the gas is given by the expression $v = \sqrt{\frac{8RT}{\pi M}}$ where T is the absolute temperature and R is the gas constant. Thus the average velocity can be taken as proportional to the square root of the absolute temperature. Hence the ratio of the average velocity at 200°C to that at

50°C will be equal to $\sqrt{\frac{273 + 200}{273 + 50}}$ which is 1.21.

29. (d) No. of atoms of hydrogen in 0.046 g of alcohol
- $$= \frac{0.046}{46} \times 6 \times 10^{23} \times 6$$
- $$= 1 \times 10^{-3} \times 6 \times 10^{23} \times 6 = 3.6 \times 10^{21}$$

30. (b) Since the gas B turns CuSO_4 solution blue, it can be NH_3 .



31. (c) (i) $\text{CH}_3 - \text{C}(=\text{O})\text{H} \xrightarrow{\substack{\text{(i)} \text{CH}_3\text{MgBr} \\ \text{(ii)} \text{H}_3\text{O}^+}} \text{CH}_3 - \text{C}(\text{H})(\text{OH})$
 2-Propanol
- (ii) $\text{CH}_3 - \text{C}(\text{H})(\text{OH}) + \text{C}_2\text{H}_5\text{OH} \xrightarrow{\text{HCl}}$
- $\begin{array}{c} \text{H} \\ | \\ \text{CH}_3 - \text{C}(\text{H})(\text{OC}_2\text{H}_5) \\ | \\ \text{OH} \end{array} \xrightarrow{\substack{\text{HCl} \\ \text{C}_2\text{H}_5\text{OH}}} \begin{array}{c} \text{H} \\ | \\ \text{CH}_3 - \text{C}(\text{H})(\text{OC}_2\text{H}_5)_2 \\ | \\ \text{Acetal} \end{array}$
32. (b) The temperature of 383 K is equal to 110°C. Although the salts will increase the boiling point of water, it should boil at or below this temperature.
33. (b) Since each of (i), (ii) and (iii) are hexa-coordinated, in the case of (ii), one of the chlorines (chloride ions) is coordinated to the central cobalt ion and in (iii), two such chlorides are coordinately linked. Thus, the ionisable chlorides in (i) is three in (ii) it is two and in (iii) it is only one.
 Primary valency means the valency of the complex cation.
34. (b) Polyethylene or Polyethene, $\left[\text{CH}_2 - \text{CH}_2 \right]_n$, is made from a single monomer, it is a homopolymer.
35. (d) Energy requirements of the body are met by glucose that is circulated through blood and glycogen stored in the muscles.
36. (c) More the *s*-character, more is the stability of the carbanion. hence the correct order is $sp > sp^2 > sp^3$.
37. (a) The total number of electrons in the molecular species given, respectively are 17, 16 and 18. Write down the electronic configuration of the molecular species and observe the number of electrons in antibonding orbitals which are respectively 7, 6 and 8.
38. (c) Deviation from ideal gas behaviour is greater, when the pressure is higher and the gas is closed to its liquefaction point or its critical temperature. Thus, the conditions of -100°C and 4 atm pressure among the sets given causes maximum deviation.
39. (b) Heat of neutralisation of strong acid and strong base is always 13.7 kcal.
40. (b) Average atomic weight = $85\left(\frac{75}{100}\right) + 87\left(\frac{25}{100}\right) = 85.5$.
41. (b) In Kjeldahl's method of estimation of nitrogen, the nitrogen present in most of the organic compounds is quantitatively converted into ammonium sulphate. The $(\text{NH}_4)_2\text{SO}_4$ so obtained is decomposed with excess of NaOH solution to give NH_3 which is absorbed in an excess of cooled HCl . H_2SO_4 reacts with Na_2CO_3 to give Na_2SO_4 which is titrated with

43. (a) Addition of a catalyst to a reaction mixture has the effect of lowering the activation energy of the reaction by changing the path or mechanism of the reaction. The reaction rate increases manifold. However, the equilibrium constant and the enthalpy (ΔH) of the reaction are unaffected.
44. (c) Be(OH)₂ is amphoteric that means it can react with both acids and alkalies
45. (d) Lithium, sodium and potassium are highly electropositive and highly reactive metals. When any of these come in contact with water, the reaction is so swift and intense that the hydrogen evolved catches fire instantaneously. The reaction thus is doubly exothermic, using water to quench fires caused by these metals makes it explosively dangerous. Likewise CO₂ and nitrogen too are reactive. Small fires can be quenched by asbestos blanket or by covering with dry sand, since these measures prevent contact with oxygen and water vapour and thus become effective.

46. (0.107) $N_1 V_1 = N_2 V_2$

$$N_{NaOH} = M_{NaOH} = 0.164$$

$$\Rightarrow 25 \times N = 32.63 \times 0.164$$

$$N = \frac{32.63 \times 0.164}{25} = 0.214 \text{ N}$$

$$\text{But } N_{H_2SO_4} = 2 \times M_{H_2SO_4}$$

$$\Rightarrow M = \frac{\text{Normality}}{2} = \frac{0.214}{2} = 0.107$$

47. (50) Eq. of KMnO₄ used = $\frac{50 \times 1}{1000 \times 10} = 0.005$

$$\therefore \text{Eq of FAS reacted} = 0.005$$

$$\begin{aligned}\therefore \text{weight of FAS needed} &= 0.005 \times 392 \\ &= 1.96 \text{ g}\end{aligned}$$

Thus, percentage purity of FAS is 50%.

48. (4) In a 'fcc' crystal atoms are located at the centre of the 6 faces and at the 8 corners.

On each face there is 1 atom which is shared by 2 cells. Hence, the no. of atoms/unit cell = $6/2 = 3$



Again the corner atom is shared by 8 other cells. Hence no. of atoms = $8/8 = 1$

No. of atoms/unit cell = $1 + 3 = 4$

49. (800) As AgNO₃ dissociates completely, therefore in 0.1 M AgNO₃ solution, [Ag⁺] = 0.1 M



$$\begin{aligned}
 K_{\text{sp}} &= [\text{Ag}^+]^2 [\text{CO}_3^{2-}] \\
 &= 8 = (0.1 + 2s)^2 \times s \\
 &= 0.01 s = 8 ; (0.1 + 2s \approx 0.1) \\
 s &= 800
 \end{aligned}$$



$$2x + 5(5-x) = 16 \Rightarrow x = 3 \text{ L}$$

$$\therefore \text{Heat released} = \frac{3}{22.4} \times 890 + \frac{2}{22.4} \times 2220 = 317.$$

MATHEMATICS

51. (d) Given $f(x) = \tan^{-1}(\sin x + \cos x)$

$$f'(x) = \frac{1}{1 + (\sin x + \cos x)^2} \cdot (\cos x - \sin x)$$

$$= \frac{\sqrt{2} \left(\frac{1}{\sqrt{2}} \cos x - \frac{1}{\sqrt{2}} \sin x \right)}{1 + (\sin x + \cos x)^2}$$

$$\therefore f'(x) = \frac{\sqrt{2} \cos \left(x + \frac{\pi}{4} \right)}{1 + (\sin x + \cos x)^2}$$

if $f'(x) > 0$ then $f(x)$ is increasing function.
Hence $f(x)$ is increasing, if

$$-\frac{\pi}{2} < x + \frac{\pi}{4} < \frac{\pi}{2} \Rightarrow -\frac{3\pi}{4} < x < \frac{\pi}{4}$$

Hence, $f(x)$ is increasing when $x \in \left(-\frac{\pi}{2}, \frac{\pi}{4} \right)$

52. (a) $\sqrt{1+x^2} + \sqrt{1+y^2} = \lambda(x\sqrt{1+y^2} - y\sqrt{1+x^2})$

$$\Rightarrow \sqrt{1+x^2}(1+\lambda y) = \sqrt{1+y^2}(\lambda x - 1)$$

$$\begin{aligned}
&\Rightarrow \frac{x^2+1}{y^2+1} = \frac{\lambda^2 x^2 - 2\lambda x + 1}{\lambda^2 y^2 + 2\lambda y + 1} \\
&\Rightarrow (y^2+1)(\lambda^2 x^2 - 2\lambda x + 1) \\
&\quad = (x^2+1)(\lambda^2 y^2 + 2\lambda y + 1) \\
&\Rightarrow \lambda^2 x^2 y^2 - 2\lambda x y^2 + y^2 + \lambda^2 x^2 - 2\lambda x + 1 \\
&\quad = \lambda^2 x^2 y^2 + 2\lambda x^2 y + x^2 + \lambda^2 y^2 + 2\lambda y + 1 \\
&\Rightarrow \lambda^2(x^2 - y^2) - 2\lambda(xy^2 + x^2 y + x + y) = 0 \\
&\Rightarrow \lambda^2(x+y)(x-y) - 2\lambda [xy(x+y) + (x+y)] = 0 \\
&\Rightarrow \lambda(x+y)[\lambda(x-y) - 2xy - 2] = 0 \\
&\Rightarrow (x+y)[\lambda(x-y) - 2xy - 2] = 0 \\
&\Rightarrow \lambda(x-y) - 2xy - 2 = 0 \\
&\Rightarrow \frac{2xy+2}{x-y} = \lambda \Rightarrow \frac{xy+1}{x-y} = \frac{\lambda}{2} \\
&\Rightarrow \frac{\left(x \frac{dy}{dx} + y\right)(x-y) - (xy+1)\left(1 - \frac{dy}{dx}\right)}{(x-y)^2} = 1
\end{aligned}$$

This is the first order differential equation and clearly degree of $\frac{dy}{dx}$ is 1.

Hence degree of the differential equation is 1.

53. (c) $I = \int_0^2 [x^2] dx$

The function $[x^2]$ varies as follows between $x = (0, 2)$

$$[x^2] = \begin{cases} 0 & \text{if } 0 \leq x^2 < 1, \text{ or } 0 \leq x < 1 \\ 1 & \text{if } 1 \leq x^2 \leq 2 \text{ or } 1 \leq x \leq \sqrt{2} \end{cases}$$

$$\Rightarrow I = \int_0^1 0 \cdot dx + \int_1^{\sqrt{2}} 1 \cdot dx + \int_{\sqrt{2}}^{\sqrt{3}} 2 \cdot dx + \int_{\sqrt{3}}^2 3 \cdot dx$$

$$= 0 + (\sqrt{2} - 1) + 2(\sqrt{3} - \sqrt{2}) + 3(2 - \sqrt{3})$$

$$= \sqrt{2} - 1 + 2\sqrt{3} - 2\sqrt{2} + 6 - 3\sqrt{3} = 5 - \sqrt{2} - \sqrt{3}$$

54. (c) $\alpha + \beta = 3; \alpha\beta = a; \gamma + \delta = +12; \gamma\delta = b$

$\alpha, \beta, \gamma, \delta$ are in increasing G.P.

$$\beta = \alpha x, \gamma = \alpha x^2, \delta = \alpha x^3$$

$$\alpha + \beta = \alpha + \alpha x = 3 = \alpha(1+x) \quad \dots\dots(1)$$

$$\gamma + \delta = \alpha x^2 + \alpha x^3 = 12 = \alpha x^2(1+x) \quad \dots\dots(2)$$

$$\text{Divding } \frac{3}{12} = \frac{\alpha(1+x)}{\alpha x^2(1+x)} \text{ or } \frac{1}{4} = \frac{1}{x^2} \text{ or } x = 2$$

$$\Rightarrow \beta = 2\alpha \text{ and } \alpha + 2\alpha = 3 \Rightarrow \alpha = 1 \text{ and } \beta = 2$$

$$\therefore a = \alpha\beta = 2$$

$$\gamma = \alpha x^2 = 1 \times 2^2 = 4; \delta = \alpha x^3 = 1 \times 2^3 = 8$$

$$\therefore b = \gamma\delta = 4 \times 8 = 32$$

55. (a) Consider the example: Let $A = \{1, 2, 3\}$,

$$R = \{(1, 1), (1, 2)\} \text{ and } S = \{(2, 2), (2, 3)\}$$

Clearly R and S are transitive relations on A.

$$R \cup S = \{(1, 1), (2, 2), (1, 2), (2, 3)\}$$

$R \cup S$ is not transitive as $(1, 3) \notin R \cup S$.

56. (d) $I = \int \log 2x \, dx = \int \log 2x \cdot 1 \, dx$


Using Integration by parts

$$I = \log 2x \cdot x - \int \frac{2}{2x} \cdot \int 1 \, dx$$

57. (a) For $k=0$,

it is obvious from the given interval that graph will be increasing from -1 to 1

Similar graphs can be obtained for all values of k .



58. (a) It can be also solved by comparing

with the linear equation $\frac{dy}{dx} + Py = Q$

The integrating factor, I.F. = $e^{\int 1 dx} = e^x$

$$\text{Therefore, } y \cdot \text{I.F.} = \int 2e^{2x} \cdot \text{I.F.} + C$$

$$y \cdot e^x = \int 2e^{2x} \cdot e^x + C$$

$$y \cdot e^x = 2 \int e^{3x} + C = \frac{2}{3} e^{3x} + C \Rightarrow y = \frac{2e^{2x}}{3} + ce^{-x}$$

$$59. \quad (c) \quad \Delta = \begin{vmatrix} a+x & a-x & a-x \\ a-x & a+x & a-x \\ a-x & a-x & a+x \end{vmatrix} = 0$$

$$\Rightarrow \Delta = \begin{vmatrix} 3a-x & a-x & a-x \\ 3a-x & a+x & a-x \\ 3a-x & a-x & a+x \end{vmatrix}, \quad C_1 \rightarrow C_1 + C_2 + C_3$$

$$= (3a-x) \begin{vmatrix} 1 & a-x & a-x \\ 1 & a+x & a-x \\ 1 & a-x & a+x \end{vmatrix} = 0$$

Using $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 - R_1$

$$\Rightarrow \Delta = (3a-x) \begin{vmatrix} 1 & a-x & a-x \\ 0 & 2x & 0 \\ 0 & 0 & 2x \end{vmatrix} = 0$$

$$\text{or, } 4x^2(3a-x) = 0 \Rightarrow x = 0 \text{ or } 3a$$

60. (c) We know that we divide the equation in the ratio $3 : 1$

$$= \frac{2}{3} \times 3a = 2a$$

Centre of the (given) circle is C(0, 0). Therefore the equation of the circle
 $(x - 0)^2 + (y - 0)^2 = (2a)^2 \Rightarrow x^2 + y^2 = 4a^2$

- 61. (b)** Its contrapositive is ‘sum of digits of n is not divisible by 9’
 $\Rightarrow n$ is not divisible by 9

- 62. (c)** $n = 7$

Prob. of getting any no. out 1, 2, 3, ... 9 is $p = 9/15$

$$\therefore q = 6/5$$

$$P(x = 7) = {}^7C_7 p^7 q^0 \quad [\text{Binomial distribution}]$$

$$= \left(\frac{9}{15}\right)^7 = \left(\frac{3}{5}\right)^7$$

- 63. (a)** For $x \geq a$, the equation becomes

$$x^2 - 2a(x-a) - 3a^2 = 0 \Rightarrow x = (1 + \sqrt{2})a, (1 - \sqrt{2})a$$

for $x \leq a$, the equation becomes

$$x^2 - 2a[-(x-a)] - 3a^2 = 0 \Rightarrow x^2 + 2ax - 5a^2 = 0$$

$$\Rightarrow x = -(1 + \sqrt{6})a, (-1 + \sqrt{6})a$$

This shows $(-1 + \sqrt{6})a$ is one of the roots.

- 64. (c)** $X \cap (X \cup Y)^c = X \cap (X^c \cap Y^c) = (X \cap X^c) \cap Y^c$

$$= \emptyset \cap Y^c = \emptyset$$


- 65. (a)** $y = \log_2 \{\log_2(x)\} = \log_2 \{\log_e x \cdot \log_2 e\}$

$$\Rightarrow y = \log_e \{\log_e x \cdot \log_2 e\} \cdot \log_2 e$$

$$\Rightarrow \frac{dy}{dx} = \log_2 e \cdot \frac{d}{dx} [\log_e \{\log_e x \cdot \log_2 e\}]$$

$$\Rightarrow \frac{dy}{dx} = \log_2 e \cdot \frac{1}{\log_e x \cdot \log_2 e} \cdot \frac{d}{dx} (\log_e x \cdot \log_2 e)$$

66. (c) The function breaks at $x = 0$ and multiples of π . Hence the function is differentiable at all other points as the function is continuous at all these pts.



At $x = 0$, for $f(x)$ to be continuous

$$\lim_{x \rightarrow 0} f(0^-) = f(x = 0) = \lim_{x \rightarrow 0} f(0^+)$$

$$f(x) = 0 \text{ at } x = 0$$

$$\text{RHL} = \lim_{x \rightarrow 0} \sin(x + h) = \sin h > 0$$

$$\text{L. H. L.} = \lim_{x \rightarrow 0} \sin(x - h) = \sin(-h) < 0$$

Hence, not differentiable at $x = 0$

Similarly, $f(x)$ is not differentiable at all multiples of π , i.e., $n\pi$ where $n = 0, 1, 2, \dots$

67. (d)
$$\int_0^{\pi/3} \frac{\cos x + \sin x}{\sqrt{1 + \sin 2x}} dx$$

$$= \int_0^{\pi/3} \frac{\cos x + \sin x}{\sqrt{\sin^2 x + \cos^2 x + 2 \sin x \cos x}} dx$$

$$= \int_0^{\pi/3} \frac{\cos x + \sin x}{\sqrt{(\cos x + \sin x)^2}} dx = \int_0^{\pi/3} dx = \frac{\pi}{3}$$

68. (c) The equation of the pair of tangents is given by $SS_1 = T^2$

$$(3x^2 + 2y^2 - 5)(3.1^2 + 2.2^2 - 5) = (3x \cdot 1 + 2y \cdot 2 - 5)^2$$

$$9x^2 - 4y^2 - 24xy + 40y + 30x - 55 = 0$$

further angle, θ between them can be found by using

$$\tan \theta = \frac{2\sqrt{h^2 - ab}}{a + b} = \frac{2\sqrt{(12)^2 - (9)(-4)}}{9 + (-4)}$$

69. (d) The roots of the equation $x^2 + x + 1$ are given as ω & ω^2 . i.e. say, $\alpha = \omega$ & $\beta = \omega^2$
 $\alpha^{19} = \omega^{19} = (\omega^3)^6\omega = \omega$; $\beta^7 = (\omega^2)^7 = \omega^{14} = (\omega^3)^4\omega^2 = \omega^2$
Hence the equation is $x^2 + x + 1 = 0$

70. (c) $\sin^{-1}(1-x) = \left(\frac{\pi}{2} - \sin^{-1}x\right) - \sin^{-1}x \quad (\because \cos^{-1}x = \frac{\pi}{2} - \sin^{-1}x)$

$$\sin^{-1}(1-x) = \frac{\pi}{2} - 2\sin^{-1}x$$


Taking sum of both sides

$$1-x = \sin\left(\frac{\pi}{2} - 2\sin^{-1}x\right) = \cos(2\sin^{-1}x)$$

$$= \cos 2\theta, \text{ where } \sin^{-1}x = \theta$$

$$1-x = 1-2\sin^2\theta = 1-2x^2 \text{ or } x(1-2x)=0 \text{ or } x=0, \frac{1}{2}$$

71. (0.33)



Solving, $y = x^2$ and $x = y^2$


$$y = y^4 \text{ or } y(y^3 - 1) = 0 \Rightarrow y = 0 \text{ or } y = 1$$

\therefore Point of intersection are $(0, 0)$ & $(1, 1)$

$$\text{To find the shaded area, } A = \int_0^1 (\sqrt{x} - x^2) dx$$

$$= \frac{2}{3} \left[x^{3/2} \right]_0^1 - \left[\frac{x^3}{3} \right]_0^1 = \frac{2}{3} - \frac{1}{3} = \frac{1}{3}$$

72. (0.63)



$$\text{Req. prob.} = P(A) + P(B) + P(C) - P(A \cap B)$$

$$- P(B \cap C) - P(C \cap A) + P(A \cap B \cap C)$$

$$= \frac{1}{4} + \frac{1}{4} + \frac{1}{4} - 0 - 0 - \frac{1}{8} + 0 = \frac{5}{8}$$

73. (1) $T_{r+1} = {}^6C_r x^{6-r} \left(\frac{1}{x^2}\right)^r = {}^6C_r (x)^{6-r-2r}$

For coefficient of x^6 , $6-r-2r=6$, or $r=0$

This means the term is the first term.

$$\Rightarrow T_1 = {}^6C_0 x^6 = 1 \cdot x^6$$

$$\Rightarrow \text{coefficient of } x^6 = 1$$

74. (3) For $f(x)$ to be continuous, $\lim_{x \rightarrow 0} f(x) = f(0)$

$$f(0) = k \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{\sin 3x}{\sin x} = \lim_{x \rightarrow 0} \frac{3 \cdot \frac{\sin 3x}{3x}}{\frac{\sin x}{x}} = 3$$

$$\left[\because \lim_{x \rightarrow 0} \frac{\sin \theta}{\theta} = 1 \right]$$

$$\Rightarrow k = 3$$

75. (1.33) Line is \perp to $3x + y = 3$

$$\therefore \text{Slope of line, } m = \frac{1}{3}$$

$$\text{Equation is, } y = mx + c = \frac{x}{3} + c$$

$$\text{It passes through } (2, 2) \Rightarrow 2 = \frac{2}{3} + c$$

$$\Rightarrow c = \frac{4}{3}$$

$$\Rightarrow y - \frac{x}{3} = 4/3 \Rightarrow 3y - x = 4$$

$$\therefore y\text{-intercept} = 4/3$$

MOCK TEST-4

PHYSICS

1. (d) $v = \sqrt{\frac{2gh}{1 + \frac{I}{mr^2}}} = \sqrt{\frac{2 \times 10 \times 3}{1 + \frac{mr^2}{2 \times mr^2}}} = \sqrt{\frac{2 \times 10 \times 3}{\frac{3}{2}}} = \sqrt{40}$


$$\Rightarrow v = r\omega \Rightarrow r = \frac{v}{\omega} = \frac{\sqrt{40}}{2\sqrt{2}} = \sqrt{\frac{40}{8}} = \sqrt{5} \text{ m.}$$

2. (a) By conservation of energy

$$mg(3h) = mg(2h) + \frac{1}{2}mv^2 \quad (\text{v} = \text{velocity at B})$$

$$mgh = \frac{1}{2}mv^2 ; \quad v = \sqrt{2gh}$$

From free body diagram of block at B



$$N + mg = \frac{mv^2}{h} = 2mg ; \quad N = mg$$

3. (b) Bulk modulus, $B = -V_0 \frac{\Delta p}{\Delta V} \Rightarrow \Delta V = -V_0 \frac{\Delta p}{B}$

$$\Rightarrow V = V_0 \left[1 - \frac{\Delta p}{B} \right]$$

$$\therefore \text{Density, } \rho = \rho_0 \left[1 - \frac{\Delta p}{B} \right]^{-1} = \rho_0 \left[1 + \frac{\Delta p}{B} \right]$$

where, $\Delta p = p - p_0 = h\rho_0 g$

= pressure difference between depth and surface of ocean

4. (b) Here, $\vec{E} = 5\hat{i} - 3\hat{j}$ kV/m


$$\begin{aligned}
 V_B - V_A &= - \int_{r_A}^{r_B} \vec{E} \cdot d\vec{r} \\
 &= - \int_{(4, 0, 3)}^{(10, 3, 0)} (5\hat{i} - 3\hat{j}) \cdot (\partial x \hat{i} + \partial y \hat{j} + \partial z \hat{k}) \\
 &= - \int_4^{10} 5 \partial x - \int_0^3 (-3) \partial y + 0 = -5[x]_4^{10} + 3[y]_0^3 \\
 &= -5(10 - 4) + 3(3 - 0) = -30 + 9 = -21 \text{ kV}
 \end{aligned}$$

5. (c) The current upto which bulb of marked 25W -220V, will not fuse

$$I_1 = \frac{W_1}{V_1} = \frac{25}{220} \text{ Amp}$$

$$\text{Similarly, } I_2 = \frac{W_2}{V_2} = \frac{100}{220} \text{ Amp}$$

The current flowing through the circuit



$$I = \frac{440}{R_{eff}}$$

$$R_{eff} = R_1 + R_2$$

$$R_1 = \frac{V_1^2}{P_1} = \frac{(220)^2}{25}; \quad R_2 = \frac{V_2^2}{P} = \frac{(220)^2}{100}$$

$$I = \frac{440}{\frac{(220)^2}{25} + \frac{(220)^2}{100}} = \frac{440}{(220)^2 \left[\frac{1}{25} + \frac{1}{100} \right]}$$

$$I = \frac{40}{220} \text{ Amp}$$


(Ans) (a) (b) (c)

6. (c) When the current flows in both wires in the same direction then magnetic field at half way due to the wire P,

$$\vec{B}_P = \frac{\mu_0 I_1}{2\pi \frac{5}{2}} = \frac{\mu_0 I_1}{\pi \cdot 5} = \frac{\mu_0}{2\pi}$$

(where $I_1 = 5$ amp)

The direction of \vec{B}_P is downward \odot



Magnetic field at half way due to wire Q

$$\vec{B}_Q = \frac{\mu_0 I_2}{2\pi \frac{5}{2}} = \frac{\mu_0}{\pi} \quad [\text{upward } \odot]$$

[where $I_2 = 5$ amp.]

Net magnetic field at half way

$$\vec{B} = \vec{B}_P + \vec{B}_Q = -\frac{\mu_0}{2\pi} + \frac{\mu_0}{\pi} = \frac{\mu_0}{2\pi} \quad (\text{upward})$$

Hence, net magnetic field at midpoint = $\frac{\mu_0}{2\pi}$

7. (b)

8. (a) Limiting friction between block and slab

$$= \mu_s m_A g = 0.6 \times 10 \times 9.8 = 58.8 \text{ N}$$

But applied force on block A is 100 N. So the block will slip over a slab.

Now kinetic friction works between block and slab

$$F_k = \mu_k m_A g = 0.4 \times 10 \times 9.8 = 39.2 \text{ N}$$

This kinetic friction helps to move the slab

9. (a) $V_p = \frac{dx}{dt} = a + 2bt$

and $V_Q = \frac{dx}{dt} = f - 2t$, Given, $V_p = V_Q$

$$\therefore a + 2bt = f - 2t \Rightarrow t = \frac{f - a}{2(b + I)}$$

10. (c) Intensity of light

$$I = \frac{\text{Watt}}{\text{Area}} = \frac{nhc}{A\lambda} \Rightarrow \text{Number of photon} = \frac{IA\lambda}{hc}$$

$$\therefore \text{Number of photoelectrons emitted} = \frac{1}{100} \times \frac{IA\lambda}{hc}$$

$$= \frac{1}{100} \times \frac{1 \times 10^{-4} \times 300 \times 10^{-9}}{6.6 \times 10^{-34} \times 3 \times 10^8} = 1.5 \times 10^{12} \text{ per sec}$$

11. (c) The wavelength of spectral line in Balmer series is given

$$\text{by } \frac{1}{\lambda} = R \left[\frac{1}{2^2} - \frac{1}{n^2} \right]$$

For first line of Balmer series, $n = 3$

$$\Rightarrow \frac{1}{\lambda_1} = R \left[\frac{1}{2^2} - \frac{1}{3^2} \right] = \frac{5R}{36}; \text{ For second line } n = 4.$$

$$\Rightarrow \frac{1}{\lambda_2} = R \left[\frac{1}{2^2} - \frac{1}{4^2} \right] = \frac{3R}{16}$$

$$\therefore \frac{\lambda_2}{\lambda_1} = \frac{20}{27} \Rightarrow \lambda_2 = \frac{20}{27} \times 6561 = 4860 \text{ Å}$$

12. (d) In pure semiconductor electron-hole pair = $7 \times 10^{15}/\text{m}^3$

$n_{\text{initial}} = n_h + n_e = 14 \times 10^{15}$ after doping donor Impurity

$$N_D = \frac{5 \times 10^{28}}{10^7} = 5 \times 10^{21} \text{ and } n_e = \frac{N_D}{2} = 2.5 \times 10^{21}$$

So, $n_{\text{final}} = n_h + n_e$

$$\Rightarrow n_{\text{final}} \approx n_e \approx 2.5 \times 10^{21} (\because n_e \gg n_h)$$

$$\text{Factor} = \frac{n_{\text{final}} - n_{\text{initial}}}{n_{\text{initial}}}$$

13. (a) According to Wien's displacement law

$$\lambda_m \propto \frac{1}{T} \Rightarrow \lambda m_2 < \lambda_{m_1} [\because T_1 < T_2]$$

Therefore I- λ graph for T_2 has lesser wavelength (λ_m) and so curve for T_2 will shift towards left side.

14. (b) At resonance, amplitude of oscillation is maximum

$$\Rightarrow 2\omega^2 - 36\omega + 9 \text{ is minimum}$$

$$\Rightarrow 4\omega - 36 = 0 \text{ (derivative is zero)}$$

$$\Rightarrow \omega = 9$$

15. (b) Average speed of gas molecules is $\sqrt{\frac{8kT}{\pi m}}$. It depends on temperature and molecules mass. So the average speed of O_2 will be same in (A) and (C).

16. (c) Apparent frequency

$$n' = n \frac{(u + v_w)}{(u + v_w - v_s \cos 60^\circ)} = \frac{510 (330 + 20)}{330 + 20 - 20 \cos 60^\circ}$$

$$= 510 \times \frac{350}{340} = 525 \text{ Hz}$$

17. (c) The area swept by radius OC in one half circle is $\pi r^2/2$. The flux change in time $T/2$ is thus $(\pi r^2 B/2)$. The induced emf is then $e = \pi r^2 B/T = B\omega r^2/2$

$$\left[\because T = \frac{2\pi}{\omega} \right]$$

The induced current is then $I = e/R = B\omega r^2/2R$

18. (a) The Instantaneous value of voltage is

$$E = 100 \sin(100t) \text{ V}$$

We get

$$E_0 = 100 \text{ V}, \omega = 100 \text{ rad s}^{-1}$$

The rms value of voltage is

$$E_{\text{rms}} = \frac{E_0}{\sqrt{2}} = \frac{100}{\sqrt{2}} \text{ V} = 70.7 \text{ V}$$

The instantaneous value of current is

$$I = 100 \sin\left(100t + \frac{\pi}{3}\right) \text{ mA}$$

Compare it with

$$I = I_0 \sin(\omega t + \phi)$$

we get

$$I_0 = 100 \text{ mA}, \omega = 100 \text{ rad s}^{-1}$$

The rms value of current is

19. (c) Incident momentum, $p = \frac{E}{c}$


For perfectly reflecting surface with normal incidence

$$\Delta p = 2p = \frac{2E}{c}$$

$$F = \frac{\Delta p}{\Delta t} = \frac{2E}{ct}$$

$$P = \frac{F}{A} = \frac{2E}{ctA}$$

20. (a)



According to New Cartesian sign convention,
Object distance $u = -15 \text{ cm}$

Focal length of a concave lens, $f = -10 \text{ cm}$

Height of the object $h_0 = 2.0 \text{ cm}$

$$\text{According to mirror formula, } \frac{1}{v} + \frac{1}{u} = \frac{1}{f}$$


$$\frac{1}{v} = \frac{1}{f} - \frac{1}{u} = \frac{1}{-10} - \frac{1}{-15} \Rightarrow v = -30 \text{ cm.}$$

This image is formed 30 cm from the mirror on the same side of the object.
It is real image.

$$\text{Magnification of the mirror, } m = \frac{-v}{u} = \frac{h_1}{h_0}$$

$$\Rightarrow \frac{-(-30)}{-15} = \frac{h_1}{2} \Rightarrow h_1 = -4 \text{ cm}$$

21. (1.319)



From the figure

$$R = mg \cos 30^\circ + qE \sin 30^\circ$$

$$\begin{aligned} &= \frac{10\sqrt{3}}{2} + \frac{0.01 \times 100}{2} \\ &= 5\sqrt{3} + 0.5 = 9.16 \text{ N} \end{aligned}$$

$$\text{Frictional force } F_s = \mu R = 0.2 \times 9.16 = 1.832 \text{ N}$$

Resultant force along the plane in the downward direction

$$F = mg \sin 30^\circ - (F_s + qE \cos 30^\circ)$$

$$\begin{aligned} &= 5 - \left(1.832 + 0.01 \times 100 \times \frac{1.732}{2} \right) \\ &= 5 - 2.698 = 2.3 \text{ N} \end{aligned}$$

$$\therefore \text{Acceleration along the plane, } f = \frac{F}{m} = 2.3 \text{ m/sec}^2$$

Distance along the plane = $1 \times \text{cosec } 30^\circ = 2 \text{ m}$

$$s = ut + (1/2) ft^2, u = 0$$

$$\therefore t = \left(\frac{2s}{f} \right)^{1/2} = \left(\frac{2 \times 2}{2.3} \right)^{1/2}$$

$$= 1.319 \text{ sec}$$

22. (3.57×10^7)

Time period of satellite,

$$T = \frac{2\pi(R_E + h)}{\sqrt{GM_E}} = \frac{2\pi(R_E + h)^{3/2}}{\sqrt{GM_E}}$$

$$T^2 = \frac{4\pi^2(R_E + h)^3}{GM_E}$$

$$(R_E + h)^3 = \frac{GM_E T^2}{4\pi^2}$$

$$(R_E + h) = \left(\frac{GM_E T^2}{4\pi^2} \right)^{1/3}$$

$$\text{or } h = \left(\frac{GM_E T^2}{4\pi^2} \right)^{1/3} - R_E$$

Here, $M_E = 6 \times 10^{24} \text{ kg}$

$$R_E = 6400 \text{ km} = 6400 \times 10^3 \text{ m} = 6.4 \times 10^6 \text{ m}$$

$$T = 24 \text{ h} = 24 \times 60 \times 60 \text{ s} = 86400 \text{ s}$$

$$G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$$

On substituting the given values, we get

$$h = \left(\frac{6.67 \times 10^{-11} \times 6 \times 10^{24} \times (86400)^2}{4 \times (3.14)^2} \right)^{1/3} - 6.4 \times 10^6 \\ = 4.21 \times 10^7 - 6.4 \times 10^6 = 3.57 \times 10^7 \text{ m}$$

23. (7) Let initial e.m.f. induced = e .

$$\therefore \text{Initial current } i = \frac{E - e}{R} \text{ i.e., } 2 = \frac{12 - e}{1}$$

This gives $e = 12 - 2 = 10 \text{ volt.}$ As $e \propto \omega.$


when speed is halved, the value of induced e.m.f. becomes

$$\frac{e}{2} = \frac{10}{2} = 5 \text{ volt}$$

\therefore New value of current

$$i' = \frac{E - e}{R} = \frac{12 - 5}{1} = 7 \text{ A}$$

24. (40)



$$\begin{aligned}\Delta Q_{ACB} &= \Delta W_{ACB} + \Delta U_{ACB} \\ \Rightarrow 60 \text{ J} &= 30 \text{ J} + \Delta U_{ACB} \\ \Rightarrow \Delta U_{ACB} &= 30 \text{ J} \\ \therefore \Delta U_{ADB} &= \Delta U_{ACB} = 30 \text{ J} \\ \Delta Q_{ADB} &= \Delta U_{ADB} + \Delta W_{ADB} \\ &= 10 \text{ J} + 30 \text{ J} = 40 \text{ J}\end{aligned}$$

25. (1.324)

Energy produced in one day = $10^6 \times 24 \times 60 \times 60$ joule

$$\eta = 0.8 = \frac{\text{output energy}}{\text{input energy}} = \frac{10^6 \times 24 \times 60 \times 60}{\text{input energy}}$$

$$\text{So input energy} = \frac{10^6 \times 24 \times 60 \times 60}{0.8} = 10.8 \times 10^{10} \text{ J}$$

Energy released in one fission

$$= 200 \times 10^6 \times 1.6 \times 10^{-19} = 3.2 \times 10^{-11} \text{ J}$$

$$\text{No. of fissions per day} = \frac{10.8 \times 10^{10}}{3.2 \times 10^{-11}} = 3.375 \times 10^{21}$$

Mass of U²³⁵ consumed per day

$$= \text{no. of nuclei disintegrating per day} \times \text{mass of U}^{235}$$

$$= 3.375 \times 10^{21} \times 235 \times 1.67 \times 10^{-27} = 1.324 \text{ mg}$$

CHEMISTRY

- 26. (b)** When the temperature is increased, energy in form of heat is supplied which increases the kinetic energy of the reacting molecules. This will increase the number of collisions and ultimately the rate of reaction will be enhanced.
- 27. (b)** In lanthanides, there is poorer shielding of $5d$ electrons by $4f$ electrons resulting in greater attraction of the nucleus over $5d$ electrons and contraction of the atomic radii.

- 28. (b)** $(\text{CH}_3)_2\text{CHCH}_2\text{MgBr} \xrightarrow{\text{C}_2\text{H}_5\text{OH}} (\text{CH}_3)_2\text{CHCH}_3 + \text{MgOC}_2\text{H}_5$
- 29. (c)** Using the relation $K_p = K_c \cdot (RT)^{\Delta n}$, we get

Thus $\frac{K_p}{K_c}$ will be highest for the reaction having highest value of Δn .

The Δn values for various reactions are

- (a) $\Delta n = 1 - \left(1 + \frac{1}{2}\right) = -\frac{1}{2}$
- (b) $\Delta n = 2 - (1 + 1) = 0$
- (c) $\Delta n = (1 + 1) - 1 = 1$
- (d) $\Delta n = (2 + 4) - (7 + 2) = -3$

Thus, maximum value of $\Delta n = 1$

- 30.** (d) According to Fajan's rule :


$$\text{Covalent character} \propto \frac{1}{\text{size of cation}} \\ \propto \text{size of anion}$$

Among the given species order of size of cations



Order of size of anions $\text{O}^{2-} > \text{Cl}^-$.

Hence the order of covalent character is



$\therefore \text{BaCl}_2$ is most ionic in nature.

- 31.** (c)

Element	%	Relative no. of atoms	Simplest ratio of atoms
C	49.3	$49.3/12 = 4.1$ $1.5 \times 2 = 3$	$4.1/2.74 = 1.5$
H	6.84	$6.84/1 = 6.84$	$6.84/2.74 = 2.5$ $= 2.5 \times 2 = 5$
O	43.86	$43.86/16 = 2.74$	$2.74/2.74 = 1$ $1 \times 2 = 2$

$\therefore \text{Empirical formula} = \text{C}_3\text{H}_5\text{O}_2$

Empirical formula mass

$$= (3 \times 12) + (5 \times 1) + (2 \times 16) = 36 + 5 + 32 = 73$$

Molecular mass = 2 \times Vapour density

$$= 2 \times 73 = 146$$

$$n = \frac{\text{molecular mass}}{\text{empirical formula mass}} = 146/73 = 2$$

Molecular formula = Empirical formula \times 2

$$= (\text{C}_3\text{H}_5\text{O}_2) \times 2 = \text{C}_6\text{H}_{10}\text{O}_4$$

33. (b) $\Delta G = \Delta H - T \Delta S$

At equilibrium, $\Delta G = 0$

$$\Rightarrow 0 = (170 \times 10^3 \text{ J}) - T(170 \text{ JK}^{-1})$$

$$\Rightarrow T = 1000 \text{ K}$$

For spontaneity, ΔG is – ve, which is possible only if $T > 1000 \text{ K}$.

34. (b) According to gas law

$$PV = nRT, n = \frac{PV}{RT}$$

$$\frac{n_A}{n_B} = \frac{\frac{P_1 V_1}{R T_1}}{\frac{P_2 V_2}{R T_2}} ; \frac{n_A}{n_B} = \frac{P_1 V_1}{T_1} \times \frac{T_2}{P_2 V_2}$$

$$\frac{n_A}{n_B} = \frac{2P \times 2V}{2T} \times \frac{T}{PV} ; \frac{n_A}{n_B} = \frac{2}{1}$$

35. (b) Due to inert pair effect oxidation state decrease by 2 while going down the group in *p*-block.

36. (a) Carbon atom is connected with four different groups in chiral structure.

37. (c) Sr^{90} is harmful radiological pollutant.


38. (d) Here, A_2B_3 can also be written as A_4B_6 .

Since, *hcp* has six atoms, so ‘B’ forms *hcp* lattice and ‘A’ is present in void.



Total tetrahedral voids = 12

\therefore Fraction of tetrahedral voids occupied by


$$A = 4/12 = \frac{1}{3}$$



39. (c) $\text{CH}_3 - \overset{|}{\text{CH}} - \overset{|}{\text{CH}} - \text{CH}_3$. Since it contains only two types of H-atoms hence it will give only two mono chlorinated compounds viz.



40. (c) Let solubility of $\text{PbCl}_2 = s$



$$K_{\text{sp}} = [\text{Pb}^{2+}] [\text{Cl}^-]^2$$

$$\therefore 1.7 \times 10^{-5} = (s)(2s)^2$$

$$\text{or } 1.7 \times 10^{-5} = 4s^3$$

$$\therefore s = \sqrt[3]{\frac{1.7 \times 10^{-5}}{4}} = 1.62 \times 10^{-2}$$

41. (d) d^4 in high spin octahedral complex





$$\text{CFSE} = (-0.4x + 0.6y)\Delta_0$$

Where, $x \rightarrow$ electrons in t_{2g} orbital

$y \rightarrow$ electrons in e_g orbital


$$\text{CFSE} = [0.6 \times 1] + [-0.4 \times 3] = -0.6 \Delta_0$$

42. (d)



43. (b) Nylon is a polyamide polymer

44. (a) Reaction involved:



45. (d) $\lambda = \frac{h}{mv}$


$$\therefore mv = \frac{h}{\lambda} = \frac{6.625 \times 10^{-34}}{0.33 \times 10^{-9}} = 2.01 \times 10^{-24} \text{ kg m sec}^{-1}$$

46. (300) ΔH = Heat of formation at constant pressure

ΔE = Heat of formation at constant volume

$$T = 27^\circ\text{C} = 27 + 273 = 300 \text{ K.}$$

$$R = 2 \text{ cal/degree/mole.}$$





$$\Delta n = n_p - n_r = 1 - \frac{1}{2} = \frac{1}{2}$$

$$\Delta H = \Delta E + \Delta n_g RT \quad \text{or} \quad \Delta H - \Delta E = \Delta n_g RT$$

$$= \frac{1}{2} \times 2 \times 300 = 300 \text{ cal}$$

\therefore Heat of formation of CO at constant pressure and at constant volume at 27°C will differ from one another by 300 cal.



48. (32) $\text{O}_2\% = 20\%$

$$\text{Metal}\% = 80\%.$$



100g of metal oxide contains 80g metal and 20g oxygen

$\therefore \text{Eq. wt. of metal} = \text{mass of metal} \times 8 / \text{mass of oxygen}$

- 49. (4)** The total number of isomers for the complex compound




These four isomers are



The isomer $[\text{Cu}(\text{NH}_3)_2\text{Cl}_2][\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ does not exist due to both parts being neutral.

- 50. (3)** 10 volume solution of H_2O_2 means that 1L of this H_2O_2 solution will give 10 L of oxygen at STP.



2 × 34g 22.7 L at STP

$$= 68\text{g}$$

Thus, 22.4L of O_2 is produced from 68 g H_2O_2 at STP. 10 L of O_2 at STP

$$\text{is produced from } \frac{68 \times 10}{22.4} \text{ g}$$

$$= 29.9\text{gH}_2\text{O}_2 = 30\text{ g}$$

Therefore, strength of H_2O_2 in 10 volume of H_2O_2 solution,

$$= 30\text{ g/L} = 3\% \text{ H}_2\text{O}_2 \text{ solution.}$$

MATHEMATICS

- 51. (a)** Since $(7 + 4\sqrt{3})(7 - 4\sqrt{3}) = 1$,

∴ The given equation becomes

$$y + \frac{1}{y} = 14 \text{ where } y = (7 - 4\sqrt{3})^{x^2 - 4x + 3}$$

$$\Rightarrow y^2 - 14y + 1 = 0 \Rightarrow y = 7 \pm 4\sqrt{3}$$

$$\text{Now } y = 7 + 4\sqrt{3} \Rightarrow x^2 - 4x + 3 = -1 \Rightarrow x = 2, 2$$

$$\text{Also } y = 7 - 4\sqrt{3} \Rightarrow x^2 - 4x + 3 = 1 \Rightarrow x = 2 \pm \sqrt{2}$$

- 52. (a)** $f(x) = x^{3/2} + x^{-3/2} - 4\left(x + \frac{1}{x}\right)$

$$\text{Let } \sqrt{x} + \frac{1}{\sqrt{x}} = t \quad (x > 0)$$

$$\text{Let } g(t) = t^3 - 3t - 4t^2 + 8$$

$$g(t) = t^3 - 4t^2 - 3t + 8$$

$$g'(t) = 3t^2 - 8t - 3 = (t-3)(3t+1)$$

$$g'(t) = 0 \Rightarrow t = 3 \quad (t \neq -1/3)$$

$$g''(t) = 6t - 8$$

$$g''(3) = 10 > 0 \Rightarrow g(3) \text{ is minimum}$$

$$g(3) = 27 - 9 - 36 + 8 = -10$$

53. (a) term of $\left(\frac{x}{2} - \frac{3}{x^2}\right)^{10}$ is ${}^{10}C_t \left(\frac{x}{2}\right)^{10-t} \left(\frac{-3}{x^2}\right)^t$.

Here, $x^{-t+10-2t} = x^4 \Rightarrow -3t + 10 = 4 \Rightarrow t = 2$

Hence coefficient of x^4 is ${}^{10}C_2 \left(\frac{1}{2}\right)^8 (3)^2 = \frac{405}{256}$

54. (b) Given plane $3x + y + 2z + 6 = 0$

and line $\frac{x-1/3}{2b/3} = \frac{y-3}{-1} = \frac{z-1}{a}$

Since plane is parallel to line, then

$$3\left(\frac{2b}{3}\right) + (1)(-1) + 2(a) = 0$$

$$\Rightarrow 2b - 1 + 2a = 0 \Rightarrow a + b = 1/2$$

$$\text{Now, } 3a + 3b = 3/2$$

55. (b) $f(x) = \sqrt{1 + \log_e(1-x)}$ value of $f(x)$ is real when

$$1 + \log_e(1-x) \geq 0 \text{ and } 1-x > 0$$

$$\Rightarrow \log_e(1-x) \geq -1 \text{ and } x < 1$$

$$\Rightarrow \log_e(1-x) \geq \log_e e^{-1} \text{ and } x < 1$$

$$\Rightarrow 1-x \geq \frac{1}{e} \text{ and } x < 1 \Rightarrow x \leq \frac{e-1}{e} \text{ and } x < 1.$$

56. (c) $f(x) = [x]^2 - [x^2]$

Check continuity at $x=0$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} [x]^2 - [x^2] = 0$$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} [x]^2 - [x^2]$$

Thus, discontinuous at $x = 0$

Check continuity at $x = 1$


$$\lim_{x \rightarrow 1^+} f(x) = 1 - 1 = 0$$

$$\lim_{x \rightarrow 1^-} f(x) = 0 - 0 = 0$$

Also $f(1) = 0$

Hence continuous at $x = 1$.

57. (a) $y = 1 + 4x - x^2 = 5 - (x - 2)^2$



We have $\int_0^{3/2} (1 + 4x - x^2) dx = 2 \int_0^{3/2} mx dx$

$$= \frac{3}{2} + 2\left(\frac{9}{4}\right) - \frac{1}{3}\left(\frac{27}{8}\right) = m \cdot \frac{9}{4}$$

On solving we get $m = \frac{13}{6}$

58. (b) Series $3 + 33 + 333 + \dots + n$ terms

Given series can be written as,

$$= \frac{1}{3}[9 + 99 + 999 + \dots + n \text{ terms}]$$

$$= \frac{1}{3}[(10 - 1) + (100 - 1) + (1000 - 1) + \dots + n \text{ terms}]$$

$$= \frac{1}{3}[10 + 10^2 + \dots + 10^n] - \frac{1}{3}[1 + 1 + 1 + \dots + n \text{ terms}]$$

$$= \frac{1}{3} \cdot \frac{10(10^n - 1)}{10 - 1} - \frac{1}{3}n = \frac{1}{3} \left[\frac{10^{n+1} - 10}{9} - n \right]$$

59. (a) Let $I = \int \frac{1}{1+\sin x} dx = \int \frac{dx}{2 \tan \frac{x}{2}}$
 $\quad\quad\quad 1 + \frac{1 + \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}$

$$\int \frac{\left(1 + \tan^2 \frac{x}{2}\right) dx}{1 + \tan^2 \frac{x}{2} + 2 \tan \frac{x}{2}} = \int \frac{\sec^2 \frac{x}{2} dx}{1 + \tan^2 \frac{x}{2} + 2 \tan \frac{x}{2}}$$

Substitute

$$\tan \frac{x}{2} = t \Rightarrow \frac{1}{2} \sec^2 \frac{x}{2} dx = dt \Rightarrow \sec^2 \frac{x}{2} dx = 2dt.$$

Then

$$I = \int \frac{2dt}{1+t^2+2t} = 2 \int \frac{dt}{(1+t)^2} = 2 \frac{-1}{(1+t)} + C$$

$$= \frac{-2}{1+\tan \frac{x}{2}} + C = 1 - \frac{2}{1+\tan \frac{x}{2}} + (C-1) = \frac{\tan \frac{x}{2} - 1}{\tan \frac{x}{2} + 1} + b,$$

Where $b = C - 1$, a new constant

$$= -\frac{1-\tan \frac{x}{2}}{1+\tan \frac{x}{2}} + b = -\tan\left(\frac{\pi}{4} - \frac{x}{2}\right) + b = \tan\left(\frac{x}{2} - \frac{\pi}{4}\right) + b.$$

Clearly $a = -\frac{\pi}{4}$ and $b \in \mathbf{R}$

60. (d) Given expression can be written as

$$\begin{aligned} y &= \tan^{-1} \left[\frac{2^x(2-1)}{1+2^x \cdot 2^{x+1}} \right] = \tan^{-1} \left[\frac{2^{x+1}-2^x}{1+2^x \cdot 2^{x+1}} \right] \\ &= \tan^{-1}(2^{x+1}) - \tan^{-1}(2^x) \end{aligned}$$

$$\Rightarrow \frac{dy}{dx} = \frac{2^{x+1} \log 2}{1+2^{2(x+1)}} - \frac{2^x \log 2}{1+2^{2x}}$$

61. (d) We have, $\cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{6\pi}{7}$

$$\begin{aligned}
 &= \frac{1}{2\sin \frac{\pi}{7}} \left[2\sin \frac{\pi}{7} \cos \frac{2\pi}{7} + 2\sin \frac{\pi}{7} \cos \frac{4\pi}{7} + 2\sin \frac{\pi}{7} \cos \frac{6\pi}{7} \right] \\
 &= \frac{1}{2\sin \frac{\pi}{7}} \left[\left(\sin \frac{3\pi}{7} - \sin \frac{\pi}{7} \right) + \left(\sin \frac{5\pi}{7} - \sin \frac{3\pi}{7} \right) + \left(\sin \frac{7\pi}{7} - \sin \frac{5\pi}{7} \right) \right] \\
 &= -\frac{1}{2} \quad \left[\because \sin \frac{7\pi}{7} = \sin \pi = 0 \right]
 \end{aligned}$$

62. (b) Consider the differential equation

$$\frac{dy}{dx} = y \tan x - y^2 \sec x$$

Divide by y^2 on both the sides, we get

$$\frac{1}{y^2} \left(\frac{dy}{dx} \right) = \frac{\tan x}{y} - \sec x \quad \dots(1)$$

$$\text{Let } \frac{1}{y} = z$$

Differentiating both sides, we get:

$$\frac{-1}{y^2} \cdot \frac{dy}{dx} = \frac{dz}{dx}$$

Put value of $\frac{1}{y^2} \frac{dy}{dx}$ in the equation(1), we get

$$-\left(\frac{dz}{dx} \right) - (\tan x)z = -\sec x$$

$$\Rightarrow \left(\frac{dz}{dx} \right) + (\tan x)z = \sec x$$

This is the linear diff equation in 'z' i.e.

This is of the form $\frac{dz}{z} + P.z = Q$


∴ In the given question

$$\text{I.F.} = e^{\int \tan x dx} = e^{\log(\sec x)} = \sec x$$

63. (c) Here equation of the circle

$$(x^2 + y^2 - 10x) + \lambda(y - 2x) = 0$$

Now centre $C(5 + \lambda, -\lambda/2)$ lies on the chord again.



$$\therefore \frac{-\lambda}{2} = 2(5 + \lambda)$$

$$\therefore \lambda = -4$$

$$\text{Hence } x^2 + y^2 = 10x + 4y - 8x$$

$$\text{or } x^2 + y^2 - 2x - 4y = 0$$

64. (c) Since, \vec{a} and $\vec{b} + \vec{c}$ are mutually perpendicular.

$$\therefore \vec{a} \cdot (\vec{b} + \vec{c}) = 0 \Rightarrow \vec{a} \cdot \vec{b} + \vec{c} \cdot \vec{a} = 0 \quad \dots(i)$$

$$\text{Similarly, } \vec{b} \cdot \vec{c} + \vec{a} \cdot \vec{b} = 0 \quad \dots(ii)$$

$$\text{and } \vec{c} \cdot \vec{a} + \vec{b} \cdot \vec{c} = 0 \quad \dots(iii)$$

On adding eqs. (i), (ii) and (iii), we get

$$2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\text{Now, } |\vec{a} + \vec{b} + \vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a})$$

$$= |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2$$

$$= 9 + 16 + 25 \quad (\because |\vec{a}| = 3, |\vec{b}| = 4, |\vec{c}| = 5)$$

$$65. \text{ (c)} \quad \begin{vmatrix} a-x & c & b \\ c & b-x & a \\ b & a & c-x \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} a+b+c-x & c & b \\ a+b+c-x & b-x & a \\ a+b+c-x & a & c-x \end{vmatrix} = 0$$

$$\Rightarrow (\Sigma a - x) \begin{vmatrix} 1 & c & b \\ 1 & b-x & a \\ 1 & a & c-x \end{vmatrix} = 0$$

$$\Rightarrow x = \Sigma a = 0$$

$$\text{or } 1\{(b-x)(c-x)-a^2\} - c\{c-x-a\} + b\{a-b+x\} = 0$$

(by expanding the determinant.)

$$\text{or } x^2 - (a^2 + b^2 + c^2) + (ab + bc + ca) = 0$$

$$\text{or } x^2 - \Sigma a^2 + \Sigma ab = 0$$

$$\text{or } x^2 - (\Sigma a^2) - \frac{1}{2}(\Sigma a^2) = 0$$

$$[\because a + b + c = 0 \Rightarrow (a + b + c)^2 = 0]$$

$$\Rightarrow \Sigma a^2 + 2\Sigma ab = 0 \Rightarrow \Sigma ab = -\frac{1}{2}\Sigma a^2$$

$$\text{or } x = \pm \sqrt{\frac{3}{2}\Sigma a^2}$$

$$\therefore \text{ the solution is } x = 0 \text{ or } \pm \sqrt{\frac{3}{2}\Sigma a^2}.$$

$$66. \text{ (b)} \quad I_1 = \int_0^1 2^{x^2} dx, \quad I_2 = \int_0^1 2^{x^3} dx, \quad I_3 = \int_1^2 2^{x^2} dx, \quad I_4 = \int_1^2 2^{x^3} dx$$

$$\forall 0 < x < 1, x^2 > x^3$$

$$\Rightarrow \int_0^1 2^{x^2} dx > \int_0^1 2^{x^3} dx \Rightarrow I_1 > I_2.$$

$$\int_2^3 x^2 dx > \int_2^3 x^3 dx$$

67. (c) Given, $f(x) = |x|$ and $g(x) = [x - 3]$

For $-\frac{8}{5} < x < \frac{8}{5}$; $0 \leq f(x) < \frac{8}{5}$

Now, for $0 \leq f(x) < 1$,

$$g(f(x)) = [f(x) - 3] = -3 \quad [\because -3 \leq f(x) - 3 < -2]$$

for $1 \leq f(x) < 1.6$

$$g(f(x)) = -2 \quad [\because -2 \leq f(x) - 3 < -1.4]$$

\therefore required set is $\{-3, -2\}$.

68. (d) We know that $P(A \cup B) \geq \max \{P(A), P(B)\} = \frac{2}{3}$

$$P(A \cap B) \leq \min \{P(A), P(B)\} = \frac{1}{2}$$

$$\text{and } P(A \cap B) = P(A) + P(B) - P(A \cup B) \geq P(A) + P(B) - 1 = \frac{1}{6}$$

$$\Rightarrow \frac{1}{6} \leq P(A \cap B) \leq \frac{1}{2}$$


$$P(A' \cap B) = P(B) - P(A \cap B)$$

$$\therefore \frac{2}{3} - \frac{1}{2} \leq P(A' \cap B) \leq \frac{2}{3} - \frac{1}{6}$$

$$\Rightarrow \frac{1}{6} \leq P(A' \cap B) \leq \frac{1}{2}$$

69. (d) Let $P(a \sec \theta, b \tan \theta)$ and $Q(a \sec \theta, -b \tan \theta)$ be end points of double ordinates and $C(0, 0)$, is the centre of the hyperbola.

Now $PQ = 2b \tan \theta$



$$\therefore OQ = OP = PQ,$$

$$\therefore 4b^2 \tan^2 \theta = a^2 \sec^2 \theta + b^2 \tan^2 \theta$$

$$\Rightarrow 3b^2 \tan^2 \theta = a^2 \sec^2 \theta \Rightarrow 3b^2 \sin^2 \theta = a^2$$

$$\Rightarrow 3a^2(e^2 - 1) \sin^2 \theta = a^2,$$

$$\left[\because e = \sqrt{1 - \frac{b^2}{a^2}} \right]$$

$$\Rightarrow 3(e^2 - 1) \sin^2 \theta = 1$$

$$\Rightarrow \frac{1}{3(e^2 - 1)} = \sin^2 \theta < 1, \quad (\because \sin^2 \theta < 1)$$

$$\Rightarrow \frac{1}{e^2 - 1} < 3 \Rightarrow e^2 - 1 > \frac{1}{3} \Rightarrow e^2 > \frac{4}{3} \Rightarrow e > \frac{2}{\sqrt{3}}.$$

70. (b) Let p be the length of the perpendicular from the origin on the given line.
Then its equation in normal form is

$$x \cos 30^\circ + y \sin 30^\circ = p \Rightarrow \sqrt{3}x + y = 2p$$

This meets the coordinate axes at $A\left(\frac{2p}{\sqrt{3}}, 0\right)$ and $B(0, 2p)$.

$$\therefore \text{Hence, area of } \Delta OAB = \frac{1}{2} \left(\frac{2p}{\sqrt{3}} \right) 2p$$

$$= \frac{2p^2}{\sqrt{3}}$$

\therefore area of triangle is $\frac{50}{\sqrt{3}}$.

$$\therefore \frac{2p^2}{\sqrt{3}} = \frac{50}{\sqrt{3}} \Rightarrow p = \pm 5.$$

Hence the lines are $\sqrt{3}x + y \pm 10 = 0$.

71. (6) The first equation can be written as

$$2 \sin \frac{1}{2}(x+y) \cos \frac{1}{2}(x-y)$$

$$= 2 \sin \frac{1}{2}(x+y) \cos \frac{1}{2}(x+y)$$

\therefore Either $\sin \frac{1}{2}(x+y) = 0$ or $\sin \frac{1}{2}x = 0$ or $\sin \frac{1}{2}y = 0$

or $x - y = -1$ which gives $\left(\frac{1}{2}, -\frac{1}{2}\right)$ or $\left(-\frac{1}{2}, \frac{1}{2}\right)$ as the possible solution.

Again solving with $x = 0$, we get $(0, \pm 1)$ and solving with $y = 0$, we get $(\pm 1, 0)$ as the other solution. Thus we have six pairs of solutions for x and y .

72. (120) Using L-Hospital's rule,

$$\begin{aligned} \lim_{x \rightarrow 0} \left\{ \frac{\sin x - x + \frac{x^3}{6}}{x^5} \right\} &= \lim_{x \rightarrow 0} \frac{\cos x - 1 + \frac{3x^2}{6}}{5x^4} \\ &= \lim_{x \rightarrow 0} \frac{-\sin x + \frac{6x}{6}}{20x^3} = \lim_{x \rightarrow 0} \frac{-\cos x + 1}{60x^2} \\ &= \lim_{x \rightarrow 0} \frac{\sin x}{120x} = \lim_{x \rightarrow 0} \frac{\cos x}{120} = \frac{1}{120} \end{aligned}$$

73. (750) Let edge of the cube be x cm.

Volume of the cube be x^3 cm 3 .

$$\text{Given, } \frac{dx}{dt} = 10 \text{ cm/sec}$$

$$\text{Now, } v = x^3 \Rightarrow \frac{dv}{dt} = 3x^2 \frac{dx}{dt}$$

$$\Rightarrow \frac{dv}{dt} = 3(5)^2 (10) \text{ cm}^3/\text{sec} = 750 \text{ cm}^3/\text{sec}.$$

$$\text{74. (0)} \quad \text{Given } 2x = -1 + \sqrt{3}i \quad \Rightarrow \quad x = \frac{-1 + \sqrt{3}i}{2} = \omega$$

$$\begin{aligned} &\text{Now } (1 - \omega^2 + \omega)^6 - (1 - \omega + \omega^2) \\ &= (-\omega^2 - \omega^2)^6 - (-\omega - \omega)^6 \quad (\because 1 + \omega + \omega^2 = 0) \\ &= (-2\omega^2)^6 - (-2\omega)^6 = (-2)^6(\omega^3)^4 - (-2)^6(\omega^3)^2 \\ &= (-2)^6 - (-2)^6 = 0 \quad (\because \omega^3 = 1) \end{aligned}$$

75. (13986)

The non-zero perfect square digits are 1, 4 and 9.


1 can occur at units place in $3 \times 3 = 9$ ways.

\therefore Sum due to 1 at units place is 1×9 . Similarly,
sum due to 1 at tens place is $1 \times 10 \times 9$ and

sum due to 1 at hundreds place $1 \times 100 \times 9$. We can deal with the digits

MOCK TEST-5

PHYSICS



Apply the law of conservation of linear momentum along a direction perpendicular to the direction of motion (i.e. along *y*-axis), we get

$$0 + 0 = 4v_1 \sin 30^\circ - v_2 \sin 60^\circ$$

$$4v_1 \sin 30^\circ = v_2 \sin 60^\circ$$

$$\frac{v_1}{v_2} = \frac{\sin 60^\circ}{4 \sin 30^\circ} = \frac{\sqrt{3}}{4}$$

2. (d) Mass per unit length of the wire = ρ

$$\text{Mass of L length, } M = \rho L$$

and since the wire of length L is bent in a form of circular loop therefore

$$2\pi R = L \Rightarrow R = \frac{L}{2\pi}$$

$$\text{Moment of inertia of loop about given axis} = \frac{3}{2} M R^2$$


$$= \frac{3}{2} \rho L \left(\frac{L}{2\pi} \right)^2 = \frac{3\rho L^3}{8\pi^2}$$

3. (c) $V_{\text{in}} = \frac{-GM}{2R} \left[3 - \left(\frac{r}{R} \right)^2 \right],$

$$V_{\text{surface}} = \frac{-GM}{R}, V_{\text{out}} = \frac{-GM}{r}$$

4. (c) Electric field, $E \propto \frac{1}{x}$

5. (a) Let each plate moves a distance x from its initial position.
Let q charge flows in the loop. Using Kirchoff's voltage law



$$\frac{\left(\frac{Q}{2}-q\right)(d_0+x)}{\epsilon_0 A} - \frac{\left(\frac{Q}{2}+q\right)(d_0-x)}{\epsilon_0 A} = 0$$

$$\therefore q = \frac{Qx}{2d_0}; I = \frac{dq}{dt} = \frac{Q}{2d_0} \left(\frac{dx}{dt} \right) = \frac{Q}{2d_0} u_0$$

6. (a) The magnetic field varies inversely with the distance for a long conductor.
That is, $B \propto \frac{1}{d}$
so, graph (a) is the correct one.

7. (d) Applying dimensional method :

$$v_c = \eta^x \rho^y r^z$$

$$[M^0 L T^{-1}] = [M L^{-1} T^{-1}]^x [M L^{-3} T^0]^y [M^0 L T^0]^z$$

Equating powers both sides

$$x + y = 0; -x = -1 \therefore x = 1$$

$$1 + y = 0 \therefore y = -1$$


$$-x - 3y + z = 1$$

$$-1 - 3(-1) + z = 1$$

$$-1 + 3 + z = 1$$

$$\therefore z = -1$$

8. (d)



In fig., $A A_1 = v_{\max} = \alpha t_1 = \beta t_2$

But $t = t_1 + t_2 = \frac{v_{\max}}{\alpha} + \frac{v_{\max}}{\beta}$

$$= v_{\max} \left(\frac{1}{\alpha} + \frac{1}{\beta} \right) = v_{\max} \left(\frac{\alpha + \beta}{\alpha \beta} \right)$$

or, $v_{\max} = t \left(\frac{\alpha \beta}{\alpha + \beta} \right)$

9. (c) Given, $u \cos \theta = \frac{\sqrt{3} u}{2}$

$$\Rightarrow \cos \theta = \frac{\sqrt{3}}{2} \Rightarrow \theta = 30^\circ$$

$$\text{Range (R)} = \frac{u^2 \sin 2\theta}{g} = \frac{u^2 \sin 60^\circ}{g} = \frac{\sqrt{3} u^2}{2g}$$

$$\text{Maximum height} = \frac{u^2 \sin^2 \theta}{2g} = \frac{u^2 \sin^2 30^\circ}{2g} = \frac{u^2}{8g}$$

Now, Range = P × H

$$\Rightarrow \frac{\sqrt{3} u^2}{2g} = P \times \frac{u^2}{8g} \Rightarrow P = 4\sqrt{3}$$

10. (d) The electron ejected with maximum speed v_{\max} are stopped by electric field $E = 4N/C$ after travelling a distance $d = 1m$

$$\frac{1}{2}mv_{\max}^2 = eEd = 4eV$$

$$\text{The energy of incident photon} = \frac{1240}{200} = 6.2 \text{ eV}$$

From equation of photo electric effect

$$\frac{1}{2}mv_{\max}^2 = h\nu - \phi_0$$

$$\therefore \phi_0 = 6.2 - 4 = 2.2 \text{ eV}$$


11. (d) Shortest wavelength comes from $n_1 = \infty$ to $n_2 = 1$ and longest wavelength comes from $n_1 = 6$ to $n_2 = 5$ in the given case.

$$\frac{1}{\lambda_{\max}} = R \left(\frac{1}{5^2} - \frac{1}{6^2} \right) = R \left(\frac{36-25}{25 \times 36} \right) = \frac{11}{900} R$$

$$\therefore \frac{\lambda_{\max}}{\lambda_{\min}} = \frac{900}{11}$$

12. (c) The range of energy of β -particles is from zero to some maximum value.
 13. (c) According to Newton's law of cooling, the temperature goes on decreasing with time non-linearly.

14. (c) The equation for the line is



$$P = \frac{-P_0}{V_0} V + 3P_0 \quad [\text{slope} = \frac{-P_0}{V_0}, c = 3P_0]$$

$$\begin{aligned} PV_0 + P_0 V &= 3P_0 V_0 \\ \text{But } PV &= nRT \end{aligned} \quad \dots(i)$$

$$\therefore P = \frac{nRT}{V} \quad \dots(ii)$$

From (i) and (ii)

$$\frac{nRT}{V} V_0 + P_0 V = 3P_0 V_0$$

$$\therefore nRT V_0 + P_0 V^2 = 3P_0 V_0 \quad \dots(iii)$$

$$\text{For temperature to be maximum } \frac{dT}{dV} = 0$$

Differentiating e.q. (iii) by 'V' we get


$$nRV \frac{dT}{dV} + P_0 (2V) = 3P_0 V_0$$

$$\frac{dT}{dV} = \frac{3P_0 V_0 - 2P_0 V}{n R V_0} = 0$$

$$V = \frac{3V_0}{2} \quad \therefore P = \frac{3P_0}{2} \quad [\text{From (i)}]$$

$$\therefore T_{\max} = \frac{9P_0 V_0}{4nR} \quad [\text{From (iii)}]$$

15. (b) $P_1 > P_2$



As $V = \text{constant} \Rightarrow P \propto T$

Hence from $V-T$ graph $P_1 > P_2$

16. (d) At $t = 2$ sec, the particle crosses mean position.

At $t = 4$ sec, its velocity is 4 ms^{-1}

For simple harmonic motion, $y = a \sin \omega t$

$$\therefore y = a \sin\left(\frac{2\pi}{T}t\right)$$

$$y_1 = a \sin\left[\left(\frac{2\pi}{16}\right) \times 2\right] = a \sin\left(\frac{\pi}{4}\right) = \frac{a}{\sqrt{2}} \quad \dots(i)$$


After 4 sec or after 2 sec from mean position, $y_1 = \frac{a}{\sqrt{2}}$,

velocity = 4 ms^{-1}

$$\therefore \text{Velocity} = \omega \sqrt{a^2 - y_1^2}$$

$$\Rightarrow 4 = \left(\frac{2\pi}{16}\right) \sqrt{a^2 - \frac{a^2}{2}} \quad [\text{from (i)}]$$

17. (d) Here, induced e.m.f.



$$e = \int_{2\ell}^{3\ell} (\omega x) B dx = B\omega \frac{[(3\ell)^2 - (2\ell)^2]}{2}$$

$$= \frac{5B\ell^2\omega}{2}$$

18. (c) Charge on the capacitor at any time t is given by

$$q = CV(1 - e^{-t/\tau})$$

at $t = 2\tau$

$$q = CV(1 - e^{-2})$$

19. (b) \because The E.M. wave are transverse in nature i.e.,

$$= \frac{\vec{k} \times \vec{E}}{\mu\omega} = \vec{H} \quad \dots(i)$$

$$\text{where } \vec{H} = \frac{\vec{B}}{\mu}$$

$$\text{and } \frac{\vec{k} \times \vec{H}}{\omega\epsilon} = -\vec{E} \quad \dots(ii)$$

\vec{k} is \perp \vec{H} and \vec{k} is also \perp to \vec{E}

or In other words $\vec{X} \parallel \vec{E}$ and $\vec{k} \parallel \vec{E} \times \vec{B}$

20. (b) Acceleration of block AB = $\frac{3mg}{3m+m} = \frac{3}{4}g$

$$\text{Acceleration of block CD} = \frac{2mg}{2m+m} = \frac{2g}{3}$$

Acceleration of image in mirror AB = $2 \times$ acceleration of mirror =

$$2 \left(\frac{-3g}{4} \right) = \frac{-3}{2}g$$


$$\text{Acceleration of image in mirror CD} = 2 \left(\frac{2g}{3} \right) = \frac{4g}{3}$$

\therefore Acceleration of the two images w.r.t. each other

21. (2.5) If C_e be the effective capacitance, then

$$V_C = \frac{1}{2} V_0$$

$$\frac{q}{C_e} = \frac{q_0}{2C_e}$$



$$\Rightarrow q_0(1 - e^{-t/RC_e}) = \frac{q_0}{2} \Rightarrow t = RC_e \ln 2$$

For parallel grouping

$$C_e = \frac{2C}{2}$$

$$\therefore t_2 = 2RC \ln 2$$

For series grouping,

$$C_e = \frac{C}{2}$$

$$\therefore t_1 = \frac{RC}{2} \ln 2$$

$$\therefore \frac{t_2}{t_1} = \frac{1}{4} \Rightarrow t_2 = 2.5s$$

22. (22) $v_e = 11 \text{ Km/s.}$

$$R_p = 2R$$

$$\rho' = \rho$$

$$\therefore g = \frac{GM}{R^2} = \frac{G \cdot \frac{4}{3}\pi R^3 \rho}{R^2} = 4\pi G R \rho$$


$$g_p = \frac{G \frac{4}{3}\pi R_p^3 \rho}{R_p^2} = 4\pi G R_p \rho$$

$$= 2 \times (4\pi G R \rho) = 2 \times g$$

$$v_e' = \sqrt{2g_p R_p} = \sqrt{2 \times 2g \times 2R} = 2\sqrt{2gR}$$

$$= 2 \times 11 = 22 \text{ km/s.}$$

23. (70)



At glass rod S_2

$1 \rightarrow n$ refraction

$$\frac{n}{\infty} - \frac{1}{u_2} = \frac{n-1}{+10}$$

$$\Rightarrow u_2 = -20 \text{ cm}$$

At glass rod S_1

For $n \rightarrow 1$ refraction

$$v_1 = d - 20$$

$$\frac{1}{d-20} - \frac{n}{(-50)} = \frac{1-n}{-10}$$

$$\frac{1}{d-20} - \frac{n}{(-50)} = \frac{1-n}{-10}$$

$$\frac{1}{d-20} + \frac{n}{50} = +\frac{1}{20}$$

$$d = 70 \text{ cm}$$

24. (0.3) The displacement of particle, executing SHM

$$y = 5 \sin\left(4t + \frac{\pi}{3}\right) \quad \dots(i)$$

$$\text{Velocity of particle, } \frac{dy}{dt} = \frac{5d}{dt} \sin\left(4t + \frac{\pi}{3}\right)$$

$$= 5 \cos\left(4t + \frac{\pi}{3}\right) 4 = 20 \cos\left(4t + \frac{\pi}{3}\right)$$

$$\text{Velocity at } t = \left(\frac{T}{4}\right)$$

$$(dy)_{t=\frac{T}{4}} = \left(, \frac{T}{4}, \frac{\pi}{3}\right)$$

$$\Rightarrow u = 20 \cos\left(T + \frac{\pi}{3}\right) \quad \dots(ii)$$

Comparing the given equation with standard equation of SHM $y = a \sin(\omega t + \phi)$, we get $\omega = 4$.

$$\text{As } \omega = \frac{2\pi}{T} \Rightarrow T = \frac{2\pi}{\omega} \Rightarrow T = \frac{2\pi}{4} \Rightarrow T = \left(\frac{\pi}{2}\right)$$

Now, putting value of T in Eq. (ii), we get

$$u = 20 \cos\left(\frac{\pi}{2} + \frac{\pi}{3}\right) = -20 \sin \frac{\pi}{3}$$

$$= -20 \times \frac{\sqrt{3}}{2} = -10 \times \sqrt{3}$$

The kinetic energy of particle,

$$KE = \frac{1}{2} mu^2$$

$$\because m = 2g = 2 \times 10^{-3} kg$$

$$= \frac{1}{2} \times 2 \times 10^{-3} \times (-10\sqrt{3})^2$$

$$= 10^{-3} \times 100 \times 3 = 3 \times 10^{-1} \Rightarrow K.E. = 0.3J$$

25. (0.144) Here, $E = 9V$; $V_z = 6V$; $R_L = 1000\Omega$ and $R_s = 100\Omega$,

Potential drop across series resistor

$$V = E - V_z = 9 - 6 = 3V$$

Current through series resistance R_s is

$$I = \frac{V}{R} = \frac{3}{100} = 0.03 A$$

Current through load resistance R_L is

$$I_L = \frac{V_z}{R_L} = \frac{6}{1000} = 0.006 A$$

Current through Zener diode is

$$I_z = I - I_L = 0.03 - 0.006 = 0.024 \text{ amp.}$$

Power dissipated in Zener diode is

26. (d) (a) $\text{Ba}(\text{N}_3)_2 \xrightarrow{\Delta} \text{Ba} + 3\text{N}_2$
 (b) $(\text{NH}_4)_2\text{Cr}_2\text{O}_7 \xrightarrow{\Delta} \text{Cr}_2\text{O}_3 + \text{N}_2 + 4\text{H}_2\text{O}$
 (c) $\text{NH}_4\text{NO}_2 \xrightarrow{\Delta} \text{N}_2 + 2\text{H}_2\text{O}$
 (d) $(\text{NH}_4)_2\text{SO}_4 \xrightarrow{\Delta} 2\text{NH}_3 + \text{H}_2\text{SO}_4$

NH_3 is evolved in case of (d).

27. (b) Aspirin is analgesic and antipyretic.

28. (b) $\Delta H = E_{a(f)} - E_{a(b)}$

Thus energy of activation for reverse reaction depend upon whether reaction is exothermic or endothermic.

If reaction is exothermic, $\Delta H = -\text{ve}$, $E_{a(b)} > E_{a(f)}$

If reaction is endothermic, $\Delta H = +\text{ve}$ $E_{a(b)} < E_{a(f)}$

29. (c) Liquation process, Mond's process and, van Arkel process are the refining processes that are applied depending upon the nature of the metal under treatment and nature of the impurities whereas amalgamation process is used for the extraction of noble metals like gold, silver, etc, from native ores. The metal is recovered from the amalgam by subjecting it to distillation, where the mercury distils over leaving behind the metal.




30. (b) PbO_2 is a powerful oxidizing agent and liberate O_2 when treated with acids.



31. (d) Soap helps to lower the surface tension of solution, thus soap get stick to the dust particles and grease, and these are removed by action of water.

32. (d) H_3BO_3 acts as a Lewis acid and accepts OH^- ions to form $[\text{B}(\text{OH})_4]^-$



$$\Rightarrow P_{\text{NH}_3}^2 = 3^3 \times 4 K_p$$

$$\Rightarrow P_{\text{NH}_3} = 3^{\frac{3}{2}} \times 2 K_p^{\frac{1}{2}}$$

$$= \frac{3^{\frac{3}{2}} \cdot P^2 K_p^{\frac{1}{2}}}{16}$$


34. (d) We can distinguish between formic acid and acetic acid by their action on Fehling's solution. Formic acid gives a red ppt of cuprous oxide but acetic acid does not give red ppt.

35. (c) $E_{\text{cell}}^\circ = \frac{0.0591}{n} \log K_{\text{eq}}$

$$\therefore 0.591 = \frac{0.0591}{1} \log K_{\text{eq}}$$

$$\text{or } \log K_{\text{eq}} = \frac{0.591}{0.0591} = 10$$

$$\text{or } K_{\text{eq}} = 1 \times 10^{10}$$



37. (c) In a DNA molecule, A = T (Two H-bond)
C ≡ G (Three H-bond)

Purine → Adenine (A), Guanine (G)


Pyrimidine → Cytosine (C), Thymine (T)

So the complimentary sequence of ATGCTTGA is TACGAACT.


38. (c) -CH₃ group is *o*, *p*-directing.

39. (b) Sodium cyanide (Na + C + N → NaCN).
(Lassaigne's test)


40. (b) Magnesium reacts with air to form oxide and nitride. On reaction with water the oxide gives hydroxide and nitride gives hydroxide and ammonia.



(X)



(Y)



(Z)

41. (c) Peptization involves conversion of freshly prepared precipitate into colloidal particles using a suitable electrolyte.

42. (b) $\Delta T_b = K_b \times m \times i = 0.52 \times 1 \times 2 = 1.04$
 $\therefore \Delta T_b = 100 + 1.04 = 101.04^\circ\text{C}$

43. (d) Oxidation state of Cr in $[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]^+$.

Let it be x , $1 \times x + 4 \times 0 + 2 \times (-1) = 1$ Therefore $x = 3$.

44. (a) Higher the value of reduction potential higher will be the oxidising power whereas lower the value of reduction potential higher will be the reducing power.

45. (b) $k = \frac{2.303}{t} \log \frac{a}{(a-x)}$
 $(a-x)$ is the concentration left after 100 sec.

$$2.7 \times 10^{-3} = \frac{2.303}{100} \log \frac{0.29}{(a-x)}$$

$$\Rightarrow \frac{0.27}{2.303} = \log \frac{0.29}{(a-x)} \Rightarrow 0.117 = \log \frac{0.29}{(a-x)}$$

$$\Rightarrow (a-x) = 0.22 \text{ M.}$$

46. (0) It is zero order reaction



47. (38) $\text{M}(\text{NO}_3)_n \rightarrow \text{M}_2(\text{SO}_4)_n$ (n = Valency of metal)

g eq. $\text{M}(\text{NO}_3)_n =$ g eq. of $\text{M}_2(\text{SO}_4)_n$


$$\frac{1.0}{\text{E}(\text{M}) + \text{E}(\text{NO}_3^-)} = \frac{0.86}{\text{E}(\text{M}) + \text{E}(\text{SO}_4^{2-})}$$

$$\Rightarrow \frac{1}{\text{E} + \frac{62}{1}} = \frac{0.86}{\text{E} + \frac{96}{2}} \Rightarrow \text{E} = 38 \text{ g}$$

48. (2)



49. (279) $\wedge^\infty \text{BaCl}_2 = \frac{1}{2} \lambda^\infty \text{Ba}^{2+} + 2\lambda^\infty \text{Cl}^-$
 $= 127 + 2 \times 76 = 279 \text{ S cm}^2 \text{ mol}^{-1}$



The resonance energy provides extra stability to the benzene molecule so it has to be overcome, for hydrogenation to take place.
So $\Delta H = -358.5 - (-150.4) = -208.1 \text{ kJ}$

MATHEMATICS

51. (a) The equation is $x^2 + px + q = 0$

Let α be one of the root, then as per problem, second root is α^2 .

From the principle of quadratic equation.

$$\alpha^2 + \alpha = -p \quad \dots(1)$$

$$\text{and } \alpha^3 = q \quad \dots(2)$$

From eq (1) + eq (2):

$$\alpha^3 + \alpha^2 + \alpha = q - p$$

$$\Rightarrow \alpha(\alpha^2 + \alpha + 1) = q - p$$

$$\Rightarrow \alpha(-p + 1) = q - p \quad [\text{since } \alpha^2 + \alpha = -p \text{ from eqn (1)}]$$

$$\Rightarrow \alpha = \frac{q-p}{1-p} = \frac{p-q}{p-1}$$

Putting this value of α in equation (1)

$$\left(\frac{p-q}{p-1} \right)^2 + \left(\frac{p-q}{p-1} \right) = -p$$

$$\Rightarrow \frac{p^2 - 2pq + q^2}{(p-1)^2} + \frac{p-q}{(p-1)} = -p$$

$$\Rightarrow \frac{p^2 - 2pq + q^2 + (p-1)(p-q)}{(p-1)^2} = -p$$

$$\Rightarrow p^2 - 2pq + q^2 + p^2 - pq - p + q = -p(p^2 - 2p + 1)$$

52. (b) Let $M(h, k)$

$$\text{Given, } AM = 2AB$$

$$\Rightarrow AB + BM = 2AB$$

$$\Rightarrow AB = BM$$

So B is mid point of AM

$$B = \left(\frac{h}{2}, \frac{k+3}{2} \right)$$


\therefore Point B lies on the circle.

$\therefore B$ satisfies the equation of circle. i.e.,

$$\left(\frac{h}{2} \right)^2 + 4 \left(\frac{h}{2} \right) + \left(\frac{k+3}{2} - 3 \right)^2 = 0$$

$$\Rightarrow \frac{h^2}{4} + \frac{8h}{4} + \frac{(k-3)^2}{4} = 0$$

or $x^2 + y^2 + 8x - 6y + 9 = 0$, which is a circle.



53. (b) We have $f(x) = \begin{cases} (x-1)\sin\left(\frac{1}{x-1}\right) & \text{if } x \neq 1 \\ 0 & \text{if } x = 1 \end{cases}$

$$Rf'(1) = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h \sin \frac{1}{h} - 0}{h} = \lim_{h \rightarrow 0} \sin \frac{1}{h}$$

which does not exist.

$\therefore f$ is not differentiable at $x = 1$

$$\text{Also } f'(0) = \left[\sin \frac{1}{(x-1)} - \frac{x-1}{(x-1)^2} \cos \left(\frac{1}{x-1} \right) \right]_{x=0} = -\sin 1 + \cos 1$$

$\therefore f$ is differentiable at $x = 0$

54. (b) $n(A) = 40\% \text{ of } 10,000 = 4,000$

$n(B) = 20\% \text{ of } 10,000 = 2,000$

$n(C) = 10\% \text{ of } 10,000 = 1,000$

$n(A \cap B) = 5\% \text{ of } 10,000 = 500$

$$n(A \cap B \cap C) = 2\% \text{ of } 10,000 = 200$$

$$\begin{aligned} \text{We want to find } n(A \cap B^c \cap C^c) &= n[A \cap (B \cup C)^c] \\ &= n(A) - n[A \cap (B \cup C)] \\ &= n(A) - n[(A \cap B) \cup (A \cap C)] \\ &= n(A) - [n(A \cap B) + n(A \cap C) - n(A \cap B \cap C)] \\ &= 4000 - [500 + 400 - 200] = 4000 - 700 = 3300. \end{aligned}$$

- 55. (d)** Given function $f(x) = Pe^{2x} + Qe^x + Rx$... (i)

$$\text{Given conditions } f(0) = -1, f'(\log 2) = 31$$

$$\text{and } \int_0^{\log 4} [f(x) - Rx] dx = \frac{39}{2}$$

differentiate equation (i)

$$f'(x) = 2Pe^{2x} + Qe^x + R \quad \dots (\text{ii})$$

Put $x = \log 2$ in equation (ii)

$$f'(\log 2) = 2Pe^{2\log 2} + Qe^{\log 2} + R \quad \dots (\text{iii})$$

$$31 = 8P + 2Q + R$$

and, put $x = 0$ in equation (i)

$$f(0) = Pe^{2 \times 0} + Qe^0 + R \cdot 0$$

$$= P + Q - 1 = P + Q$$

$$\Rightarrow P = -1 - Q \quad \dots (\text{iv})$$

$$\text{Thus } \int_0^{\log 4} [f(x) - Rx] dx = \frac{39}{2}$$

$$\Rightarrow \int_0^{\log 4} [Pe^{2x} + Qe^x + Rx - Rx] dx = \frac{39}{2}$$

$$\Rightarrow \int_0^{\log 4} [Pe^{2x} + Qe^x] dx = \frac{39}{2}$$

$$\Rightarrow \left[\frac{Pe^{2x}}{2} + Qe^x \right]_0^{\log 4} = \frac{39}{2}$$

$$\Rightarrow \boxed{P = -1, Q = 0}$$

$$\Rightarrow \frac{15P}{2} + 3Q = \frac{39}{2} \quad \dots (v)$$

From (iv) and (v), we get

$$\frac{15P}{2} + 3(-1 - P) = \frac{39}{2}$$

$$\Rightarrow \frac{9P}{2} = \frac{45}{2} \Rightarrow P = 5$$

$$\text{and } Q = -1 - P = -1 - 5 = -6$$

and from equation (iii)

$$31 = 8 \times 5 + 2 \times -6 + R$$

$$31 = 40 - 12 + R$$

$$\therefore P = 5; Q = -6, R = 3$$

56. (a) $\lim_{x \rightarrow 0^+} x^m (\log x)^n = \lim_{x \rightarrow 0^+} \frac{(\log x)^n}{x^{-m}}, \left(\frac{\infty}{\infty} \text{ Form} \right)$

$$= \lim_{x \rightarrow 0^+} \frac{n(\log x)^{(n-1)} \frac{1}{x}}{-mx^{-m-1}} \quad [\text{Using L-Hospital's rule}]$$

$$= \lim_{x \rightarrow 0^+} \frac{n(\log x)^{(n-1)}}{-mx^{-m}}, \left(\frac{\infty}{\infty} \text{ Form} \right)$$

$$= \lim_{x \rightarrow 0^+} \frac{n(n-1)(\log x)^{(n-2)} \frac{1}{x}}{(-m)^2 x^{-m-1}}$$

[Again using L-Hospital's rule]

$$= \lim_{x \rightarrow 0^+} \frac{n(n-1)(\log x)^{n-2}}{m^2 x^{-m}}, \left(\frac{\infty}{\infty} \text{ Form} \right)$$

.....
.....

$$= \lim_{x \rightarrow 0^+} \frac{n!}{(-m)^n x^{-m}} = 0$$

57. (a) We have ; $f(x) = \sin x - \cos x - ax + b$

$$\Rightarrow f'(x) = \cos x + \sin x - a$$

As the max. value of $(\cos x + \sin x)$ is $\sqrt{2}$

The above is possible when $a \geq \sqrt{2}$

$$\begin{aligned} 58. \quad (d) \quad & \frac{\sin 3B}{\sin B} = \frac{3 \sin B - 4 \sin^3 B}{\sin B} = 3 - 4 \sin^2 B \\ & = 3 - 4 + 4 \cos^2 B = -1 + \frac{4(a^2 + c^2 - b^2)^2}{4(ac)^2} \\ & = -1 + \frac{\left(\frac{a^2 + c^2}{2}\right)^2}{(ac)^2} = -1 + \frac{(a^2 + c^2)^2}{4(ac)^2} \\ & = \frac{(a^2 + c^2)^2 - 4a^2c^2}{4(ac)^2} = \left(\frac{c^2 - a^2}{2ac}\right)^2. \end{aligned}$$

59. (a) We have, $y = (1+x)^y + \sin^{-1}(\sin^2 x)$... (i)

when $x = 0$, we have $y = 1$

Differentiating (i) w.r.t. x we get

$$\frac{dy}{dx} = (1+x)^y \left\{ \frac{dy}{dx} \log(1+x) + \frac{y}{1+x} \right\} + \frac{\sin 2x}{\sqrt{1-\sin^4 x}}$$

$$\Rightarrow \left(\frac{dy}{dx} \right)_{(0,1)} = 1 \Rightarrow - \left(\frac{dx}{dy} \right)_{(0,1)} = -1.$$

So the equation of the normal at $(0, 1)$ is

$$y - 1 = -1(x - 0) \Rightarrow x + y = 1$$

60. (b) Parametric equation of the hyperbola $xy = c^2$ is $(ct, c/t)$
and equation of circle is $x^2 + y^2 = a^2$... (i)
Put $x = ct$ and $y = c/t$ in (i)

$$(ct)^2 + \left(\frac{c}{t}\right)^2 = a^2$$
$$c^2 t^4 + c^2 - a^2 t^2 = 0 \quad \text{(ii)}$$

61. (b) Integration by parts is given as

$$\int_{\text{I II}} u v \, dx = u \int v \, dx - \int \left[\frac{d}{dx}(u) \int v \, dx \right] dx$$

$$\text{Let } I = \int 32x^3 (\log x)^2 \, dx$$

Integrate it by parts, using ILATE so, we choose $(\log x)^2$ as Ist function and x^3 as IInd function

$$= 32 \left\{ (\log x)^2 \frac{x^4}{4} - \int 2 \log x \frac{1}{x} \cdot \frac{x^4}{4} \, dx \right\}$$

$$= \frac{32}{4} x^4 (\log x)^2 - 16 \int x^3 \log x \, dx$$

$$= 8x^4 (\log x)^2 - 16 \left\{ \log x \cdot \frac{x^4}{4} - \int \frac{1}{x} \cdot \frac{x^4}{4} \, dx \right\}$$

$$= 8x^4 (\log x)^2 - 4x^4 \log x + 4 \int x^3 \, dx$$

$$= 8x^4 (\log x)^2 - 4x^4 \log x + x^4 + C$$

$$= x^4 \{8(\log x)^2 - 4 \log x + 1\} + C$$

62. (a) Let $u = \tan^{-1} \frac{2x}{1-x^2}$ (i)

and $v = \sin^{-1} \frac{2x}{1+x^2}$ (ii)

In equation (i) put, $x = \tan \theta$

$$\therefore u = \tan^{-1} \left[\frac{2 \tan \theta}{1 - \tan^2 \theta} \right] = \tan^{-1} (\tan 2 \theta)$$

$$\Rightarrow u = 2 \theta \Rightarrow \frac{du}{d\theta} = 2 \quad \dots \dots \text{(iii)}$$

In equation (ii), put $x = \tan \theta$


$$\therefore v = \sin^{-1} \left[\frac{2 \tan \theta}{1 - \tan^2 \theta} \right] = \sin^{-1} (\sin 2\theta)$$

From equations (iii) and (iv),

$$\frac{du}{dv} = \frac{du}{d\theta} \times \frac{d\theta}{dv} = 2 \times \frac{1}{2} = 1$$

\therefore required differential coefficient will be 1.

63. (c)



$$\text{Parabola: } y^2 = \frac{-x}{2} \text{ and } y^2 = \frac{1}{3}(1-x)$$

On solving, we get $x = -2, y = \pm 1$

$$\therefore \text{required Area} = 2 \left[\frac{1}{\sqrt{3}} \int_{-2}^1 \sqrt{(1-x)} dx - \frac{1}{\sqrt{2}} \int_{-2}^0 \sqrt{-x} dx \right]$$

$$= 2 \left\{ \left[\frac{1}{\sqrt{3}} \times \frac{-2}{3} (1-x)^{3/2} \right]_{-2}^1 - \left[\frac{1}{\sqrt{2}} \times \frac{-2}{3} (-x)^{3/2} \right]_{-2}^0 \right\}$$

$$= 2 \left\{ \left(\frac{2}{3\sqrt{3}} \cdot 3\sqrt{3} \right) - \left(\frac{2}{3\sqrt{2}} \cdot 2\sqrt{2} \right) \right\} = \frac{4}{3}.$$

64. (c) The inverse of the proposition $(p \wedge \sim q) \rightarrow r$ is

$$\sim(p \wedge \sim q) \rightarrow \sim r$$

$$\equiv \sim p \vee \sim(\sim q) \rightarrow \sim r$$

$$\equiv \sim p \vee \sim q \rightarrow \sim r$$

65. (c) The r^{th} term in the expansion of $\left(\frac{3}{2}x^2 - \frac{1}{3x}\right)^9$ is

$$T_{r+1} = {}^9C_r \left(\frac{3}{2}x^2\right)^{9-r} \left(-\frac{1}{3x}\right)^r$$

The coefficient of the term independent of x in the expansion of $(1 + x + 2x^3)$

$$\left(\frac{3}{2}x^2 - \frac{1}{3x}\right)^9 \quad \dots \text{(ii)}$$

= Sum of the coefficient of the terms x^0, x^{-1} and x^{-3} in

$$\left(\frac{3}{2}x^2 - \frac{1}{3x}\right)^9.$$

For x^0 in (i) above, $18 - 3r = 0 \Rightarrow r = 6$.

for x^{-1} in (i) above, there exists no value of r and hence no such term exists.

For x^{-3} in (i), $18 - 3r = -3 \Rightarrow r = 7$

\therefore for term independent of x , in (ii) the coefficient

$$= 1 \times {}^9C_6(-1)^6 \left(\frac{3}{2}\right)^{9-6} \left(\frac{1}{3}\right)^6 + 2 \times {}^9C_7(-1)^7 \left(\frac{3}{2}\right)^{9-7} \left(\frac{1}{3}\right)^7 \\ = \frac{9.8.7}{1.2.3} \cdot \frac{3^3}{2^3} \cdot \frac{1}{3^6} + 2 \frac{9.8}{1.2} (-1) \frac{3^2}{2^2} \cdot \frac{1}{3^7} = \frac{7}{18} - \frac{2}{27} = \frac{17}{54}.$$

66. (a) We have $\frac{dy}{dx} = \frac{f'(x)}{f(x)}y - \frac{y^2}{f(x)} \Rightarrow \frac{dy}{dx} - \frac{f'(x)}{f(x)}y = -\frac{y^2}{f(x)}$

Divide by y^2

$$y^{-2} \frac{dy}{dx} - y^{-1} \frac{f'(x)}{f(x)} = -\frac{1}{f(x)}$$

$$\text{Put } y^{-1} = z \Rightarrow -y^{-2} \frac{dy}{dx} = \frac{dz}{dx}$$

$$-\frac{dz}{dx} - \frac{f'(x)}{f(x)}(z) = -\frac{1}{f(x)} \Rightarrow \frac{dz}{dx} + \frac{f'(x)}{f(x)}z = \frac{1}{f(x)}$$

$$\text{I.F.} = e^{\int \frac{f'(x)}{f(x)} dx} = e^{\log f(x)} = f(x)$$


$$\therefore \text{The solution is } z(f(x)) = \int \frac{1}{f(x)}(f(x)) dx + c$$

67. (d) Given $\angle A - \angle B = \theta \Rightarrow \tan(A - B) = \tan \theta$

$$\Rightarrow \frac{\tan A - \tan B}{1 + \tan A \tan B} = \tan \theta \quad \dots(i)$$

In right angled triangle CDA ,

$$\tan A = \frac{k}{a - h}$$



Similarly in triangle CDB ,

$$\tan B = \frac{k}{a + h}$$

Substitute the values of $\tan A$ and $\tan B$ in (i), we get

$$h^2 - k^2 + 2hk \cot \theta = a^2$$

Hence the locus is $x^2 - y^2 + 2xy \cot \theta = a^2$.

68. (a) Equation of planes passing through intersecting the planes $3x - y - 4z = 0$ and $x + 3y + 6 = 0$ is,

$$(3x - y - 4z) + \lambda(x + 3y + 6) = 0$$

$$(3 + \lambda)x + (3\lambda - 1)y + (-4 + 6\lambda)z = 0 \quad \dots\dots(i)$$

Given, distances of plane (i) from origin is 1.

$$\therefore \frac{6\lambda}{\sqrt{(3 + \lambda)^2 + (3\lambda - 1)^2 + (-4 + 6\lambda)^2}} = 1$$

$$\text{or } 36\lambda^2 = 10\lambda^2 + 26 \text{ or } \lambda = \pm 1$$

Put the value of λ in (i),

$$\therefore (3x - y - 4z) \pm (x + 3y + 6) = 0$$

$$\text{or } 4x + 2y - 4z + 6 = 0 \text{ or } 2x + y - 2z + 3 = 0$$

$$\text{and } 2x - 4y - 4z - 6 = 0 \text{ or } x - 2y - 2z - 3 = 0$$

Thus the required planes are $x - 2y - 2z - 3 = 0$ and $2x + y - 2z + 3 = 0$.

69. (a) Since, angles of Δ are in AP.

Use cosine law in ΔABC

$$\cos 60^\circ = \frac{(10)^2 + (9)^2 - x^2}{2.(10).(9)} \Rightarrow \frac{1}{2} = \frac{181 - x^2}{2.(90)}$$

$$\Rightarrow x^2 = 91 \Rightarrow x = \sqrt{91}.$$

70. (a) $\vec{a} = (1, -1, 2)$, $\vec{b} = (-2, 3, 5)$, $\vec{c} = (2, -2, 4)$

$$\text{So, } \vec{a} = (1, -1, 2) \equiv \hat{i} - \hat{j} + 2\hat{k}; \vec{b}$$

$$= (-2, 3, 5) \equiv -2\hat{i} + 3\hat{j} + 5\hat{k}$$

$$\text{and } \vec{c} = (2, -2, 4) \equiv 2\hat{i} - 2\hat{j} + 4\hat{k}$$

$$\Rightarrow \vec{a} - 2\vec{b} + 3\vec{c} = (\hat{i} - \hat{j} + 2\hat{k}) - 2(-2\hat{i} + 3\hat{j} + 5\hat{k})$$

$$+ 3(2\hat{i} - 2\hat{j} + 4\hat{k})$$

$$= 11\hat{i} - 13\hat{j} + 4\hat{k} \text{ and } (\vec{a} - 2\vec{b} + 3\vec{c}).\hat{i} = 11.$$

71. (120) As the greater side of a triangle has greater angle opposite to it.

\therefore The angle (say C) opposite to $\sqrt{a^2 + b^2 + ab} = c$ (say) is the greatest in this case.

$$\begin{aligned} \text{Now, } \cos C &= \frac{a^2 + b^2 - c^2}{2ab} \\ &= \frac{a^2 + b^2 - (a^2 + b^2 + ab)}{2ab} \quad [\because c^2 = a^2 + b^2 + ab] \\ &= \frac{-ab}{2ab} = \frac{-1}{2}; C = 120^\circ \end{aligned}$$

72. (198) $\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} a & d & g \\ b & e & h \\ c & f & i \end{bmatrix}$

Sum of diagonal elements,

$$a^2 + b^2 + c^2 + d^2 + e^2 + f^2 + g^2 + h^2 + i^2 = 5$$

Case – I: Five (1's) and four (0's)

$${}^9C_5 = 126$$

Case – II: One (2) and one (1)

73. (2) We know that, $|z_1 - z_2| \geq \|z_1\| - \|z_2\|$... (i)

Here $|z_1| = 12$ and $|z_2 - 3 - 4i| = 5$

but $|z_2 - (3 + 4i)| \geq \|z_2\| - \|3 + 4i\|$

$$\Rightarrow 5 \geq \|z_2\| - 5$$

$$\Rightarrow \|z_2\| \leq 10$$

Also from (i) $|z_1 - z_2|$ will have least value when $\|z_2\|$ has greatest value i.e. 10

$$\therefore |z_1 - z_2| \geq 12 - 10 = 2$$

Thus min. value of $|z_1 - z_2|$ is 2.

74. (1875) $x_1 x_2 x_3 x_4 x_5 = 2 \times 3 \times 5^2 \times 7$ we can assign 2, 3 or 7 to any of variable.

We can assign entire 5^2 to just one variable in 5 ways or can assign.

$5^2 = 5 \times 5$ to two variables in 5C_2 ways

$${}^5C_1 + {}^5C_2 = 5 + 10 = 15 \text{ ways}$$

Required number of solutions $= 5 \times 5 \times 5 \times 15 = 1875$

75. (0.55) Total number of cases obtained by taking multiplication of only two numbers out of 100 $= {}^{100}C_2$.

Out of hundred (1, 2, ..., 100) given numbers, there are the numbers 3, 6, 9, 12, ..., 99, which are 33 in number such that when any one of these is multiplied with any one of remaining 67 numbers or any two of these 33 are multiplied, then the resulting products is divisible by 3. Then the number of numbers which are the products of two of the given number are divisible by 3 $= {}^{33}C_1 \times {}^{67}C_1 + {}^{33}C_2$.

Hence the required probability

$$= \frac{{}^{33}C_1 \times {}^{67}C_1 + {}^{33}C_2}{{}^{100}C_2} = \frac{2739}{4950} = 0.55$$