

Quantum Computational Physics

Problem Statements

Numerical Methods in Quantum Mechanics

July 14, 2026

Contents

1	Problem 1: The Truncated Parabola in a Harmonic Oscillator	2
2	Problem 2: The Shifted and Squeezed Gaussian	3
3	Problem 3: Tunneling in the Quartic Double Well	4
4	Problem 4: Numerical Investigation of Quantum Tunneling in a Quartic Double-Well Potential using the Eigenstate Expansion	5
4.1	Stage I: Validation of the Numerical Solver	5
4.2	Stage II: Eigenstate Analysis of the Quartic Double-Well Potential	5
4.3	Stage III: Gaussian Wave Packet Dynamics	6

1 Problem 1: The Truncated Parabola in a Harmonic Oscillator

Problem Statement

Consider a particle of mass m subject to a one-dimensional quantum harmonic oscillator potential:

$$V(x) = \frac{1}{2}m\omega^2 x^2 \quad (1)$$

At time $t = 0$, the particle is prepared in the initial state:

$$\psi(x, 0) = \begin{cases} N(A - Bx^2) & \text{for } x^2 \leq \frac{A}{B} \\ 0 & \text{for } x^2 > \frac{A}{B} \end{cases} \quad (2)$$

where A and B are positive real constants, and N is the normalization constant.

Analytical Tasks

- 1. Normalization Constant:** Determine the normalization constant N in terms of A and B by enforcing the normalization condition:

$$\int_{-\infty}^{\infty} |\psi(x, 0)|^2 dx = 1 \quad (3)$$

- 2. Eigenstate Expansion:** Calculate the probability $P_n = |c_n|^2$ of finding the particle in the energy eigenstate $|\phi_n\rangle$ with energy eigenvalue $E_n = (n + \frac{1}{2})\hbar\omega$. The expansion coefficients are given by the projection:

$$c_n = \langle \phi_n | \psi(x, 0) \rangle = \int_{-\sqrt{A/B}}^{\sqrt{A/B}} \phi_n^*(x) \psi(x, 0) dx \quad (4)$$

where $\phi_n(x)$ are the Hermite-Gaussian eigenfunctions of the quantum harmonic oscillator.

- 3. Time Evolution:** Write down the formal expression for the time-evolved wave function $\psi(x, t)$ for $t > 0$:

$$\psi(x, t) = \sum_{n=0}^{\infty} c_n \phi_n(x) e^{-iE_n t/\hbar} \quad (5)$$

2 Problem 2: The Shifted and Squeezed Gaussian

Problem Statement

A particle of mass m in a harmonic oscillator potential $V(x) = \frac{1}{2}m\omega^2 x^2$ is initially prepared in a displaced (shifted) and squeezed Gaussian wave packet:

$$\psi(x, 0) = N e^{-\frac{(x-x_0)^2}{a^2}} \quad (6)$$

where:

- x_0 is the initial spatial displacement (shift) of the wave packet center.
- a is the initial width of the wave packet, such that $a \neq \sqrt{\frac{\hbar}{m\omega}}$ (meaning the wave packet width differs from the ground-state characteristic length scale $x_c = \sqrt{\frac{\hbar}{m\omega}}$, resulting in a squeezed state).
- N is the normalization constant.

Analytical Tasks

- 1. Time Evolution:** Determine the full time-evolved wave packet $\psi(x, t)$ under the harmonic oscillator Hamiltonian. Show that the wave packet remains Gaussian in form:

$$\psi(x, t) = N(t) e^{-A(t)(x-\langle x \rangle(t))^2 + \frac{i}{\hbar} \langle p \rangle(t)x + i\phi(t)} \quad (7)$$

where $A(t)$ is a complex parameter related to the time-dependent width and phase.

- 2. Expectation Values:** Calculate the expectation values of position $\langle x \rangle(t)$ and momentum $\langle p \rangle(t)$ as functions of time. Verify that they satisfy Ehrenfest's theorem and display classical oscillatory behavior:

$$\langle x \rangle(t) = x_0 \cos(\omega t) \quad (8)$$

$$\langle p \rangle(t) = -m\omega x_0 \sin(\omega t) \quad (9)$$

- 3. Uncertainty Relations and Breathing Mode:** Determine the position uncertainty $\Delta x(t) = \sqrt{\langle x^2 \rangle - \langle x \rangle^2}$ and momentum uncertainty $\Delta p(t)$. Show that the wave packet width oscillates (or "breathes") at a frequency of 2ω .

3 Problem 3: Tunneling in the Quartic Double Well

Problem Statement

A particle of mass m moves in a one-dimensional quartic double-well potential described by:

$$V(x) = -\frac{1}{2}\alpha x^2 + \beta x^4 \quad (10)$$

where α and β are positive constants. This potential features two symmetric minima at $x_{\min} = \pm\sqrt{\frac{\alpha}{4\beta}}$ separated by a potential barrier of height $V_0 = \frac{\alpha^2}{16\beta}$ at $x = 0$.

Physical and Computational Tasks

- 1. Eigenstate Calculation:** Find the lowest energy eigenstates by solving the time-independent Schrödinger equation:

$$\left[-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x) \right] \psi_n(x) = E_n \psi_n(x) \quad (11)$$

numerically on a discretized spatial grid of size N over the domain $x \in [-L, L]$ using finite difference approximations.

- 2. Energy Eigenvalues and Degeneracy Splitting:** Determine the ground state energy E_0 (symmetric state, even parity) and the first excited state energy E_1 (antisymmetric state, odd parity). Calculate the energy splitting:

$$\Delta E = E_1 - E_0 \quad (12)$$

Explain how quantum tunneling lifts the classical degeneracy of the two isolated wells.

- 3. Tunneling Time:** Construct a state initially localized in the right well:

$$\psi_R(x, 0) = \frac{1}{\sqrt{2}} (\psi_0(x) + \psi_1(x)) \quad (13)$$

Show that the wave function evolves in time as:

$$\psi(x, t) = \frac{1}{\sqrt{2}} \left(\psi_0(x) e^{-iE_0 t/\hbar} + \psi_1(x) e^{-iE_1 t/\hbar} \right) \quad (14)$$

Calculate the tunneling time τ required for the particle to tunnel completely from the right well to the left well, where:

$$\tau = \frac{\pi\hbar}{\Delta E} \quad (15)$$

- 4. Grid Convergence Study:** Conduct a numerical grid convergence test by solving the eigenvalue problem for varying spatial discretization grid points

$$N \in \{200, 400, 800, 1600, 3200\}$$

to verify the stability and accuracy of the computed eigenvalues E_0 and energy splitting ΔE .

4 Problem 4: Numerical Investigation of Quantum Tunneling in a Quartic Double-Well Potential using the Eigenstate Expansion

The one-dimensional time-independent Schrödinger equation describes the quantum behavior of a particle of mass m moving under the influence of a potential $V(x)$,

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x), \quad (16)$$

where $\psi(x)$ is the wavefunction and E is the corresponding energy eigenvalue. Analytical solutions to this equation exist only for a limited number of idealized potentials. For more realistic systems, numerical techniques become indispensable.

One important quantum system is the quartic double-well potential,

$$V(x) = -\frac{1}{2}\alpha x^2 + \beta x^4, \quad (17)$$

which exhibits two stable minima separated by a finite potential barrier. Although classically a particle localized in one well cannot cross the barrier if its energy is below the barrier height, quantum mechanics predicts the phenomenon of tunneling, whereby the particle has a finite probability of appearing in the opposite well. This phenomenon gives rise to characteristic energy-level splitting and time-dependent oscillatory dynamics.

The primary objective of this work is to develop and validate a finite-difference numerical solver for the one-dimensional Schrödinger equation and subsequently employ it to investigate the stationary and dynamical properties of a particle confined in a quartic double-well potential.

The study is divided into three stages.

4.1 Stage I: Validation of the Numerical Solver

The numerical solver will first be validated using the harmonic oscillator potential,

$$V(x) = \frac{1}{2}m\omega^2 x^2, \quad (18)$$

whose exact analytical solutions are well established. The validation consists of

1. Computing the first twenty energy eigenvalues.
2. Computing the corresponding twenty eigenfunctions.
3. Constructing an error table by comparing numerical and analytical eigenvalues.
4. Plotting numerical and analytical energy spectra for comparison.
5. Comparing numerical and analytical eigenfunctions graphically.

Successful agreement with the analytical solution verifies the correctness, accuracy, and convergence of the numerical implementation.

4.2 Stage II: Eigenstate Analysis of the Quartic Double-Well Potential

Following validation, the numerical solver will be applied to the quartic double-well potential,

$$V(x) = -\frac{1}{2}\alpha x^2 + \beta x^4. \quad (19)$$

The objectives of this stage are

1. Compute the first twenty eigenvalues.
2. Compute the corresponding twenty eigenfunctions.
3. Perform a grid-convergence study to verify numerical stability.
4. Visualize the potential together with the bound-state wavefunctions.
5. Analyze the energy spectrum, with particular emphasis on tunneling-induced energy splitting.

4.3 Stage III: Gaussian Wave Packet Dynamics

To investigate quantum tunneling dynamically, an initial Gaussian wave packet will be constructed near the left minimum of the double-well potential,

$$\Psi(x, 0) = A \exp \left[-\frac{(x - x_0)^2}{2\sigma^2} \right] e^{ik_0 x}, \quad (20)$$

where A is the normalization constant, x_0 denotes the left potential minimum, σ is the packet width, and k_0 is the initial wave number.

The initial wave packet will be expanded in terms of the stationary eigenstates,

$$\Psi(x, 0) = \sum_{n=0}^{N-1} c_n \phi_n(x), \quad (21)$$

where the expansion coefficients are

$$c_n = \int_{-\infty}^{\infty} \phi_n^*(x) \Psi(x, 0) dx. \quad (22)$$

The subsequent time evolution is obtained from

$$\Psi(x, t) = \sum_{n=0}^{N-1} c_n \phi_n(x) e^{-iE_n t/\hbar}. \quad (23)$$

The dynamical investigation consists of

1. Locating the minimum of the double-well potential.
2. Constructing a Gaussian wave packet localized near the left minimum.
3. Expanding the wave packet in the stationary eigenstate basis.
4. Computing the expansion coefficients c_n .
5. Plotting the probability distribution $|c_n|^2$.
6. Computing the time evolution of $\Psi(x, t)$.
7. Visualizing the probability density $|\Psi(x, t)|^2$ through plots and animations.
8. Analyzing the tunneling dynamics between the two wells and relating the observed behavior to the computed energy spectrum.

The successful completion of these three stages provides a comprehensive numerical framework for solving the one-dimensional Schrödinger equation, validating the computational approach, and investigating quantum tunneling phenomena in the quartic double-well potential. This section focuses on the numerical solution of the Time-Independent Schrödinger Equation (TISE) for different potentials, followed by the analysis of quantum dynamics using the computed eigenstates.