



# Fresnel Reflection & Transmission

TE<sub>z</sub> Transverse

Electric to z

S-polarization or  
Perpendicular pol.

$$\Gamma_{TE} = \frac{|\vec{E}_r|}{|\vec{E}_i|}, \quad \tau_{TE} = \frac{|\vec{E}_t|}{|\vec{E}_i|}$$

$$\vec{E}_i = E_0 e^{i\vec{k}_{1i} \cdot \vec{r}} \hat{y}$$


$$\vec{E}_r = \Gamma_{TE} E_0 e^{i\vec{k}_{1r} \cdot \vec{r}} \hat{y}$$

$$\vec{E}_t = \tau_{TE} E_0 e^{i\vec{k}_{2t} \cdot \vec{r}} \hat{y}$$

$$\vec{k}_{1i} \cdot \vec{r} = k_{1x} x - k_{1z} z = k_1 x \sin \theta_i - k_1 z \cos \theta_i$$

$$\vec{k}_{1r} \cdot \vec{r} = k_{1x} x + k_{1z} z = k_1 x \sin \theta_r + k_1 z \cos \theta_r$$

$$\vec{k}_{2t} \cdot \vec{r} = k_{2x} x - k_{2z} z = k_2 x \sin \theta_t - k_2 z \cos \theta_t$$



$$\vec{H}_i = \frac{E_0}{\eta_1} e^{i \vec{k}_i \cdot \vec{r}} (\cos \theta_i \hat{x} + \sin \theta_i \hat{z})$$

$$\vec{H}_r = \frac{\Gamma_{TE} E_0}{\eta_1} e^{i \vec{k}_r \cdot \vec{r}} (-\cos \theta_r \hat{x} + \sin \theta_r \hat{z})$$

$$\vec{H}_t = \frac{\tau_{TE} E_0}{\eta_2} e^{i \vec{k}_t \cdot \vec{r}} (\cos \theta_t \hat{x} + \sin \theta_t \hat{z})$$

Boundary Conditions

$$E_0 e^{i k_1 (x \sin \theta_i)} + \Gamma_{TE} E_0 e^{i k_1 (x \sin \theta_r)} = \tau_{TE} E_0 e^{i k_2 (x \sin \theta_t)}$$

$$x=0 \Rightarrow \boxed{1 + \Gamma_{TE} = \tau_{TE}}$$

$$\boxed{k_1 \sin \theta_i = k_1 \sin \theta_r = k_2 \sin \theta_t}$$

PHASE MATCHING!

$$k = \frac{\omega}{v} = \frac{\omega}{c} n \quad \rightarrow \text{refractive index}$$


$$k = \omega \sqrt{\mu \epsilon} = \omega \sqrt{\mu_0 \epsilon_0} \sqrt{\epsilon_r} = \frac{\omega}{c} n.$$

$$n = \frac{c}{v} = \frac{\sqrt{\mu_0 \epsilon_0}^{-1}}{\sqrt{\mu_0 \epsilon_0} \sqrt{\epsilon_r}^{-1}} = \sqrt{\epsilon_r}$$

$$\theta_i = \theta_r$$

$$n_1 \sin \theta_i = n_2 \sin \theta_t$$

SNELL'S LAWS OF REFLECTION  
& TRANSMISSION.



Applying BC for  $\vec{H}$  we get

$$\frac{\cos \theta_i}{\eta_1} - \frac{\Gamma_{TE} \cos \theta_r}{\eta_1} = \frac{\tau_{TE} \cos \theta_t}{\eta_2}$$

$$\frac{\cos \theta_i}{\eta_1} (1 - \Gamma_{TE}) = \frac{\cos \theta_t}{\eta_2} (1 + \Gamma_{TE})$$

$$\Gamma_{TE} = \frac{\eta_2 \cos \theta_i - \eta_1 \cos \theta_t}{\eta_2 \cos \theta_i + \eta_1 \cos \theta_t}$$


Fresnel  
Reflection  
&

$$\tau_{TE} = \frac{2 \eta_2 \cos \theta_i}{\eta_2 \cos \theta_i + \eta_1 \cos \theta_t}$$

Transmission  
Coefficients.

$$\eta = \sqrt{\frac{\mu}{\epsilon}} = \sqrt{\frac{\mu_0}{\epsilon_0}} \frac{1}{\sqrt{\epsilon_r}} = \sqrt{\frac{\mu_0}{\epsilon_0}} \cdot \frac{1}{n}$$

TM<sub>z</sub> pol.



$$\Gamma_{TM} = \frac{-\eta_1 \cos \theta_i + \eta_2 \cos \theta_t}{\eta_1 \cos \theta_i + \eta_2 \cos \theta_t}$$


$$\tau_{TM} = 2 \eta_2 \cos \theta_i / (\eta_1 \cos \theta_i + \eta_2 \cos \theta_t)$$

# Brewster Angle


$$\Gamma_{TM} = 0 \Rightarrow \cos \theta_i = \frac{n_2}{n_1} \cos \theta_t$$

$$\sqrt{1 - \sin^2 \theta_i} = \frac{n_2}{n_1} \sqrt{1 - \frac{k_1^2}{k_2^2} \sin^2 \theta_i}$$

$$\theta_i^{Br} = \sin^{-1} \sqrt{\frac{\epsilon_2 (\mu_1 \epsilon_2 - \epsilon_1 \mu_2)}{\mu_1 (\epsilon_2^2 - \epsilon_1^2)}} \stackrel{\mu_1 = \mu_2}{=} \sqrt{\frac{\epsilon_2}{\epsilon_1 + \epsilon_2}}$$




$$\Gamma_{TE} = 0 \Rightarrow \theta_i^{Br} = \sin^{-1} \sqrt{\frac{\mu_2 (\mu_2 \epsilon_1 - \mu_1 \epsilon_2)}{\epsilon_1 (\mu_2^2 - \mu_1^2)}} = \infty \quad (\mu_1 = \mu_2)$$



# Total Internal Reflection

$\mu_1 = \mu_2 \rightarrow$  assume.

$n_1 > n_2 \rightarrow$  WLOG



$$\theta_t = \sin^{-1} \left( \frac{n_1}{n_2} \sin \theta_i \right) = 90^\circ$$

$$\theta_i^c = \theta_c = \sin^{-1} \left( \frac{n_2}{n_1} \right) \rightarrow \text{Critical Angle.}$$

What happens to  $E_t$  beyond  $\theta_c$ ?

$$\cos \theta_t = \sqrt{1 - \sin^2 \theta_t} = \sqrt{1 - \frac{\sin^2 \theta_i}{\sin^2 \theta_c}}$$

Since  $\theta_i > \theta_c \Rightarrow \cos \theta_t$  is now imaginary!

$$\Rightarrow \cos \theta_t = i\alpha = i \sqrt{\frac{\sin^2 \theta_i}{\sin^2 \theta_c} - 1}$$

$$\begin{aligned} \alpha &= \sqrt{\frac{\sin^2 \theta_i}{\sin^2 \theta_c} - 1} = \frac{n_1}{n_2} \sqrt{\sin^2 \theta_i - \frac{n_2^2}{n_1^2}} \\ &= \frac{n_2}{n_1} \sqrt{\sin^2 \theta_i - \frac{n_2^2}{n_1^2}} \end{aligned}$$

$$\begin{aligned}
 E_t &= \tau_{TM} E_0 e^{i \vec{k}_t \cdot \vec{r}} \\
 &= \tau_{TM} E_0 e^{i k_t (x \sin \theta_t + z \cos \theta_t)} \\
 &= \tau_{TM} E_0 e^{i k_t x \sin \theta_t} \underbrace{e^{-\alpha k_t z}}_{\text{evanescent along } z}.
 \end{aligned}$$

$$\begin{aligned}
 \Gamma_{TM} &= \frac{\eta_2 \cos \theta_t - \eta_1 \cos \theta_i}{\eta_2 \cos \theta_t + \eta_1 \cos \theta_i} = \frac{i \overbrace{\alpha \eta_2}^A - \overbrace{\eta_1 \cos \theta_i}^B}{i \alpha \eta_2 + \eta_1 \cos \theta_i} \\
 &= \frac{iA - B}{iA + B} = - \frac{B - iA}{B + iA} = -e^{-i2 \tan^{-1}(\frac{A}{B})} \\
 &= e^{i(\pi - 2 \tan^{-1}(\frac{A}{B}))}
 \end{aligned}$$

$$\Rightarrow |\Gamma_{TM}| = 1 \Rightarrow \text{TIR.}$$

$$\begin{aligned}
 \phi_{TM}(\theta_i) &= \pi - 2 \tan^{-1}\left(\frac{A}{B}\right) \\
 &= 2 \tan^{-1}\left(\frac{B}{A}\right) \quad \text{when } \frac{B}{A} > 0
 \end{aligned}$$





$$\phi_{TM}(\theta_i) = 2 \tan^{-1} \left( \frac{\eta_1 \cos \theta_i}{\alpha \eta_2} \right)$$


$$\Rightarrow \phi_{TM}(\theta_i) = 2 \tan^{-1} \left( \frac{\eta_1^2 \cos \theta_i}{\eta_2^2 \sqrt{\sin^2 \theta_i - \frac{\eta_1^2}{\eta_2^2}}} \right)$$

Goos-Hanchen Phase Shift.

$$\phi_{TE} = 2 \tan^{-1} \left( \frac{\eta_2 \cos \theta_i}{\eta_1 \sqrt{\sin^2 \theta_i - \left( \frac{\eta_1}{\eta_2} \right)^2}} \right)$$



# Three Dielectric Interface



$$E_{1t} = E_{in} t_{12} e^{i\phi} t_{23}$$

$$E_{2t} = E_{in} t_{12} e^{i\phi} r_{23} e^{i\phi} r_{21} e^{i\phi} t_{23}$$

$$E_{(m+1)t} = E_{in} t_{12} t_{23} e^{i\phi} (r_{23} r_{21} e^{2i\phi})^m$$

$$E_t = \sum_{m=0}^{\infty} E_{mt} = E_{in} t_{12} t_{23} e^{i\phi} \sum_{m=0}^{\infty} (r_{12} r_{23} e^{2i\phi})^m$$

$$t = \frac{E_t}{E_{in}} = \frac{t_{12} t_{23} e^{i\phi}}{1 - r_{21} r_{23} e^{2i\phi}}$$

$$E_{or} = r_{12} E_{in}$$

$$E_{1r} = t_{12} r_{23} t_{21} e^{2i\phi} E_{in}$$

$$E_{mr} = t_{12} t_{21} r_{23} e^{2i\phi} E_{in} (r_{21} r_{23} e^{2i\phi})^m$$

$$\Rightarrow r = r_{12} + \frac{t_{12} t_{21} r_{23} e^{2i\phi}}{1 - r_{21} r_{23} e^{2i\phi}}$$

Let  $3=1$ , note  $t_{12} t_{21} = 1 - r_{12}^2$  &  $r_{21} = -r_{12}$

$$r = \frac{r_{12} (1 - e^{2i\phi})}{1 - r_{12}^2 e^{2i\phi}}$$

When  $2i\phi = 2m\pi \Rightarrow \boxed{k_2 d \cos \theta_t = m\pi}$

Fabry Perot Resonator!

$$FSR = \frac{c}{2n_2 d \cos \theta}$$

