EE24BTECH11001 - Aditya Tripathy

Question:

Find the local minimum/maximum of the given function:

$$f(x) = x^2$$

Solution:

We use the method of gradient descent to find the minimum/maximum of the given function, since the objective function is convex. Since the coefficient of $x^2 > 0$, we expect to find a local minimum.

$$x_{n+1} = x_n - \mu f'(x_n) \tag{0.1}$$

$$f'(x_n) = 2x_n \tag{0.2}$$

$$\to x_{n+1} = x_n - 2\mu x_n \tag{0.3}$$

$$= (1 - 2\mu) x_n \tag{0.4}$$

1

Applying unilateral Z-transform,

$$zX(z) - zx_0 = (1 - 2\mu)X(z)$$
(0.5)

$$(z - (1 - 2\mu)) X(z) = zx_0$$
(0.6)

$$X(z) = \frac{zx_0}{z - (1 - 2\mu)} \tag{0.7}$$

$$X(z) = \frac{x_0}{1 - (1 - 2\mu)z^{-1}} \tag{0.8}$$

$$=\sum_{0}^{\infty} (1-2\mu)^{n} z^{-n} \tag{0.9}$$

From the last equation, ROC is

$$|z| > |1 - 2\mu| \tag{0.10}$$

$$\rightarrow |1 - 2\mu| > 0 \tag{0.11}$$

$$\to \mu \in \mathcal{R} \setminus \left\{ \frac{1}{2} \right\} \tag{0.12}$$

Now, if μ satisfies the previous condition,

$$\lim_{n \to \infty} ||x_{n+1} - x_n|| = 0 \tag{0.13}$$

$$\to \lim_{n \to \infty} \|(1 - 2\mu) x_n\| = 0 \tag{0.14}$$

$$\to \|(1 - 2\mu)\| \lim_{n \to \infty} \|x_n\| = 0 \tag{0.15}$$

$$\to \lim_{n \to \infty} ||x_n|| = 0 \tag{0.16}$$

$$\to \lim_{n \to \infty} x_n = 0 \tag{0.17}$$

Taking initial guess = 8

step size = 0.01

tolerance(minimum value of gradient) = 1e-5

We get

 $x_{min} = 4.910792425174398e - 06$

As a fun exercise, the given question can be posed as the following quadratic programming problem:

Find the point lying on the line y = 1, which is nearest to origin.

We can formulate the problem as follows:

$$\min_{\mathbf{x}} \|\mathbf{x} - \mathbf{O}\|^2 = \min_{\mathbf{x}} \|\mathbf{x}\|^2
= \min_{\mathbf{x}} x^2 + y^2$$
(0.18)

$$= \min_{\mathbf{y}} x^2 + y^2 \tag{0.19}$$

$$s.t. \ e_2^{\mathsf{T}} \mathbf{x} = 1 \tag{0.20}$$

Using cvxpy to solve the convex optimization problem, we get Optimal value of x: [-0. 1.] Optimal objective value: 1.0




Fig. 0.1: Minimum value of objective function