

# Trigonometric Functions and Equations

EE24BTECH11001- ADITYA TRIPATHY

## A: FILL IN THE BLANKS

- 1) Suppose  $\sin^3 x \sin 3x = \sum_{m=0}^n C_m \cos x$  is an identity in  $x$ , where  $C_0, C_1, \dots, C_n$  are constants and  $C_n \neq 0$  then the value of  $n$  is (1981 – 2Marks)
- 2) The solution set of the system of equations  $x + y = \frac{2\pi}{3}$ ,  $\cos x + \cos y = \frac{3}{2}$ , where  $x$  and  $y$  are real, is (1987 – 2Marks)
- 3) The set of all  $x$  in the interval  $[0, \pi]$  for which  $2 \sin^2 x - 3 \sin x + 1 \geq 0$ , is (1987 – 2Marks)
- 4) The sides of a triangle in a given circle subtend angles  $\alpha, \beta, \gamma$ . The minimum value of arithmetic mean of  $\cos\left(\alpha + \frac{\pi}{2}\right)$ ,  $\cos\left(\beta + \frac{\pi}{2}\right)$ ,  $\cos\left(\gamma + \frac{\pi}{2}\right)$  is equal to (1987 – 2Marks)
- 5) The value of  $\sin \frac{\pi}{14} \sin \frac{3\pi}{14} \sin \frac{5\pi}{14} \sin \frac{7\pi}{14} \sin \frac{9\pi}{14} \sin \frac{11\pi}{14} \sin \frac{13\pi}{14}$  is equal to (1991 – 2Marks)
- 6) If  $K = \sin\left(\frac{\pi}{18}\right) \sin\left(\frac{5\pi}{18}\right) \sin\left(\frac{7\pi}{18}\right)$  then the numerical value of  $K$  is (1993 – 2Marks)
- 7) If  $A > 0, B > 0$  and  $A + B = \frac{\pi}{3}$ , then the maximum value  $\tan A \tan B$  is (1993 – 2Marks)
- 8) General value of  $\theta$  satisfying the equation  $\tan^2 \theta + \sec 2\theta = 1$  is (1996 – 1Mark)
- 9) The real roots of the equation  $\cos^7 x + \sin^4 x = 1$  in the interval  $(-\pi, \pi)$  are (1997 – 2Marks)

## B: TRUE / FALSE

- 1) If  $\tan A = \frac{1 - \cos B}{\sin B}$ , then  $\tan 2A = \tan B$  (1981 – 1Mark)
- 2) There exists a value of  $\theta$  between 0 and  $2\pi$  that satisfies the equation  $\sin^4 \theta - 2 \sin^2 \theta - 1 = 0$ . (1984 – 1Mark)

## C :MCQs WITH ONE CORRECT ANSWER

- 1) If  $\tan \theta = -\frac{4}{3}$  then  $\sin \theta$  is (1979)
  - a)  $-\frac{4}{5}$  but not  $\frac{4}{5}$
  - b)  $\frac{4}{5}$  or  $-\frac{4}{5}$
  - c)  $\frac{4}{5}$  but not  $-\frac{4}{5}$
  - d) None of These
- 2) If  $\alpha + \beta + \gamma = 2\pi$  (1979)
  - a)  $\tan \frac{\alpha}{2} + \tan \frac{\beta}{2} + \tan \frac{\gamma}{2} = \tan \frac{\alpha}{2} \tan \frac{\beta}{2} \tan \frac{\gamma}{2}$
  - b)  $\tan \frac{\alpha}{2} \tan \frac{\beta}{2} + \tan \frac{\beta}{2} \tan \frac{\gamma}{2} + \tan \frac{\gamma}{2} \tan \frac{\alpha}{2} = 1$
  - c)  $\tan \frac{\alpha}{2} + \tan \frac{\beta}{2} + \tan \frac{\gamma}{2} = -\tan \frac{\alpha}{2} \tan \frac{\beta}{2} \tan \frac{\gamma}{2}$
  - d) None of These
- 3) Given  $A = \sin^2 \theta + \cos^4 \theta$  then for all real values of  $\theta$  (1980)
  - a)  $1 \leq A \leq 2$
  - b)  $\frac{3}{4} \leq A \leq 1$
  - c)  $\frac{13}{16} \leq A \leq 1$
  - d)  $\frac{3}{4} \leq A \leq \frac{13}{16}$
- 4) The equation  $2 \cos^2 \frac{x}{2} \sin^2 x = x^2 + x^{-2}$  (1980)
  - a) no real solution

- b) one real solution
  - c) more than one real solution
  - d) None of these
- 5) The general solution to the trigonometric equation  $\sin x + \cos x = 1$  is given by (1981 – 2Marks)
- a)  $x = 2n\pi; n = 0, \pm 1, \pm 2 \dots$
  - b)  $x = 2n\pi + \frac{\pi}{2}, n = 0, \pm 1, \pm 2 \dots$
  - c)  $x = n\pi + (-1)^n \frac{\pi}{4}, n = 0, \pm 1, \pm 2 \dots$
  - d) none of these
- 6) The value of the expression  $\sqrt{3} \operatorname{cosec} 20^\circ - \sec 20^\circ$  is equal to (1988 – 2Marks)
- a) 2
  - b)  $2 \sin 20^\circ / \sin 40^\circ$
  - c) 4
  - d)  $2 \sin 20^\circ / \sin 40^\circ$