

Probability Assignment 3 (12.13.5.7)

Aditya Varun V (AI22BTECH11001)

Question

In an examination, 20 questions of true-false type are asked. Suppose a student tosses a fair coin to determine his answer to each question. If the coin falls heads, he answers 'true'; if it fails, he answers 'false'. Find the probability that he answers at least 12 questions correctly.

Solution

VARIABLE	DESCRIPTION
n	Number of questions in the examination
p	Probability of coin showing heads
X	Number of questions answered correctly

TABLE 1

X follows a binomial distribution,

$$X = \text{Bin}(n, p) \quad (1)$$

$$= \text{Bin}(20, 0.5) \quad (2)$$

The mean of X,

$$\mu = n \times p \quad (3)$$

$$= 10 \quad (4)$$

The Variance of X,

$$\sigma^2 = n \times p \times (1 - p) \quad (5)$$

$$= 5 \quad (6)$$

Applying the normal approximation to X for large values of n,

$$\Pr(a \leq X \leq b) = \int_{a-0.5}^{b+0.5} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(k-\mu)^2}{2\sigma^2}} dk \quad (7)$$

$$\Pr(12 \leq X \leq 20) = \int_{11.5}^{20.5} \frac{1}{\sqrt{10\pi}} e^{-\frac{(k-10)^2}{10}} dk \quad (8)$$

$$(9)$$

On computation,

$$\Pr(12 \leq X \leq 20) \approx 0.2512 \quad (10)$$