Solution Gate EC 29.2022

Aditya Vikram Singh*

Question 29 Let H(X) denote the entropy of a discrete random variable X taking K possible distinct real values. Which of the following statements is/are necessarily true?

(A) $H(X) \leq \log_2 K$ bits

(B) $H(X) \leq H(2X)$

(C) $H(X) \leq H(X^2)$

(D) $H(X) \leq H(2^X)$

Solution: If all the K distinct values of the variable have the same probability,

Random independent variable	value of R.V	Description
X	$X \in (x_1, x_2..., x_K)$	Value of the discrete variable X
K	$K \in N$	Total no. of distinct values

then Entropy will be maximum

$$H(X)max = \log_2 K \qquad (1)$$

but, we know that every value will have a different value so,

$$H(X) \leq \log_2 K \qquad (2)$$

Hence, Option(A) is correct

$$H(X) = \sum_{i=0}^{k} P_X x_i \log_2 \frac{1}{P_X x_i} \qquad (3)$$

$$(4)$$

Let's consider the discrete variable as follows

$X \in x_i$	$P_X x_i$
-1	$\frac{1}{4}$
0	$\frac{1}{2}$
1	$\frac{1}{4}$

$$H(X) = \frac{1}{4} \log_2 4 + \frac{1}{2} \log_2 2 + \frac{1}{4} \log_2 4 \qquad (5)$$

$$H(X) = 1.5 units \qquad (6)$$

Now $Y = 2X$

*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU GPL. Free and open source.

$Y \in y_i$	$P_Y y_i$
-2	$\frac{1}{4}$
0	$\frac{1}{2}$
2	$\frac{1}{4}$

$$H(Y) = \sum_{i=0}^{2} P_Y y_i \log_2 \frac{1}{P_Y y_i} \tag{7}$$

$$H(Y) = 1.5 units \tag{8}$$

$$H(Y) = H(2X) = H(X) \tag{9}$$

Hence, Option(B) is correct
Similarly on substituting $Y = X^2$

$Y \in y_i$	$P_Y y_i$
0	$\frac{1}{2}$
1	$\frac{1}{2}$

$$H(Y) = \sum_{i=0}^{1} P_Y y_i \log_2 \frac{1}{P_Y y_i} \tag{10}$$

$$H(Y) = 1 units \tag{11}$$

$$H(Y) = H(X^2) \leq H(X) \tag{12}$$

Hence, Option(C) is incorrect
Now for $Y = 2^X$
As the function Y is monotonically increasing, so for every distinct X, we get a distinct Y

$$H(Y) = H(2^X) = H(X) \tag{13}$$

Hence, Option(D) is correct
The ans is (A), (B), (D)