

Mapping the Topography of Earth's Core: Relationships Between Topography Variations and Seismic Waveforms



Marin Adrian Mag¹, Paula Koelemeijer^{1,2}, Tarje Nissen-Meyer^{1,3}, Cristophe Zaroli⁴


¹University of Oxford, ²RHUL, ³University of Exeter, ⁴University of Strasbourg

Contact: marin.mag@stx.ox.ac.uk


Motivation and Aim

- The Core-Mantle Boundary (CMB) is the most drastic discontinuity inside the Earth.
- Its shape affects the mantle dynamics and evolution.
- Even though its topography has been studied for more than 40 years, models still don't converge¹.
- We aim to produce a robust, multi-scale model of the CMB topography.

A collection of 6 published CMB topography models.
(Koelemeijer, 2020)





Ingredients for Inversion



CMB topography effect on waveforms

- Small perturbations in the topography of the CMB lead to waveform perturbations well approximated by time-domain shifts.



Ray paths of ScS phase. CMB topography is color-coded: blue=-0, white = 0.


- Backus-Gilbert SOLA^{2,3,4} inversion requires a linear relationship between the data and the model.
- Relation between waveforms (data) and CMB topography (model) can be linearized and used for BG-SOLA inversion.

Linearizing the relation between topography and waveforms

- The relation between waveform phase shifts and CMB topography perturbation in the linearized form is given by the following integral:

$$data = \int_{\Sigma} \underbrace{\text{Geometric sensitivity kernel } (K_d)}_{\text{(blue = positive, red = negative)}} \times \underbrace{\text{(Exaggerated) CMB topography } (\delta d)}_{\text{(blue = positive, red = negative)}} dS$$

- Computation of geometric kernels is done via the adjoint method^{5,6}
- Forward and backward wavefields are computed using the powerful yet efficient wave solver AxiSEM3D^{7,8}



Left: diagram showing the interaction between forward and backward fields for ScS phase at 20 degrees distance between station and event. Below: Shear wave volumetric sensitivity kernel



Geometric Sensitivity kernels

- The geometric sensitivity kernel can be split in 3 components:



$$K_d = K_{dV} + K_{d\Sigma_n} + K_{d\Sigma_t}$$

- $K_{d\Sigma_t}$ vanishes on solid-solid boundaries due to the tighter boundary conditions present there.
- K_{dV} is formed purely from volumetric sensitivities (density and elastic):


$$K_{dV} = \rho K_{\rho} + C :: K_C$$



K_{dV} for ScS at 20° and 30s main period evaluated at the CMB




K_d for PP at 94° and 30s main period evaluated at the 670km discontinuity



BG-SOLA Inversion


- The linearized relationship will be inverted using the Backus-Gilbert SOLA method (BG-SOLA)
- This method computes averages of possible models and provides a unique set of advantages:
 - Provides un-biased averages
 - Can invert locally
 - Easily parallelizable
 - Full uncertainty and resolution quantification



Comparison of inversion between SOLA and Damped Least Squared (DLS) methods (Zaroli, C., Koelemeijer, P. and Lambotte, S., 2017).

Future Work

- Finish kernel code framework
- Create framework for BG-SOLA
- Gather and process data
- Run BG-SOLA inversion
- Apply further constraints on model from independent normal modes data
- Compare with models from geodynamics simulations



Visual representation of the main steps of the Ph.D. project

References

- Koelemeijer, P., 2021. Toward consistent seismological models of the core-mantle boundary landscape. *Mantle convection and surface expressions*, pp.229-255.
- Backus, G., & Gilbert, F. (1968). The resolving power of gross earth data. *Geophysical Journal International*, 16 (2), 169–205.
- Zaroli, C., Koelemeijer, P. and Lambotte, S., 2017. Toward seeing the Earth's interior through unbiased tomographic lenses. *Geophysical Research Letters*, 44(22), pp.11-399.
- Zaroli, C., 2016. Global seismic tomography using Backus-Gilbert inversion. *Geophysical Supplements to the Monthly Notices of the Royal Astronomical Society*, 207(2), pp.876– 888.
- Liu, Q. and Tromp, J., 2008. Finite-frequency sensitivity kernels for global seismic wave propagation based upon adjoint methods. *Geophysical Journal International*, 174(1), pp.265-286.
- Fichtner, A., 2010. *Full seismic waveform modelling and inversion*. Springer Science & Business Media.
- Leng, K., Nissen-Meyer, T., & van Driel, M. (2016). Efficient global wave propagation adapted to 3-d structural complexity: a pseudospectral/spectral-element approach. *Geophysical Supplements to the Monthly Notices of the Royal Astronomical Society*, 207 (3), 1700–1721.
- Leng, K., Nissen-Meyer, T., Van Driel, M., Hosseini, K., & Al-Attar, D. (2019). AxiSEM3d: broad-band seismic wavefields in 3-d global earth models with undulating discontinuities. *Geophysical Journal International*, 217 (3), 2125–2146.

Acknowledgments

