



# Combining SOLA and Deterministic Linear Inferences

Adrian Marin Mag<sup>1</sup>, Christophe Zaroli<sup>2</sup>, Paula Koelemeijer<sup>1</sup>

<sup>1</sup>University of Oxford, <sup>2</sup>University of Strasbourg



1

## Introduction

### The Problem

- Poor seismic data coverage leads to non-uniqueness<sup>1</sup>.
- Non-uniqueness is typically mitigated by regularisations that impose strong prior constraints.
- Robust uncertainty and resolution quantification is often difficult and absent in inverse methods.

### Motivation

- Instead of finding a model, focus on constraining desired properties (Figure 1) of the model using mathematical inferences.
- This allows for a relaxation of prior regularisations (Figure 2).

2

## Methods (noise-free)

SOLA<sup>2</sup> (Subtractive Optimally Localized Averages) and DLI<sup>3</sup> are linear inference methods stemming from the work of Backus and Gilbert<sup>1</sup>.

### Research Question

- How are SOLA and Deterministic Linear Inferences (DLI) related and how can they be combined?

### SOLA

- Provides an *approximate property of the true model* without the need for prior model constraints<sup>3</sup>.
- Non-uniqueness errors are quantified in the resolving kernels<sup>5</sup> (Figure 4) (approximate mapping).

### DLI

- Provides bounds for the *true property of the true model* using deterministic prior constraints (hard prior, Figure 2).
- Non-uniqueness errors are quantified in property bounds.
- Results are interpreted through target kernels.




Figure 1: Example of properties. Column 1: True model. Column 2: Three examples of target kernels centered at 0.5 with width 0.2, each associated with a type of local average. Column 3: Each of the three types of local averages evaluated against the true model at 100 points in the domain of the model

Nonuniqueness: there are many (often infinite) models that fit the data. We assume that the true model fits the data.




Figure 2: Graph showing the relationships between the main spaces used in the noise-free SOLA and DLI methods. Ellipses represent spaces, subspaces and/or subsets. Dots represent elements of the space. Arrows indicate the direction of the relation between spaces, sets, or elements. \*SOLA inverse is for the case when unimodularity is not imposed on resolving kernels.

### Combined SOLA-DLI method

- Provides bounds on the true property of the true model as solution, but also uses resolving kernels to analyse trade-offs between physical parameters.

3

## Results




Figure 3: Least norm solution

- Least norm solution provides unique answer due to strong regularizations (Figure 3), while inferences give bounds thanks to more relaxed regularization (Figure 4).
- Inferences take into account lack of data sensitivity by construction and provide broad bounds in such cases (Figure 4).
- Some properties are better constrained than others (Figure 4).
- We can extract more than just local averages<sup>6</sup> (for eg. gradients and coefficients of basis functions) (Figure 4).
- Inference methods work naturally on undiscretised model spaces.




Figure 4: SOLA-DLI solutions for two different types of local average and a local gradient. First column: property bounds for the second physical parameter evaluated at 100 evenly spaced enquiry points. Second column: target and resolving kernels for each type of property

4


## Conclusions and Discussion

- Combining SOLA and DLI we can place the result interpretation on the target kernels, which are well known and consistent.
- Smart choices of target kernels can improve the precision of the results and their interpretability.
- Resolving kernels can provide additional information about spatial trade-offs and trade-offs between physical parameters.
- Currently we are working on introducing data noise and probabilistic prior information.
- These methods are best suited for obtaining rigorous uncertainty and resolution quantification on a linear problem or a linearizable problem that has small amounts of data and would otherwise rely on strong regularizations.


## References and Acknowledgements

- Backus, G.E. and Gilbert, J.F., 1967. Numerical applications of a formalism for geophysical inverse problems. *Geophysical Journal International*, 13(1-3), pp.247-276.
- Pijpers, F.P. and Thompson, M.J., 1994. The SOLA method for helioseismic inversion. *Astronomy and Astrophysics* (ISSN 0004-6361), vol. 281, no. 1, p. 231-240, 281, pp.231-240.
- Al-Attar, D., 2021. Linear inference problems with deterministic constraints. arXiv preprint arXiv:2104.12256
- Zaroli, C., 2019. Seismic tomography using parameter-free Backus-Gilbert inversion. *Geophysical Journal International*, 218(1), pp.619-630.
- Latalierre, F., Zaroli, C., Lambotte, S. and Maggi, A., 2022. Analysis of tomographic models using resolution and uncertainties: a surface wave example from the Pacific. *Geophysical Journal International*, 230(2), pp. 893-907.
- Backus, G.E., 1988. Bayesian inference in geomagnetism. *Geophysical Journal International*, 92(1), pp. 125-142.

AMM received funding from Royal Society grant RF\ER\RE210182 awarded to PK. PK acknowledges financial support from a Royal Society University Research Fellowship (URF\R1\180377). AMM acknowledges the funding received from the UKRI NERC DTP NE/S007474/1



See paper in  
revision



See GitHub page