

Part IA - Analysis I

Definitions

Lectured by W. T. Gowers

Lent 2015

Limits and convergence

Sequences and series in \mathbb{R} and \mathbb{C} . Sums, products and quotients. Absolute convergence; absolute convergence implies convergence. The Bolzano-Weierstrass theorem and applications (the General Principle of Convergence). Comparison and ratio tests, alternating series test. [6]

Continuity

Continuity of real- and complex-valued functions defined on subsets of \mathbb{R} and \mathbb{C} . The intermediate value theorem. A continuous function on a closed bounded interval is bounded and attains its bounds. [3]

Differentiability

Differentiability of functions from \mathbb{R} to \mathbb{R} . Derivative of sums and products. The chain rule. Derivative of the inverse function. Rolle's theorem; the mean value theorem. One-dimensional version of the inverse function theorem. Taylor's theorem from \mathbb{R} to \mathbb{R} ; Lagrange's form of the remainder. Complex differentiation. [5]

Power series

Complex power series and radius of convergence. Exponential, trigonometric and hyperbolic functions, and relations between them. *Direct proof of the differentiability of a power series within its circle of convergence*. [4]

Integration

Definition and basic properties of the Riemann integral. A non-integrable function. Integrability of monotonic functions. Integrability of piecewise-continuous functions. The fundamental theorem of calculus. Differentiation of indefinite integrals. Integration by parts. The integral form of the remainder in Taylor's theorem. Improper integrals. [6]

Contents

1	The real number system	3
2	Convergence of sequences	4
2.1	Sums, products and quotients	4

1 The real number system

Definition (Field). A *field* is a set X with two binary operations $+$ and \times that satisfies all the familiar properties satisfied by addition and multiplication in \mathbb{Q} , namely

- Associativity: $\forall a, b, c \in X, a + (b + c) = (a + b) + c$ and $a \times (b \times c) = (a \times b) \times c$
- Commutativity: $\forall a, b \in X, a + b = b + a$ and $a \times b = b \times a$
- Identity: $\exists 0, 1 \in X$ such that $\forall a, a + 0 = a$ and $a \times 1 = a$.
- Inverses: $\forall a \in X, \exists (-a) \in X$ such that $a + (-a) = 0$. If $a \neq 0$, then $\exists a^{-1}$ such that $a \times a^{-1} = 1$.
- Distributivity: $\forall a, b, c \in F, a \times (b + c) = (a \times b) + (a \times c)$

Definition (Totally ordered set). An (*totally*) *ordered set* is a set X with a relation $<$ that satisfies

- (i) Transitivity: if $x, y, z \in X, x < y$ and $y < z$, then $x < z$
- (ii) Trichotomy: if $x, y \in X$, exactly one of $x < y, x = y, y < x$ holds

Definition (Ordered field). An *ordered field* is a field \mathbb{F} with a relation $<$ that makes \mathbb{F} into an ordered set such that

- (i) if $x, y, z \in \mathbb{F}$ and $x < y$, then $x + z < y + z$
- (ii) if $x, y, z \in \mathbb{F}, x < y$ and $z > 0$, then $xz < yz$

Definition (Least upper bound). Let X be an ordered set and let $A \subseteq X$. An *upper bound* for A is an element $x \in X$ such that $\forall a \in A (a \leq x)$. If A has an upper bound, then we say that A is *bounded above*.

An upper bound x for A is a *least upper bound* or *supremum* if nothing smaller than x is an upper bound. That is, we need

- (i) $\forall a \in A (a \leq x)$
- (ii) $\forall y < x (\exists a \in A (a \geq y))$

We usually write $\sup A$ for the supremum of A when it exists. If $\sup A \in A$, then we call it $\max A$, the maximum of A .

Definition (Least upper bound property). An ordered field has the *least upper bound property* if every non-empty subset of \mathbb{F} that is bounded above has a supremum.

2 Convergence of sequences

Definition (Sequence). A *sequence* is, formally, a function $a : \mathbb{N} \rightarrow \mathbb{R}$ (or \mathbb{C}). Usually (i.e. always), we write a_n instead of $a(n)$. Instead of a , (a_n) , $(a_n)_1^\infty$ or $(a_n)_{n=1}^\infty$ to indicate it is a sequence.

Definition (Convergence of sequence). Let (a_n) be a sequence and $\ell \in \mathbb{R}$. Then a_n *converges to ℓ* , *tends to ℓ* , or $a_n \rightarrow \ell$, if

$$\forall \varepsilon > 0 \exists N \forall n \geq N : |a_n - \ell| < \varepsilon.$$

Definition (Bounded sequence). A sequence (a_n) is *bounded*

$$\exists C \forall n : |a_n| \leq C.$$

A sequence is *eventually bounded* if

$$\exists C \exists N \forall n \geq N : |a_n| \leq C.$$

2.1 Sums, products and quotients