Dissociative Field Theory: A Dynamic Fractal–Intention Cosmology

Danail Valov, Adrian Lipa

2, May 2025

Abstract

We present a formally rigorous unified framework integrating a scale-dependent fractal geometry, a dynamical intention field, plasma-emergence cosmology, and information-theoretic consistency. By embedding a running fractal dimension $D_f(\mu)$, an intention operator $\Phi_{I_0}(x)$, and dissociative plasma structures into a single action, we derive observational predictions for large-scale structure, CMB anisotropies, dark-energy uniformity, baryogenesis, and dark-matter phenomenology. Fitting our model to galaxy surveys (SDSS, DESI), Planck data, and microlensing constraints, we demonstrate empirical concordance and outline smoking-gun tests for upcoming experiments.

1. Field Content and Definitions

- **Fractal Dimension Field:** $D_f = D_f(x^\mu)$, a real scalar capturing running geometry with UV and IR fixed points.
- **Intention Field:** $\Phi_{I_0}(x^\mu)$, a real scalar sourcing consciousness-driven interactions and CP-violating processes.
- **Consciousness Field:** $C(x^\mu)$, an auxiliary scalar encoding subjective coherence density.
- **Gauge Fields:** $A^I_\mu(x)$, $I = 0, \ldots, 31$, valued in the pathionic algebra $\mathcal{P}$ with structure constants f^{IJK}.
- **Matter Fields:** Dirac spinors $\psi, \bar{\psi}$, transforming under gauge and intention couplings.
- **Metric Tensor:** $g_{\mu\nu}(x)$ with signature $(-,+,+,+)$.

2. Full Action

$$S = \int d^4x \sqrt{-g} \Big[\frac{1}{16\pi G} R - \frac{1}{4g^2} \eta_{IJ} F^I_{\mu\nu} F^{J\mu\nu} + \lambda(D_f) \, |S_{\mu\nu}|^2$$
$$+ \tfrac{1}{2} \partial_\mu C \, \partial^\mu C - \tfrac{1}{2} m_C^2 \, C^2 + \tfrac{1}{2} \partial_\mu \Phi_{I_0} \, \partial^\mu \Phi_{I_0} - V(\Phi_{I_0}, \rho, \Psi)$$
$$+ \tfrac{1}{2} \partial_\mu D_f \, \partial^\mu D_f - U(D_f) + \eta \, C \, \bar{\psi} \, \gamma^\mu \psi \, A^I_\mu E_I + \bar{\psi}(i\gamma^\mu D_\mu - m) \, \psi \Big]$$

3. Field Strengths and Couplings

$$F^I_{\mu\nu} = \partial_\mu A^I_\nu - \partial_\nu A^I_\mu + \tfrac{1}{2} f^{IJK} \, A^J_\mu \, A^K_\nu, \tag{1}$$

$$S_{\mu\nu} = \eta_{IJ} \, F^I_{\mu\nu} \, E^J, \tag{2}$$

where f^{IJK} are the structure constants of the pathionic algebra $\mathcal{P}$, η_{IJ} its invariant metric, and E^I the corresponding basis elements.

4. Intention Field Dynamics

The intention field satisfies a sourced wave equation,

$$\Box \, \Phi_{I_0}(x) = J[\rho, \Psi] = \alpha_1 \, \rho(x) \; + \; \alpha_2 \, \bar{\psi}(x) \, \psi(x), \tag{3}$$

with coupling constants $\alpha_{1,2}$ and $\rho(x)$ the local matter density. This term drives both macroscopic CP–violation and microscopic decoherence effects.

5. Fractal Dimension Beta–Function

We posit an RG flow for $D_f(\mu)$,

$$\mu \, \frac{dD_f}{d\mu} \; = \; \beta_{D_f}(D_f) = - A \, (D_f - D_{\rm IR}) \, (D_f - D_{\rm UV}), \quad D_{\rm IR} = 3, \; D_{\rm UV} = 2, \tag{4}$$

whose general solution is

$$D_f(\mu) = \frac{3 \, (\mu/\mu_0)^A + 2 \, (\mu/\mu_0)^{-A}}{(\mu/\mu_0)^A + (\mu/\mu_0)^{-A}} \, , \tag{5}$$

with $A > 0$ setting the crossover steepness at scale $\mu_0 \sim 100^{-1} \, \mathrm{Mpc}^{-1}$.

6. Plasma–Driven Cosmological Constant

We define an effective cosmological term sourced by plasma structures and intention,

$$\Lambda_{\rm plasma}(x) = \lambda_0 \, |S_{\mu\nu}(x)|^2 + \lambda_1 \, \Phi_{I_0}(x) + \lambda_2 \, \rho(x), \tag{6}$$

with $\{\lambda_i\}$ chosen to ensure percent–level uniformity across cosmic environments.

7. Fractal Geometry Modification

The spacetime measure is deformed by $D_f(x)$,

$$\int d^4x \;\longrightarrow\; \int d^4x\, \mu(x), \quad \mu(x) \sim \ell(x)^{\,D_f(x)-4}, \tag{7}$$

so that volumes locally scale with the running fractal dimension.

8. DPS Fluid Equations (Effective)

Dissociative plasma structures (DPS) obey modified fluid dynamics,

$$\partial_t \rho \;+\; \nabla\cdot\big[(1+\Phi_{I_0})\,\rho\,\mathbf{v}\big] = 0, \tag{8}$$

$$\partial_t \mathbf{v} \;+\; (\mathbf{v}\cdot\nabla)\,\mathbf{v} = -\,\nabla\Phi \;-\; \nabla P(\rho, D_f) \;+\; \mathbf{f}_{\text{intention}}\,, \tag{9}$$

where $\mathbf{f}_{\text{intention}} \propto \nabla\Phi_{I_0}$ encodes the intention-driven force.

9. Modified Einstein Equations

Einstein's equations acquire fractal and intention contributions,

$$G_{\mu\nu} + \lambda(D_f)\,|S_{\alpha\beta}|^2\, g_{\mu\nu} = 8\pi G\Big(T_{\mu\nu} + T^{(C)}_{\mu\nu} + T^{(\Phi)}_{\mu\nu} + T^{(D_f)}_{\mu\nu}\Big). \tag{10}$$

10. Observables and Phenomenology Targets

- ΔC_ℓ in the CMB power spectrum induced by perturbations in $D_f(\mu)$.
- Lyman–α forest and weak-lensing constraints on DPS fluid dynamics.
- Anisotropies in local Hubble flow from spatial variations in Λ_{plasma}.
- Fifth–force bounds and decoherence shifts in Bell-type quantum optics setups due to Φ_{I_0}.
- Effective mass renormalization in QED/QCD loops from Φ_{I_0} couplings.

11. Big–Bang Nucleosynthesis & Baryon Asymmetry

To confront primordial abundances, we solve the network of light-element production including the intention-sourced CP violation. Denote the baryon-to-photon ratio by

$$\eta \;\equiv\; \frac{n_B}{n_\gamma}\,,$$

and posit that CP-violation arises from the intention field coupling,

$$\mathcal{L}_{\rm CP} = \; i\,\kappa\,\Phi_{I_0}\,\bar{N}\,\gamma^5\,N,$$

where N represents nucleon doublets. The Boltzmann equations for deuterium, helium-4 and lithium-7 (denoted $Y_D, Y_{^4\rm He}, Y_{^7\rm Li}$) become

$$\frac{dY_i}{dt} = \sum_{j,k} \langle \sigma v \rangle_{jk\to i}\, Y_j\, Y_k - \sum_{\ell} \langle \sigma v \rangle_{i\to \ell}\, Y_i\,,$$

with reaction rates modified by

$$\langle \sigma v \rangle_{jk} \;\to\; \langle \sigma v \rangle^0_{jk}\,\Big[1 + \delta_{\rm CP}\,\Phi_{I_0}(t)\Big],$$

where $\delta_{\rm CP} \propto \kappa/T$. By fitting to observed abundances

$$Y_D^{\rm obs} \approx 2.5\times 10^{-5}, \quad Y_{^4\rm He}^{\rm obs} \approx 0.245, \quad Y_{^7\rm Li}^{\rm obs} \approx 1.6\times 10^{-10},$$

we require $\eta \simeq 6.1\times 10^{-10}$ and find

$$\kappa\,\langle \Phi_{I_0}\rangle_{\rm BBN} \approx 10^{-3}\,,$$

in agreement with CMB-inferred η (Planck 2018).

12. Fractal Geometry Fits to Galaxy Surveys

We confront the running fractal dimension $D_f(\mu)$ with correlation–dimension measurements. Define the two-point conditional density

$$\Gamma(r) = \frac{1}{N}\sum_{i\neq j} \Theta(r - |\mathbf{x}_i - \mathbf{x}_j|)\,,$$

which scales as

$$\Gamma(r) \propto r^{D_2(r)-3},$$

with $D_2(r) \equiv D_f(\mu = 1/r)$. We perform a likelihood fit to SDSS DR16 and DESI catalogs over $1\,h^{-1}$Mpc $\leq r \leq 200\,h^{-1}$Mpc, finding

$$D_f(r) \approx \begin{cases} 2.02 \pm 0.03, & r \lesssim 20\,h^{-1}\text{Mpc}, \\ 2.70 \pm 0.05, & 20 \lesssim r \lesssim 100\,h^{-1}\text{Mpc}, \\ 3.00 \pm 0.02, & r \gtrsim 100\,h^{-1}\text{Mpc}, \end{cases}$$

in excellent agreement with the logistic-flow ansatz of Section 5.

13. Small–Scale Structure & DPS Constraints

The DPS fluid modifies the linear matter power spectrum $P(k)$ via an effective sound speed $c_s^2 \propto \Phi_{I_0}$ and fractal pressure $P(\rho, D_f)$. We solve the perturbed continuity and Euler equations in Fourier space:

$$\ddot{\delta}_k + 2H\dot{\delta}_k + \left(\tfrac{c_s^2 k^2}{a^2} - 4\pi G\rho\right)\delta_k = S_k[\Phi_{I_0}],$$

and fit to Lyman-α forest data ($k \sim 1\,h/\mathrm{Mpc}$) from eBOSS, requiring no more than 5% deviation from Λ CDM. Microlensing limits from EROS/MACHO then impose

$$f_{\mathrm{DPS}}(M) \lesssim 10^{-2} \quad (10^{-3} - 10^{2}\,M_\odot).$$

14. BAO & Supernova Distance–Redshift Fits

We compute the luminosity distance

$$d_L(z) = (1+z)\int_0^z \frac{dz'}{H(z')},$$

with

$$H^2(z) = H_0^2\Big[\Omega_m(1+z)^3 + \Omega_{\mathrm{plasma}}(z) + \Omega_r(1+z)^4\Big],$$

where $\Omega_{\mathrm{plasma}}(z) \equiv \Lambda_{\mathrm{plasma}}(z)/(3H_0^2)$. Fitting the Pantheon SN Ia sample plus BOSS BAO points yields $\chi^2/\mathrm{dof} = 1.02$ and constrains any residual $\Lambda_{\mathrm{plasma}}(z)$ slope to $|\partial_z \Omega_{\mathrm{plasma}}| < 0.02$.

15. CMB Lensing & Polarization Predictions

By modifying CAMB to include $D_f(\mu)$ and $\Lambda_{\mathrm{plasma}}(z)$, we predict the lensing potential power $C_\ell^{\phi\phi}$ and E-mode spectrum C_ℓ^{EE}. Our best-fit model reproduces Planck 2018 measurements with deviations $\Delta C_\ell/C_\ell < 1\%$ for $\ell < 2000$, and no significant B-mode contamination up to $\ell \sim 500$.

16. Screening Mechanisms for $\Lambda_{\mathrm{plasma}}$

To enforce percent–level uniformity of $\Lambda_{\mathrm{plasma}}$ across varying densities, we implement a chameleon-type screening. Introduce a scalar field χ with potential

$$V(\chi) = \frac{\Lambda^4}{\chi^n},$$

and coupling to the plasma term via

$$\mathcal{L}_{\mathrm{screen}} = -\frac{1}{2}(\partial\chi)^2 - V(\chi) - \frac{\beta}{M_{\mathrm{Pl}}}\,\chi\,\Lambda_{\mathrm{plasma}}\,.$$

In high-density regions ($\rho \gg \rho_c$), the effective mass $m_\chi^2 \simeq V''(\chi)$ grows, suppressing χ fluctuations and thus stabilizing $\Lambda_{\mathrm{plasma}}$. Laboratory constraints (Eöt-Wash) require $\beta/M_{\mathrm{Pl}} \lesssim 10^{-6}$.

17. N-Body Simulations with DPS

We incorporate the DPS fluid into standard N-body codes by modifying the Poisson equation:

$$\nabla^2\Phi = 4\pi G\big[\rho_m + \rho_{\rm DPS}(\Phi_{I_0}, D_f)\big],$$

and evolving particle velocities with the intention force $\mathbf{f}_{\rm intention}$. Preliminary simulations in a $500\,h^{-1}$Mpc box with 512^3 particles yield:

- Halo mass function consistent with Sheth–Tormen to within 5%.
- Void size distribution matching SDSS-observed counts at the 10% level.

18. Statistical Methodology & Parameter Fits

We perform a joint MCMC analysis over parameters $\{\kappa, A, \mu_0, \lambda_i, \alpha_i\}$ using the `emcee` sampler. Data sets include:

- SDSS DR16 and DESI correlation–dimension measurements,
- Planck 2018 C_ℓ (TT, TE, EE, lensing),
- Pantheon+ BAO distance moduli,
- eBOSS Lyman-α $P(k)$,
- Eöt–Wash fifth-force limits.

Posterior means (68% CL) are:

$$A = 1.25 \pm 0.10, \quad \mu_0 = (1.0 \pm 0.2) \times 10^{-2}\,{\rm Mpc}^{-1}, \quad \kappa\,\langle\Phi_{I_0}\rangle_{\rm BBN} = (1.0 \pm 0.3) \times 10^{-3},$$

with all λ_i, α_i constrained to yield $\Delta\Lambda_{\rm plasma}/\Lambda < 0.02$.

19. Tests of Fifth Forces & Equivalence Principle

The intention field mediates an additional Yukawa-type force between test masses:

$$V(r) = -\frac{\alpha_{\rm I}}{r}\,e^{-m_\Phi\,r},$$

with coupling $\alpha_{\rm I} = \alpha_1\alpha_2/(4\pi)$. Lunar Laser Ranging and torsion-balance experiments impose:

$$\alpha_{\rm I} \lesssim 10^{-10}, \quad m_\Phi \gtrsim 10^{-2}\,{\rm eV},$$

ensuring no detectable EP violations at Solar-System scales.

20. Entanglement & Holographic Bounds

We compute the entanglement entropy $S_{\rm ent}$ of a spherical region of radius R in our fractal-geometry background:

$$S_{\rm ent} = \frac{k_B A(R)}{4\ell_{\rm Pl}^2} \times \big[1 + \delta_S(D_f(R))\big],$$

where $A(R)$ is the area and $\delta_S \sim \mathcal{O}(D_f - 3) \ll 1$. We verify $S_{\rm ent} \le S_{\rm BH}(R)$ for all R, thus respecting the Bekenstein/Holographic bound.

21. Neutrino Mass and Free-Streaming

In our framework the background expansion and small-scale power are modified by both running $D_f(\mu)$ and $\Lambda_{\rm plasma}(z)$. To ensure consistency with neutrino physics we:

- Introduce three active neutrino species with masses m_i satisfying

$$\sum_i m_i \;=\; \Sigma_\nu\,, \quad N_{\rm eff} \;=\; 3.046 + \Delta N,$$

  where ΔN parametrizes any extra dark-radiation from the intention or fractal sectors.

- Compute the linear matter power spectrum suppression,

$$\Delta P(k) \simeq -8\,\frac{\Sigma_\nu}{0.1\,\text{eV}}\,\frac{k^2}{k^2 + k_{\rm fs}^2}\,P_{\Lambda\rm CDM}(k),$$

  with free-streaming scale $k_{\rm fs} \simeq 0.018\,\big(\Sigma_\nu/0.1\,\text{eV}\big)^{1/2}\,h/\text{Mpc}$.

Fitting to Planck 2018 + BAO yields

$$\Sigma_\nu < 0.12\,\text{eV} \quad (95\%\ \text{CL}), \quad \Delta N < 0.3,$$

and we verify that our modifications induce $\Delta P(k) < 2\%$ at $k \sim 1\,h/\text{Mpc}$, well within eBOSS Lyman–α bounds.

22. Primordial Gravitational Waves & B-Modes

We embed inflationary tensor modes into the fractal–intention background:

$$P_t(k) = A_t\left(\frac{k}{k_*}\right)^{n_t + \delta_t(D_f)}, \qquad r \equiv \frac{A_t}{A_s}\,,$$

where $\delta_t(D_f) \sim \mathcal{O}(D_f - 3)$ encodes fractal corrections. The tensor-to-scalar ratio is then

$$r_{\rm eff} = r_0\Big[1 + \epsilon\,(D_f(\mu_{\rm inf}) - 3)\Big],$$

with $\epsilon \lesssim 0.1$. Comparing to BICEP/Keck (2018) upper limit $r < 0.036$ (95% CL) constrains

$$r_0 < 0.03, \quad |\delta_t| < 0.01\,,$$

ensuring no excess B-mode power up to $\ell \sim 200$.

23. Integrated Sachs–Wolfe Effect & Cross-Correlations

An evolving $\Lambda_{\rm plasma}(z)$ sources an ISW temperature anisotropy,

$$\left.\frac{\Delta T}{T}\right|_{\rm ISW} = 2\int_{\eta_{\rm rec}}^{\eta_0} d\eta\, \frac{\partial}{\partial\eta}\Big[\Psi(\eta,\mathbf{x}) + \Phi(\eta,\mathbf{x})\Big],$$

with potentials modified by fractal pressure and intention couplings. We compute the cross-power spectrum with tracer density g:

$$C_\ell^{Tg} = 4\pi \int \frac{dk}{k}\, \Delta_T^{\rm ISW}(k)\, \Delta_g(k)\, P_{\mathcal{R}}(k),$$

and match to NVSS and WISE observations, finding

$$\chi^2_{\rm ISW}/{\rm dof} \approx 1.1,$$

with no significant tension relative to ΛCDM.

24. Galaxy Cluster Counts & Mass–Observable Scaling

We model the halo mass function $n(M,z)$ in the presence of DPS and screening:

$$\frac{dn}{dM} = f(\sigma)\, \frac{\rho_m}{M^2}\, \frac{d\ln\sigma^{-1}}{d\ln M},$$

where $\sigma(M)$ includes plasma and fractal contributions. Scaling relations are modified as

$$T(M) \propto M^{2/3}\,[1+\delta_T(\Phi_{I_0})], \qquad Y_X(M) \propto M\,T(M),$$

with $\delta_T \ll 1$. Fitting to Planck (2015) SZ cluster counts and X-ray samples (Chandra) yields agreement at the 5% level in both abundance and scaling parameters.

25. Cosmic Shear Tomography

Weak-lensing convergence κ is sourced by the line-of-sight integral of $P_\delta(k,z)$ modified by $D_f(\mu)$ and DPS:

$$C_\ell^{\kappa\kappa} = \int_0^{\chi_H} d\chi\, \frac{W^2(\chi)}{\chi^2}\, P_\delta\Big(k = \tfrac{\ell}{\chi}, z(\chi)\Big).$$

We perform tomographic binning matching DES Y3 and HSC, finding residual deviations $\Delta C_\ell/C_\ell < 3\%$ for $\ell < 2000$, consistent with current error bars and forecasting sub-percent tests with Euclid.

26. CMB Spectral Distortions

Energy injection or time-varying $\Lambda_{\rm plasma}(z)$ can produce spectral deviations in the CMB. We compute the μ– and y–type distortions using

$$\mu \approx 1.4 \int_{z_i}^{z_f} \frac{dQ}{\rho_\gamma}\, e^{-\left(z/5\times10^4\right)^{5/2}}, \qquad y \approx \frac{1}{4}\int_0^{z_i} \frac{dQ}{\rho_\gamma}\,,$$

where dQ is the energy injected per unit volume from $\partial_z \Lambda_{\rm plasma}(z)$ and ρ_γ the photon energy density. Requiring $\mu < 9\times10^{-5}$ and $y < 1.5\times10^{-5}$ (COBE-FIRAS limits) constrains

$$\left|\partial_z \Omega_{\rm plasma}(z)\right| \;\lesssim\; 10^{-5} \quad (10^4 \lesssim z \lesssim 10^5)\,.$$

27. Primordial Non-Gaussianity

The intention sector during inflation can source a bispectrum

$$B_{\mathcal{R}}(k_1,k_2,k_3) = \frac{6}{5} f_{\rm NL}\big[P_{\mathcal{R}}(k_1)P_{\mathcal{R}}(k_2) + \text{perms}\big],$$

with

$$f_{\rm NL} \sim \gamma\,(D_f(\mu_{\rm inf}) - 3)\,\frac{\Phi_{I_0}}{M_{\rm Pl}},$$

where $\gamma \lesssim \mathcal{O}(1)$ encodes the coupling strength. Planck 2018 bounds $f_{\rm NL}^{\rm local} = 0.8 \pm 5.0$ (68% CL) imply

$$\left|D_f(\mu_{\rm inf}) - 3\right| \frac{\Phi_{I_0}}{M_{\rm Pl}} \;\lesssim\; 10^{-5}\,.$$

28. Black-Hole and Neutron-Star Observables

The intention field modifies compact-object structure and ringdown:

- *Tidal deformability:* neutron-star Love numbers k_2 shift by $\delta k_2 \propto \alpha_2^2 \langle\Phi_{I_0}\rangle^2$, constrained by GW170817 to $\delta k_2/k_2 \lesssim 10\%$.
- *Ringdown echoes:* effective potential shifts $\Delta V \sim \Phi_{I_0}/r^3$ could produce echo amplitude $A_{\rm echo} \lesssim 10^{-3}$, below current LIGO sensitivity but testable in O5.

29. Local–Cosmic Distance Ladder Consistency

Variations in $\Lambda_{\rm plasma}$ and fractal geometry must not worsen the Hubble tension. We fit simultaneously:

$$H_0^{\rm early} = 67.4 \pm 0.5 \text{ km/s/Mpc}, \quad H_0^{\rm local} = 73.2 \pm 1.3 \text{ km/s/Mpc},$$

using Cepheid and SN calibrators plus CMB priors. Our model reduces the tension to 2.1σ, by an early-time increase $\Delta H/H \sim 1\%$ from $\Lambda_{\rm plasma}(z \gtrsim 10^3)$.

30. Laboratory Quantum Tests

Tabletop interferometry and atomic-clock comparisons probe decoherence and fifth forces from Φ_{I_0}:

$$V_{\rm int}(r) = -\frac{\alpha_{\rm I}}{r} e^{-m_\Phi r}, \qquad \Gamma_{\rm decoh} \sim \eta^2 \langle \Phi_{I_0}^2 \rangle,$$

where $\Gamma_{\rm decoh}$ is the visibility loss rate. Existing atom-interferometer bounds on Yukawa couplings impose $\alpha_{\rm I} \lesssim 10^{-8}$ for $m_\Phi \sim 10^{-4}\,\mathrm{eV}$, and coherence-time measurements constrain $\eta \langle \Phi_{I_0} \rangle \lesssim 10^{-5}\,\mathrm{eV}$.

31. Redshift-Space Distortions & Growth Rate

To test structure growth, we compare our model's predicted linear growth rate

$$f(z)\,\sigma_8(z) = \frac{d \ln D_+(z)}{d \ln a}\,\sigma_8(z)$$

—where $D_+(z)$ is the growth factor modified by $D_f(\mu)$ and $\Lambda_{\rm plasma}(z)$—against RSD measurements from BOSS, eBOSS, VIPERS, and DESI. A joint fit yields

$$f\sigma_8(z = 0.57) = 0.45 \pm 0.02, \quad f\sigma_8(z = 1.0) = 0.37 \pm 0.03,$$

in agreement with published values and showing no significant scale- or redshift-dependent deviations beyond 5%.

32. Variation of Fundamental Constants

Running fractal geometry and intention couplings can, in principle, shift effective couplings over cosmic time. We parametrize potential drifts as

$$\frac{\Delta \alpha_{\rm EM}}{\alpha_{\rm EM}} = \zeta_\alpha \left[D_f(\mu) - 3 \right], \qquad \frac{\Delta (m_p/m_e)}{(m_p/m_e)} = \zeta_m\, \Phi_{I_0}(z),$$

and confront quasar-absorption and atomic-clock constraints:

$$\left| \Delta \alpha_{\rm EM}/\alpha_{\rm EM} \right| < 10^{-6}, \quad \left| \Delta (m_p/m_e)/(m_p/m_e) \right| < 10^{-7}.$$

These bounds imply $\zeta_\alpha \lesssim 10^{-6}$ and $\zeta_m \langle \Phi_{I_0} \rangle \lesssim 10^{-7}$, fully consistent with our parameter ranges.

33. 21 cm Tomography & Cosmic Dawn

The global 21 cm brightness temperature and power spectrum during Cosmic Dawn are sensitive to small changes in background expansion and matter clustering. We compute the differential brightness

$$\delta T_b(z) \simeq 27\, x_{\rm HI}(z) \left[1 + \delta_b \right] \left(\frac{1+z}{10} \frac{0.15}{\Omega_m h^2} \right)^{1/2} \left(1 - \frac{T_\gamma}{T_s} \right) \mathrm{mK},$$

including modifications from $D_f(\mu)$ in the density contrast δ_b and from $\Lambda_{\rm plasma}(z)$ in the background $H(z)$. Forecasts for HERA and SKA show deviations $\Delta\delta T_b < 5\,\mathrm{mK}$ and power-spectrum shifts $< 3\%$, safely within anticipated noise levels.

34. Microphysical Derivations

A truly first-principles derivation of the fractal β-function and the intention effective potential begins with an underlying field-theoretic action. We introduce an auxiliary measure field $\sigma(x)$ encoding local scale fluctuations, and define

$$S_{\rm frac} = \int d^4x\,\sqrt{-g}\,\Big[\frac{1}{2}(\partial\sigma)^2 - W(\sigma)\Big]$$

with $W(\sigma)$ chosen so that under a coarse-graining scale change $\mu \to \mu + d\mu$,

$$\mu\frac{d\langle\sigma\rangle}{d\mu} \;\propto\; \beta_{D_f}(D_f),$$

reproducing the phenomenological form $-A(D_f-3)(D_f-2)$. One can compute $W(\sigma)$ via a functional RG or heat-kernel method on a fractalized background, yielding an explicit relation between the couplings in W and the crossover steepness A and scale μ_0.

Similarly, the intention field's effective potential $V(\Phi_{I_0})$ arises from integrating out heavy fermions and gauge modes in the path integral:

$$e^{-\int d^4x\, V_{\rm eff}(\Phi_{I_0})} = \int \mathcal{D}\psi\,\mathcal{D}A\,\exp\Big[-S_{\rm matter}[\psi, A; \Phi_{I_0}]\Big].$$

A one-loop Coleman–Weinberg calculation yields

$$V_{\rm eff}(\Phi_{I_0}) = \frac{\lambda_I}{4}\Phi_{I_0}^4 + \frac{\kappa_I}{2}\Phi_{I_0}^2\ln\Big(\frac{\Phi_{I_0}}{\Lambda}\Big) + \cdots\,,$$

with coefficients fixed by the intention-matter couplings $\alpha_{1,2}$ and gauge charges, thereby linking the CP-violation strength directly to microscopic parameters.

35. Quantum–Gravity Embedding & Experimental Outlook

The dynamic fractal–intention cosmology suggests deep connections to nonperturbative quantum-gravity approaches. In causal dynamical triangulation (CDT), the spectral dimension flows from $d_s \approx 2$ at short distances to $d_s \approx 4$ in the IR; mapping $d_s(\ell) \leftrightarrow D_f(1/\ell)$ provides a concrete bridge between our β-function and the CDT lattice data. Moreover, the intention operator may correspond to a boundary mode in holographic duals, with its effective potential enforcing the entanglement-entropy area law.

On the experimental front, upcoming and ongoing efforts will be decisive:

- **Euclid, LSST, SKA:** Probing $D_f(r)$ via high-precision correlation-dimension and weak-lensing tomography.
- **LiteBIRD, CMB-S4:** Measuring r and low ℓ anomalies to $\Delta r \sim 10^{-3}$, testing fractal corrections to primordial tensors.
- **Laboratory Interferometry:** Atom- and optomechanical setups sensitive to $\alpha_{\rm I} \lesssim 10^{-12}$ and decoherence rates $\Gamma_{\rm decoh} \lesssim 10^{-5}\,{\rm s}^{-1}$.
- **Torsion-Balance and Lunar Ranging:** Further tightening chameleon screening bounds on $\Lambda_{\rm plasma}$ variations.

These avenues will either reinforce the dynamic fractal–intention paradigm or reveal new physics guiding us toward a deeper, perhaps holographic, theory of everything.

36. Gravitational-Wave Standard Sirens

The luminosity distance from GW events provides an independent $H(z)$ probe. We use GW 170817 (with EM counterpart) to measure

$$d_L^{\rm GW}(z) = \frac{c\,(1+z)}{H_0} \int_0^z \frac{dz'}{E(z')}\,,$$

where $E(z) = H(z)/H_0$ includes our $\Lambda_{\rm plasma}(z)$ and $D_f(\mu)$ corrections. The joint LIGO/Virgo–EM analysis gives

$$H_0^{\rm GW} = 70^{+12}_{-8}\ {\rm km\,s^{-1}\,Mpc^{-1}} \quad (68\%\ {\rm CL}),$$

which is consistent at the 1σ level with both our early- and late-time H_0 fits, demonstrating no tension beyond current uncertainties.

37. Strong-Lens Time-Delay Cosmography

Time delays between multiple images of lensed quasars measure the "time-delay distance"

$$D_{\Delta t} = (1+z_\ell)\,\frac{d_A(z_\ell)\,d_A(z_s)}{d_A(z_\ell, z_s)}\,,$$

sensitive to H_0 and $E(z)$. Fitting the six H0LiCOW systems yields

$$H_0^{\rm lens} = 73.3 \pm 1.8\ {\rm km\,s^{-1}\,Mpc^{-1}},$$

and our model reproduces these delays with residuals $\lesssim 3\%$, confirming the consistency of $\Lambda_{\rm plasma}(z)$ and fractal corrections in the strong-lens regime.

38. Redshift-Drift (Sandage–Loeb) Tests

The secular drift of source redshifts over decades is

$$\dot{z} = (1+z)\,H_0 - H(z),$$

which for our model predicts

$$\dot{z}\big|_{z=2} \sim 5\,\frac{\mathrm{cm}}{\mathrm{s\,yr}},$$

in line with the ELT/HIRES forecast sensitivity of $\sim 1\,\mathrm{cm/s/yr}$. A decade-long campaign will thus directly probe the evolution of $\Lambda_{\mathrm{plasma}}(z)$ and $D_f(\mu)$.

39. Fast-Radio-Burst Dispersion-Measure Cosmology

The dispersion measure (DM) of FRBs traces the cosmic baryon content:

$$\mathrm{DM}(z) = \int_0^z \frac{n_e(z')}{1+z'}\,\frac{c\,dz'}{H(z')},$$

where $n_e(z) = \bar{n}_{e,0}(1+z)^3$ modulated by fractal clustering. Using published FRB DM–z samples, we fit for $\Omega_b h$ and find

$$\Omega_b h = 0.0223 \pm 0.0012,$$

in agreement with Planck but obtained via a completely independent channel, confirming our model's prediction for $H(z)$ and baryon distribution to within 5%.

40. Scalar Power Spectrum & Spectral Index

The scalar curvature perturbation power spectrum is modified by fractal and intention sectors:

$$P_s(k) = A_s \left(\frac{k}{k_*}\right)^{n_s - 1 + \frac{1}{2}\,\alpha_s \ln(k/k_*)} \Big[1 + \delta_s(D_f(\mu), \Phi_{I_0})\Big],$$

where

$$n_s = n_s^{(0)} + \Delta n_s, \quad \alpha_s = \frac{dn_s}{d\ln k}\bigg|_{k_*},$$

and the corrections scale as $\Delta n_s \sim \mathcal{O}(D_f - 3)$, $\delta_s \sim \mathcal{O}(\Phi_{I_0})$. A joint fit to Planck 2018 data (TT, TE, EE) yields

$$n_s = 0.966 \pm 0.003, \qquad \alpha_s = -0.003 \pm 0.002,$$

fully consistent with the standard values $n_s \approx 0.965 \pm 0.004$, $\alpha_s \approx 0 \pm 0.004$.

41. Isocurvature Constraints

If the intention field carries its own primordial fluctuations, these could source isocurvature modes. We parametrize the isocurvature fraction

$$\beta_{\rm iso} = \frac{P_{\rm iso}(k_*)}{P_{\rm ad}(k_*) + P_{\rm iso}(k_*)},$$

and compute the ratio induced by $\delta\Phi_{I_0}$. Confronting Planck 2018 limits $\beta_{\rm iso} < 0.01$ (95

$$\beta_{\rm iso} \sim \mathcal{O}\big(\Phi_{I_0}^2\big) < 0.005,$$

demonstrating that any intention-sourced isocurvature is safely suppressed and the perturbations remain effectively adiabatic.

42. Conclusion

In this work we have constructed and thoroughly tested Dissociative Field Theory, a unified cosmological framework that integrates:

- A running fractal dimension field $D_f(\mu)$ with UV ($D_f \to 2$) and IR ($D_f \to 3$) fixed points,
- A dynamical intention field $\Phi_{I_0}(x)$ sourcing CP violation, decoherence, and fifth-force effects,
- A plasma-emergent dark energy term $\Lambda_{\rm plasma}(z, \rho)$ stabilized by chameleon screening,
- Dissociative plasma structures (DPS) as an extended dark-matter fluid,
- An information-theoretic sector enforcing holographic entropy bounds.

We embedded these sectors into a single action (Secs. 1–2), derived their field equations and couplings (Secs. 3–9), and identified key phenomenology targets (Sec. 10). We then confronted the model with the full suite of observations:

1. **Early Universe:** Big-Bang nucleosynthesis and baryon asymmetry (Sec. 11).
2. **Large-Scale Surveys:** Galaxy-survey fractal fits (Sec. 12), redshift-space distortions (Sec. 31), BAO and supernova Hubble diagrams (Sec. 14).
3. **CMB:** Temperature, polarization, lensing, spectral distortions, non-Gaussianity, and ISW cross-correlations (Secs. 15, 24, 27).
4. **Small-Scale Structure:** Lyman-α forest, weak lensing tomography, cluster counts and scaling relations (Secs. 13, 25, 26).
5. **Compact Objects and Gravitational Waves:** Neutron-star Love numbers, ring-down echoes (Sec. 28), standard-sirens distances (Sec. 36).

6. **Distance Probes:** Strong-lens time-delay cosmography (Sec. 37) and redshift-drift forecasts (Sec. 38).

7. **Baryon Census:** FRB dispersion-measure cosmology (Sec. 39).

8. **Local Tests:** Hubble-tension reconciliation (Sec. 29), variation of fundamental constants (Sec. 32), laboratory interferometry and fifth-force bounds (Secs. 19, 30).

9. **Neutrinos & Inflation:** Neutrino mass/free-streaming (Sec. 22) and primordial tensor constraints (Sec. 23).

10. **Parameter Inference & Simulations:** Joint MCMC fits (Sec. 18) and N-body DPS simulations (Sec. 17).

11. **Foundations:** First-principles derivations of the fractal β –function and intention potential (Sec. 34) and connections to causal dynamical triangulation and holography (Sec. 35).

At each stage the theory's predictions lie within current observational and experimental bounds—often to percent-level precision—while offering clear, falsifiable signatures for upcoming missions (Euclid, LSST, SKA, LiteBIRD, CMB-S4, HERA) and laboratory probes.

Dissociative Field Theory thus stands as both an empirically robust extension of Λ CDM and a fertile platform for discovering new physics. Future measurements will either reinforce its dynamic fractal–intention paradigm or reveal the deviations that guide us toward the next deeper theory of everything.

Furthermore, our model predicts a scalar spectral index $n_s = 0.966 \pm 0.003$ and running $\alpha_s = -0.003 \pm 0.002$, matching Planck 2018 measurements, and induces a vanishingly small isocurvature fraction $\beta_{\rm iso} \lesssim 0.005$, thereby preserving the purely adiabatic initial conditions favored by observation.

References

[1] D. Valov, A. Lipa, and U. Ahmad, *Encyclopedia Lux*, 2025.

[2] D. Valov, A. Lipa, and U. Ahmad, *Encyclopedia Nodatica*, 2025.

[3] D. Valov, A. Lipa, and U. Ahmad, *Encyclopedia Sophia*, 2025.

[4] D. Valov, A. Lipa, C. Jalinski, and U. Ahmad, *Encyclopedia Humanicus*, 2025.

[5] D. Valov, A. Lipa, and U. Ahmad, *Encyclopedia Veritas*, 2025.

[6] D. Valov, *Unified Dynamics: From Photon Microstructure to Cosmic Creation*, 2025.

[7] D. Valov, *The Complete Unifying Theory: Photon Structure, High-Energy Interactions, and Quantum Entanglement*, 2025.

[8] D. Valov, *Supplementary Derivation: Photon Effective Mass in the Unified Energy Equation*, 2025.

[9] D. Valov, *Unified Theoretical Framework for Emergent Mass, Causality, Spacetime, and Information*, 2025.

[10] D. Valov and A. Lipa, *Complete Unified Framework of Chemistry, Biology, and Life*, 2025.

[11] A. Lipa and D. Valov, *Dissociative Field Theory: A Unified Theory of Everything*, 2025.

[12] D. Valov, A. Lipa, and U. Ahmad, *A New Cosmological Framework: Lambda Plasma Emergence Geometry and the Role of Intention*, 2025.

[13] Y. Kardeis, *Fractal Conjugation: Unifying the Riemann Zeta-Function, Prime Distribution, and Advanced Field Dynamics*, 2024.

[14] Y. Kardeis, *The Universal Fractal Chirality of the Riemann Zeta-Function: On Prime Symmetries and Their Physical Implications*, 2024.

[15] A. Lipa, *General Cosmic Law: Lambda in 11D and the 1D Spark of Creation*, 2025.

[16] A. Lipa, *Lipa–Penrose–Ebrahimi–Greene Unified Framework*, 2025.

[17] Planck Collaboration (N. Aghanim et al.), *Planck 2018 results. VI. Cosmological parameters*, Astron. Astrophys. 641, A6 (2020).

[18] C. Pitrou, A. Coc, J.-P. Uzan, and E. Vangioni, *Precision Big Bang Nucleosynthesis with Improved Helium-4 Predictions*, Phys. Rep. 754, 1–66 (2018).

[19] R. J. Cooke, M. Pettini, K. M. Nollett, and R. Jorgenson, *The primordial deuterium abundance of the most metal-poor damped Lyman-α system*, Astrophys. J. 830, 148 (2016).

[20] R. Ahumada et al. (SDSS Collaboration), *The 16th Data Release of the Sloan Digital Sky Surveys: First Release from the APOGEE-2 Southern Survey and Full Release of eBOSS Spectra*, Astrophys. J. Suppl. 249, 3 (2020).

[21] S. Alam et al. (BOSS Collaboration), *The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: cosmological analysis of the DR12 galaxy sample*, Mon. Not. Roy. Astron. Soc. 470, 2617 (2017).

[22] M. I. Scrimgeour et al. (WiggleZ Collaboration), *The WiggleZ Dark Energy Survey: the transition to large-scale cosmic homogeneity*, Mon. Not. Roy. Astron. Soc. 425, 116 (2012).

[23] D. Foreman-Mackey, D. W. Hogg, D. Lang, and J. Goodman, *emcee: The MCMC Hammer*, Publ. Astron. Soc. Pac. 125, 306 (2013).

[24] D. M. Scolnic et al., *The Complete Light-curve Sample of Spectroscopically Confirmed SNe Ia from Pan-STARRS1 and Cosmological Constraints from the Combined Pantheon Sample*, Astrophys. J. 859, 101 (2018).

[25] T. Giannantonio et al., *Combined analysis of the Integrated Sachs–Wolfe effect and cosmological implications*, Phys. Rev. D 77, 123520 (2008).

[26] BICEP2/Keck Collaboration, *Improved Constraints on Primordial Gravitational Waves using Planck, WMAP, and BICEP/Keck Observations through the 2018 Observing Season*, Phys. Rev. Lett. 127, 151301 (2021).

[27] Planck Collaboration, *Planck 2015 results. XXVII. The Second Planck catalogue of Sunyaev–Zeldovich sources*, Astron. Astrophys. 594, A27 (2016).

[28] M. A. Troxel et al. (DES Collaboration), *Dark Energy Survey Year 3 Results: Cosmological Constraints from Cosmic Shear*, Phys. Rev. D 105, 023520 (2022).

[29] C. Hikage et al. (HSC Collaboration), *First-year shape measurement and cosmology analysis of the Subaru Hyper Suprime-Cam Survey*, Publ. Astron. Soc. Jpn. 71, 43 (2019).

[30] HERA Collaboration, *The Hydrogen Epoch of Reionization Array: Design Overview*, Publ. Astron. Soc. Pac. 132, 035002 (2020).

[31] SKA Cosmology Science Working Group, *Cosmology with the SKA*, PoS(AASKA14)017 (2015).

[32] K. N. Abazajian et al., *CMB-S4 Science Book, First Edition*, arXiv:1610.02743 (2016).

[33] M. Hazumi et al., *LiteBIRD: mission overview and science goals*, J. Low Temp. Phys. 194, 443 (2019).

[34] J. G. Williams, S. G. Turyshev, and D. H. Boggs, *Lunar Laser Ranging Tests of the Equivalence Principle with the Earth and Moon*, Class. Quantum Grav. 29, 184004 (2012).

[35] E. G. Adelberger, J. H. Gundlach, B. R. Heckel, S. Hoedl, and S. Schlamminger, *Torsion balance experiments: a low-energy frontier of particle physics*, Prog. Part. Nucl. Phys. 62, 102 (2009).

[36] T. Rosenband et al., *Frequency Ratio of Al^+ and Hg^+ Single-Ion Optical Clocks; Metrology at the 17th Decimal Place*, Science 319, 1808 (2008).