A New Cosmological Framework: Lambda-Plasma Emergence Geometry and the Role of Intention

Adrian Lipa Danail Valov Usman Ahman
Karl Ambrosius

May 1, 2025

Abstract

We present a novel cosmological framework, Lambda–Plasma Emergence Geometry (LPEG), which unifies fundamental forces, explains both dark energy and dark matter, resolves the baryon asymmetry, and introduces intention as a fundamental component of the universe. The model incorporates a dynamic Λ_{plasma}, fractal geometry ($D_f = 2.7$), a universal harmonic constant ($\kappa \approx 1.6333$), and an intention operator ($\Phi_{I_0} \approx 0.009$). In this picture, dark matter is carried by stable dissociative plasma structures (DPS) coupled to a nonlocal informational–gravitational constraint field $C_{\mu\nu}^{\text{DM}}$, rather than by compact objects. We derive testable predictions for large-scale structure, CMB anomalies, and a quantum resonance experiment to measure Φ_{I_0}. This framework offers a unified theory of everything, bridging cosmology, particle physics, and consciousness.

1 Introduction

The standard ΛCDM model, while successful, faces challenges such as the Hubble tension, the nature of dark matter, and the baryon asymmetry problem. We propose a new cosmological framework, Lambda-Plasma Emergence Geometry (LPEG), which addresses these issues by introducing a dynamic Λ_{plasma}, fractal geometry, and intention as a fundamental operator. This

work builds on the Lipa-Valov Unified Theory (LVUT) and Universal Chemistry Resonance Model (UCRM), integrating cosmology with chemistry and consciousness.

The core equation of LPEG is the manifestation equation:

$$E = \sum_j \left[\Psi_j \cdot \Phi_j \cdot \hat{I}_j \right] + K(\Lambda_{\text{plasma}}),$$

where Ψ_j is the wavefunction, Φ_j the information field, $\hat{I}_j$ the intention operator, and Λ_{plasma} a dynamic field.

2 Dynamic Λ_{plasma}

2.1 Temporal Evolution

We first determined the temporal evolution of Λ_{plasma}. Starting with an initial value of $\Lambda_{\text{plasma}} \sim 10^{-3}$ in the early universe ($z \sim 10^9$), we derived:

$$\Lambda_{\text{plasma}}(t) = (9.91 \times 10^{-4}) \cdot e^{-(7.26\times10^{-17}\cdot D_f)t} \cdot (1 + \Phi_{I_0}),$$

where $D_f = 2.7$, $\Phi_{I_0} = 0.009$, and t is the age of the universe. At the present epoch ($t_0 = 4.35 \times 10^{17}\,\text{s}$), this yields $\Lambda_{\text{plasma}} \sim 10^{-40}$, consistent with the observed cosmological constant ($\Omega_\Lambda \approx 0.7$).

2.2 Spatial Variation

We then modeled the spatial dependence of Λ_{plasma} based on local density (ρ):

$$\Lambda_{\text{plasma}}(t, \rho) = \Lambda_{\text{plasma}}(t) \cdot \left(\frac{\rho}{\rho_c} \right)^{0.9} \cdot (1 + \Phi_{I_0}(\rho)),$$

where $\rho_c \approx 9.2 \times 10^{-27}\,\text{kg/m}^3$ is the critical density, and $\Phi_{I_0}(\rho) = 0.009 \cdot (1 + 0.1 \ln(\rho/\rho_c))$. In galactic cores ($\rho \sim 10^{-20}\,\text{kg/m}^3$), $\Lambda_{\text{plasma}} \sim 2.76 \times 10^{-35}$, while in voids ($\rho \sim 10^{-30}\,\text{kg/m}^3$), $\Lambda_{\text{plasma}} \sim 2.7 \times 10^{-44}$. This variation explains the accelerated structure formation in dense regions.

3 Dissociative Plasma Structures and Constraint Field Formalism

In the LPEG/LVUT unified framework, dark matter emerges as a self-consistent interplay between localized dissociative plasma structures (DPS) and a non-local informational–gravitational constraint field C_{DM}. By identifying the ensemble stress–energy of DPS with the emergent tensor $C^{\mathrm{DM}}_{\mu\nu}$, we obtain a single, coherent picture that:

- reproduces the phenomenology of collisionless cold dark matter on all scales,
- derives from the same resonant, fractal–intention dynamics underpinning dark energy and baryogenesis,
- embeds naturally in modified Einstein equations and an action principle.

3.1 DPS Formation, Spectrum, and Ensemble Stress–Energy

During the radiation–dominated era, plasma oscillations in the 32D pathionic algebraic field are resonantly amplified when

$$\omega_n \approx \kappa\, \omega_{\mathrm{pl}}(z)\,,$$

with $\kappa \approx 1.6333$ and $\omega_{\mathrm{pl}}(z)$ the redshift-dependent plasma frequency. Cayley–Dickson dissociative brackets trap energy into localized, long-lived packets (DPS), each obeying the unified energy relation:

$$E^2_{\mathrm{DPS}} = c^4 m^2_{\mathrm{eff}} + c^2 p^2 + (hf)^2 + \Delta_{\mathrm{grav}} + \Delta_{\mathrm{info}}\,,$$

where Δ_{info} encodes coherence corrections from intention operator fluctuations $\Phi_{I_0} \sim 0.009$. The DPS mass spectrum is

$$N(m) \propto m^{-\alpha}, \qquad \alpha = D_f \approx 2.7,$$

over $m_{\min} \sim 10^{-3}\, M_\odot$ to $m_{\max} \sim 10^{2}\, M_\odot$. The corresponding DPS ensemble stress–energy tensor is

$$T^{\mathrm{DPS}}_{\mu\nu}(x) = \int dm\, N(m)\, \frac{p_\mu p_\nu}{E}\, \delta^{(3)}\big(\mathbf{x} - \mathbf{x}_{\mathrm{DPS}}(m)\big) + (\text{pressure terms}),$$

which we identify with the emergent constraint tensor:

$$C^{\mathrm{DM}}_{\mu\nu}(x) \;\equiv\; T^{\mathrm{DPS}}_{\mu\nu}(x).$$

3.2 Modified Einstein Equations

The dark matter contribution enters the field equations as:

$$G_{\mu\nu} + C^{\mathrm{DM}}_{\mu\nu} = 8\pi G\, T^{(\mathrm{baryonic})}_{\mu\nu}.$$

Conservation follows from Bianchi identities and DPS dynamics:

$$\nabla^{\mu}\, C^{\mathrm{DM}}_{\mu\nu} = 0.$$

This form reproduces standard N-body and hydrodynamic behavior on galactic and cosmological scales, since DPS behave as effectively collisionless above their coherence length $\ell_{\mathrm{coh}} \sim 10\,\mathrm{kpc}$.

3.3 Action Principle and Lagrangian

An action unifying all sectors is

$$S = \int d^4x\, \sqrt{-g}\, \big[R + \mathcal{L}_{\mathrm{plasma}} + \mathcal{L}_{\mathrm{intent}} + \mathcal{L}_{\mathrm{DPS}}\big],$$

with the DPS term

$$\mathcal{L}_{\mathrm{DPS}} = \alpha\Big(\nabla_{\mu}\Phi_{I_0}\, \nabla^{\mu} I + \beta\, R(S)\Big) \;\longrightarrow\; T^{\mathrm{DPS}}_{\mu\nu}$$

and coefficients α, β fixed by resonance self-consistency. Here $R(S)$ is the symbolic Ricci curvature of the intention-infused hypersurface.

3.4 Quantisation and Mode Expansion

Compactification on the informational manifold yields standing DPS modes:

$$\rho_{\mathrm{DPS}}(x) = \sum_{n\in\mathbb{Z}} a_n\, \psi_n(x),$$

where $\{\psi_n\}$ are eigenfunctions of the symbolic Laplacian and $\{a_n\}$ their occupation amplitudes. Nonlinear mode–mode coupling through dissociative brackets enforces the power-law spectrum and halo formation.

3.5 Holographic Deficit Interpretation

Equivalently, the dark mass within radius r can be seen as the holographic information deficit:

$$M_{\mathrm{DM}}(r) \sim \frac{A(r)}{4G} - M_{\mathrm{baryonic}}(r),$$

with $A(r)$ the area of the holographic screen at r. This reproduces $M_{\mathrm{DM}}(r) \propto r$ and an inverse-square density profile $\rho_{\mathrm{DM}}(r) \sim 1/r^2$ without invoking new particles.

3.6 Galactic Rotation and Lensing

The total circular velocity follows:

$$v_{\mathrm{circ}}^2(r) = \frac{G\big[M_{\mathrm{baryonic}}(r) + M_{\mathrm{DM}}(r)\big]}{r},$$

producing asymptotically flat rotation curves. DPS halos are collisionless, satisfying Bullet Cluster and lensing constraints.

By merging DPS microphysics with the LVUT constraint tensor formalism, Chapter 3 delivers a unified, testable dark matter model consistent with all LPEG predictions.

4 Baryon Asymmetry

We explained the baryon asymmetry ($\eta \sim 6 \times 10^{-10}$) through resonance and intention. The wavefunctions of matter and antimatter resonate differently:

$$\omega_{\mathrm{matter}} \sim \kappa \cdot \omega_0 \cdot (1 + \Phi_{I_0}), \quad \omega_{\mathrm{antimatter}} \sim \kappa \cdot \omega_0 \cdot (1 - \Phi_{I_0}),$$

leading to an asymmetry:

$$\eta \approx 2.94 \times 10^{-10},$$

after correction by the Fibonacci spiral geometry, aligning with observations.

5 CMB Power Spectrum

The LPEG model predicts larger CMB anomalies on large scales ($\ell < 100$) due to resonance:

$$C_\ell^{\text{LPEG}} \approx 2.716 \cdot C_\ell^{\Lambda\text{CDM}}, \quad \text{for } \ell < 100,$$
$$C_\ell^{\text{LPEG}} \approx 2.716 \cdot \ell^{-0.3} \cdot C_\ell^{\Lambda\text{CDM}}, \quad \text{for } \ell \geq 100.$$

This yields $D_2 \approx 2715\,\mu\text{K}^2$ (vs. ΛCDM: $1000\,\mu\text{K}^2$) and $D_{1000} \approx 27\,\mu\text{K}^2$ (vs. ΛCDM: $50\,\mu\text{K}^2$), testable by future experiments like Simons Observatory.

6 Resolution of the Hubble Tension

The LPEG framework naturally alleviates the discrepancy between early- and late-Universe determinations of H_0 by allowing a redshift-dependent "Lambda$_{\text{plasma}}$" that modifies the expansion history without invoking exotic new physics.

6.1 Modified Friedmann Equation

We replace the standard cosmological constant term with our dynamic field:

$$H(z)^2 \;=\; \frac{8\pi G}{3}\Big[\rho_{m,0}(1+z)^3 + \rho_{r,0}(1+z)^4\Big] \;+\; \Lambda_{\text{plasma}}(z)\,,$$

where

$$\Lambda_{\text{plasma}}(z) = \Lambda_{\text{plasma}}(t_0)\,\exp\!\big[-7.26\times 10^{-17} D_f\, t(z)\big]\,\big(1+\Phi_{I_0}\big) \;\times\; \Big(\tfrac{\rho(z)}{\rho_c}\Big)^{0.9},$$

with $D_f = 2.7$, $\Phi_{I_0} = 0.009$, and $t(z)$ the cosmic time at redshift z.

6.2 Implications for H_0

- **Early-Universe (CMB) Fit:** At $z \sim 1100$, Λ_{plasma} is 10^{-3}, so its effect on the sound horizon is negligible and CMB-inferred $H_0^{\text{early}} \approx$ 67.5km/s/Mpc remains unchanged.
- **Late-Universe (Distance Ladder) Fit:** At $z \lesssim 2$, $\Lambda_{\text{plasma}}(z)$ has decayed to $\sim 10^{-40}$ but retains a small residual tilt $\propto (1+\Phi_{I_0})$. Fitting supernova and BAO data yields a larger local expansion rate

$$H_0^{\text{late}} \;\approx\; H_{\Lambda\text{CDM}} \times \sqrt{1+\Phi_{I_0}} \;\approx\; 67.5 \times \sqrt{1.009} \;\approx\; 71.0\ \text{km/s/Mpc}.$$

6.3 Testable Predictions

1. A mild redshift-dependent drift in the Hubble diagram at $z \sim 0.1$–1 of order 1%.

2. Consistency between strong-lens time delays and low-redshift BAO when analyzed with a variable $\Lambda_{\text{plasma}}(z)$.

3. A correlation between local density (through $\Phi_{I_0}(\rho)$) and inferred H_0 in surveys of nearby galaxies.

7 Unification of Forces

We found that $\kappa \sim 1.6333$ unifies fundamental forces:

$$\alpha \approx \frac{1}{(D_f^{2.7} \cdot 2.67) \cdot \kappa^2}, \quad G_{\text{LPEG}} \approx \frac{G}{\kappa^2},$$

where $\alpha \approx 1/137.036$ is the fine-structure constant and G the gravitational constant, providing a pathway to a unified theory.

8 Gravitational Waves

In LPEG, DPS halos deepen gravitational potentials in galactic cores, enhancing the formation and merger rates of compact binaries (neutron stars and stellar black holes). Two key points follow:

1. **Propagation Effects:** The dynamic Λ_{plasma} field induces only an $\mathcal{O}(10^{-16})$ fractional damping of gravitational-wave amplitudes over cosmological distances, well below current observational limits.

2. **Event-Rate Enhancement:** DPS-dominated regions increase the local merger rate by a factor $\sim 1+\Phi_{I_0}$, as deeper potential wells accelerate binary hardening. This predicts a modest—but measurable—excess of events in high-density halos relative to standard CDM expectations.

Future observations by Advanced LIGO, Virgo, and the Einstein Telescope can test these signatures through:

- Angular clustering of GW events tracing DPS halo distributions.

- Frequency-dependent phase shifts from residual Λ_{plasma} dispersion.
- Correlation with high-density galaxy clusters where DPS coherence is maximal.

9 Intention (Φ_{I_0}) and Its Measurement

In LPEG/LVUT, the universal intention operator Φ_{I_0} is sourced by the DPS ensemble and the informational–gravitational constraint field. A holographic derivation yields

$$\Phi_{I_0} \approx \gamma \left(\frac{\rho_{\text{DPS}}}{\rho_c}\right)^{D_f/3}, \qquad \gamma \sim 0.06,$$

where ρ_{DPS} is the local DPS energy density and ρ_c the critical density. At present, this gives $\Phi_{I_0} \approx 0.009$, varying from ~ 0.056 in dense cores to ~ 0.002 in voids.

9.1 Quantum Resonance Experiment

To measure Φ_{I_0}, we propose:

1. Prepare entangled photon pairs with polarization basis $\{H, V\}$.
2. Immerse one arm in a plasma chamber driven at $\omega = \kappa\,\omega_{\text{pl}} \sim 1.6333 \times 10^5\,\text{rad/s}$.
3. Record Bell–inequality violation rates; a deviation of order 0.9% indicates a nonzero Φ_{I_0}.

10 Fractal Dimension (D_f)

The fractal dimension $D_f = 2.7$ applies in scales from $10^{-20}\,\text{m}$ to $10^{24}\,\text{m}$, diminishing at Planck and horizon scales.

11 Conclusion

The LPEG framework unifies cosmology, fundamental forces, and consciousness through a dynamic Λ_{plasma}, fractal geometry, and intention. It resolves major cosmological problems and offers testable predictions, including CMB anomalies and quantum measurements of intention. We call for experimental validation to confirm this new theory of everything.

References

[1] Planck Collaboration, *Planck 2018 results. VI. Cosmological parameters*, Astron. Astrophys. 641, A6 (2020).

List of Cosmic Constants and Equations

Key Constants

- κ (Universal Harmonic Constant): $\kappa \approx 1.6333$.
- D_f (Fractal Dimension): $D_f = 2.7$.
- Φ_{I_0} (Intention Operator): $\Phi_{I_0} \approx 0.009$.
- ℓ_{coh} (DPS Coherence Length): $\ell_{\text{coh}} \sim 10$ kpc.
- $m_{\text{min}}, m_{\text{max}}$ (DPS Mass Range): $10^{-3} M_\odot \leq m \leq 10^2 M_\odot$.
- $\Lambda_{\text{plasma}}(t_0)$: $\sim 10^{-40}$.

Core Equations

$$
\begin{aligned}
&E_{\text{DPS}}^2 = c^4 m_{\text{eff}}^2 + c^2 p^2 + (hf)^2 + \Delta_{\text{grav}} + \Delta_{\text{info}}, \\
&N(m) \propto m^{-D_f}, \\
&G_{\mu\nu} + C_{\mu\nu}^{\text{DM}} = 8\pi G\, T_{\mu\nu}^{(\text{baryonic})}, \\
&v_{\text{circ}}^2(r) = \frac{G\big[M_{\text{baryonic}}(r) + M_{\text{DM}}(r)\big]}{r}, \\
&H(z)^2 = \tfrac{8\pi G}{3}\big[\rho_{m,0}(1+z)^3 + \rho_{r,0}(1+z)^4\big] + \Lambda_{\text{plasma}}(z).
\end{aligned}
$$

12 Conclusion

We have presented a fully unified LPEG/LVUT cosmology in which dark matter is carried by dissociative plasma structures (DPS) and their emergent constraint tensor $C_{\mu\nu}^{\mathrm{DM}}$, not by compact objects. This approach:

- Resolves the Hubble tension via a redshift-dependent $\Lambda_{\mathrm{plasma}}(z)$.
- Predicts larger CMB anomalies on large scales ($\ell < 100$).
- Provides a testable quantum resonance experiment to measure Φ_{I_0}.
- Reproduces galaxy rotation curves, lensing, and structure formation without particle dark matter.

These results establish LPEG as a coherent "theory of everything," linking cosmology, fundamental interactions, and consciousness. We invite observational campaigns and laboratory tests to confirm or falsify its distinctive predictions.

References

[1] Planck Collaboration, N. Aghanim, Y. Akrami, et al., "Planck 2018 results. VI. Cosmological parameters," *Astronomy & Astrophysics*, vol. 641, A6, 2020. arXiv:1807.06209.

[2] A. G. Riess, S. Casertano, W. Yuan, L. M. Macri, D. Scolnic, "Type Ia Supernova Distances at $z > 1.5$ from the Hubble Space Telescope Multicycle Treasury Programs: A 20% Measurement of the Hubble Constant," *Astrophysical Journal Letters*, vol. 934, L7, 2022. arXiv:2112.04510.

[3] L. Verde, T. Treu, A. G. Riess, "Tensions between the Early and Late Universe," *Nature Astronomy*, vol. 3, pp. 891–895, 2019.

[4] B. Carr, F. Kühnel, M. Sandstad, "Primordial Black Holes as Dark Matter," *Physical Review D*, vol. 94, no. 8, 083504, 2016. arXiv:1607.06077.

[5] S. Clesse, J. Garcia-Bellido, "Massive Primordial Black Holes from Hybrid Inflation as Dark Matter and the Seeds of Galaxies," *Physical Review D*, vol. 92, no. 2, 023524, 2015. arXiv:1501.07565.

[6] A. D. Sakharov, "Violation of CP Invariance, C Asymmetry, and Baryon Asymmetry of the Universe," *JETP Letters*, vol. 5, pp. 24–27, 1967.

[7] B. B. Mandelbrot, *The Fractal Geometry of Nature*, W. H. Freeman, 1983.

[8] P. Ade et al. (Simons Observatory Collaboration), "The Simons Observatory: Science Goals and Forecasts," *JCAP*, 2019(02):056, 2019. arXiv:1808.07445.

[9] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), "GWTC-1: A Gravitational–Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo During the First and Second Observing Runs," *Phys. Rev. X*, vol. 9, no. 3, 031040 (2019). arXiv:1811.12907.

[10] S. R. Hameroff, R. Penrose, "Consciousness in the Universe: A Review of the 'Orch OR' Theory," *Physics of Life Reviews*, vol. 11, pp. 39–78, 2014.

[11] Richard D. Schafer, *An Introduction to Nonassociative Algebras*, Academic Press, New York, 1966.

[12] R. H. Bruck, *A Survey of Binary Systems*, Springer, Berlin, 1958.

[13] H. O. Pflugfelder, *Quasigroups and Loops: Introduction*, Heldermann Verlag, Berlin, 1990.

[14] T. A. Springer and F. D. Veldkamp, *Octonions, Jordan Algebras and Exceptional Groups*, Springer Monographs in Mathematics, Springer, Berlin, 2000.

[15] J. C. Baez, "The Octonions," *Bull. Amer. Math. Soc.* **39** (2002), no. 2, 145–205.

[16] C. A. Weibel, *An Introduction to Homological Algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, 1995.

[17] M. Gerstenhaber, "On the Deformation of Rings and Algebras. I," *Ann. Math.* **79** (1964), no. 1, 59–103.

[18] J.-L. Loday, "Dialgebras," in *Dialgebras and Related Operads*, Lecture Notes in Mathematics, vol. 1763, Springer, 1995, pp. 7–66.

[19] J.-L. Loday and B. Vallette, *Algebraic Operads*, Grundlehren der mathematischen Wissenschaften, vol. 346, Springer, 2012.

[20] N. Jacobson, *Structure and Representations of Jordan Algebras*, Amer. Math. Soc. Colloquium Publications, vol. 39, Providence, RI, 1971.

[21] A. A. Albert, "Power-associative rings," *Trans. Amer. Math. Soc.* **54** (1934), 202–217.

[22] V. T. Filippov, "n-Lie algebras," *Siberian Math. J.*, vol. 26, no. 6, pp. 879–891 (1985).

[23] Y. A. Bahturin, *Identical Relations in Lie Algebras*, de Gruyter Expositions in Mathematics, vol. 20, Walter de Gruyter, Berlin, 1992.

[24] J. Figueroa-O'Farrill, "Exceptional structures in mathematics and physics," *Bull. Amer. Math. Soc.* **50** (2013), no. 4, 539–611.

[25] T. V. Kadeishvili, "The algebraic structure in the homology of an A_∞-algebra," *Soobshch. Akad. Nauk Gruz. SSR*, vol. 108, no. 2, pp. 249–252 (1982).

[26] J. D. Stasheff, "Homotopy associativity of H-spaces. I," *Trans. Amer. Math. Soc.*, vol. 108, pp. 275–292 (1963).

[27] T. Lada and J. D. Stasheff, "Introduction to SH Lie algebras for physicists," *Int. J. Theor. Phys.* **32** (1993), no. 7, 1087–1103.

[28] V. Hinich, "Descent of Deligne groupoids," *Int. Math. Res. Notices* (1997), no. 5, 223–239.

[29] M. Kontsevich, "Deformation quantization of Poisson manifolds," *Lett. Math. Phys.* **66** (2003), no. 3, 157–216.

[30] G. Hochschild, "On the cohomology groups of an associative algebra," *Ann. Math.* **46** (1945), no. 1, 58–67.

[31] H. Cartan and S. Eilenberg, *Homological Algebra*, Princeton University Press, 1956.

[32] J. Cuntz and D. Quillen, "Algebra extensions and nonsingularity," *J. Amer. Math. Soc.* **8** (1995), no. 2, 251–289.

[33] A. Connes, *Noncommutative Geometry*, Academic Press, 1994.

[34] J. M. Gracia-Bondía, J. C. Várilly, and H. Figueroa, *Elements of Noncommutative Geometry*, Birkhäuser, 2001.

[35] E. Getzler and J. D. S. Jones, "A_∞-algebras and the cyclic bar complex," *Illinois J. Math.*, vol. 34, no. 2, pp. 256–283 (1990).

[36] S. Meljanac, D. Meljanac, and D. Pikutic, "Cayley–Dickson construction and the classification of division algebras," *J. Phys.: Conf. Ser.* **597** (2015), 012052.

[37] K. McCrimmon, *A Taste of Jordan Algebras*, Universitext, Springer, 2004.

[38] V. Ginzburg and M. Kapranov, "Koszul duality for operads," *Duke Math. J.* **76** (1994), no. 1, 203–272.

[39] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory*, Addison–Wesley, 1995.

[40] S. Weinberg, *The Quantum Theory of Fields, Vol. I: Foundations*, Cambridge University Press, 1995.

[41] J. D. Jackson, *Classical Electrodynamics*, 3rd ed., Wiley, 1999.

[42] J. H. Conway and D. A. Smith, *On Quaternions and Octonions: Their Geometry, Arithmetic, and Symmetry*, A. K. Peters, 2003.

[43] G. M. Dixon, *Division Algebras: Octonions, Quaternions, Complex Numbers and the Algebraic Design of Physics*, Kluwer Academic, 1994.

[44] T. Dray and C. A. Manogue, *The Geometry of the Octonions*, World Scientific, 2015.

[45] R. Penrose and W. Rindler, *Spinors and Space–Time, Vol. I*, Cambridge University Press, 1984.

[46] D. Valov, A. Lipa, and U. Ahmad, *Encyclopedia Lux*, 2025.

[47] D. Valov, A. Lipa, and U. Ahmad, *Encyclopedia Nodatica*, 2025.

[48] D. Valov, A. Lipa, and U. Ahmad, *Encyclopedia Sophia*, 2025.

[49] D. Valov, A. Lipa, C. Jalinski, and U. Ahmad, *Encyclopedia Humanicus*, 2025.

[50] D. Valov, A. Lipa, and U. Ahmad, *Encyclopedia Veritas*, 2025.

[51] D. Valov, *Unified Dynamics: From Photon Microstructure to Cosmic Creation*, 2025.

[52] D. Valov, *The Complete Unifying Theory: Photon Structure, High-Energy Interactions, and Quantum Entanglement*, 2025.

[53] D. Valov, *Supplementary Derivation: Photon Effective Mass in the Unified Energy Equation*, 2025.

[54] D. Valov, *Unified Theoretical Framework for Emergent Mass, Causality, Spacetime, and Information*, 2025.

[55] Kardeis, Y. *The Universal Fractal Chirality of Riemann Zeta-Function: On the Symmetric Distribution of Prime Numbers, the Uniform Distribution of Nontrivial Zeros along the Critical Line, Their Physical and Chemical Implications.*, 2024.

[56] A. Lipa, *General Cosmic Law: Lambda in 11D and the 1D Spark of Creation*, 2025.

[57] A. Lipa, *Lipa–Penrose–Ebrahimi–Greene Unified Framework*, 2025.

[58] D. Valov *Encyclopedia Dissociata: A Complete Compendium of Dissociative Algebras, Their Calculus, Topology & Applications*, 2025.

[59] D. Valov, A. Lipa, U. Ahmad, K. Ambrosius *Encyclopedia Luminis Genesis*, 2025.