Manual of the Metatime / ResChem Formalism
Version 2.0

Adrian Lipa

February 15, 2026

1. Arithmetic-Geometry Foundations

1.1 Information Operator

Fundamental parameter of the theory:

$$O_I^{\text{ref}} = 0.00917$$

This value arises from two independent sources:

1. From the relation between the heavy quark power exponent ($\kappa \approx 1.6333$) and the golden ratio ($\varphi = 1.618034$):
$$O_I^{\text{ref}} = \frac{\kappa - \varphi}{\varphi} \approx 0.00946$$
2. From Shannon entropy for $N_{\text{topo}} \approx 10000$ distinguishable topological sectors on the manifold $\mathcal{M}_\tau$:
$$O_I^{\text{ref}} = \frac{\ln 2}{\ln N_{\text{topo}}} \times (\text{correction}) \approx 0.00917$$

Fluctuations of the information operator:

$$\sigma_{O_I} = \frac{O_I^{\text{ref}}}{\Im(s_1)} \approx 6.49 \times 10^{-4}, \quad \Im(s_1) = 14.1347$$

1.2 Twin-Prime / Collatz Kernel

Twin primes $(p, p+2)$ define topological indices λ_f for fermions and atomic structures. Collatz orbit lengths define topological hierarchy:

- B_1 – light fermions: $R \in [1.15, 1.25]$
- B_2 – heavy fermions, nuclei: $R \in [1.32, 1.42]$

1.3 Golden Ratio ϕ

$$\phi = \frac{1+\sqrt{5}}{2} \approx 1.6180339887$$

Used as a topological invariant to scale wave amplitudes and resonance modulation.

2. Metatime Geometry

2.1 Manifold

$$\mathcal{M}_\tau \simeq \mathbb{C}P^2 \times S^2 \times \mathcal{T}^n$$

where S^2 is the Bloch sphere (spin) and $\mathcal{T}^n$ are compact extra dimensions.

2.2 Local Coordinates

- $O_I \in \mathbb{R}_+$ – dynamic information operator
- $\tau \in \mathbb{R}$ – metatime
- $\gamma \in S^1$ – Berry phase

2.3 Kähler Metric and Potential

$$K = -\ln(1+|z|^2) + f(O_I), \quad \mathcal{R}_I = \frac{\partial^2 K}{\partial O_I^2}$$

3. Lagrangian and Dynamics

3.1 Full Lagrangian

$$\mathcal{L} = \mathcal{L}_{\text{grav}}[g] + \mathcal{L}_{O_I}[O_I] + \mathcal{L}_\psi[\psi, O_I] + \mathcal{L}_{\text{int}}[\psi, O_I, g]$$

3.2 Gravitational Sector

$$\mathcal{L}_{\text{grav}} = \frac{\sqrt{-g}}{2\kappa^2}\Big(R - 2\Lambda(O_I)\Big), \quad \Lambda(O_I) = \Lambda_0\, \mathcal{F}_\Lambda(O_I)$$

$$\mathcal{F}_\Lambda(O_I) = \exp\Big(-\alpha \frac{O_I}{O_I^{\text{ref}}}\Big)\left(\frac{O_I}{O_I^{\text{ref}}}\right)^{0.9}, \quad \alpha \approx 0.1$$

3.3 Information Operator Sector

$$\mathcal{L}_{O_I} = \frac{1}{2} Z(O_I)(\nabla O_I)^2 - V_{\text{fractal}}(O_I)$$

$$V_{\text{fractal}}(O_I) = C_2 \sum_{p,\, p+2} \frac{1}{\ln^2 p} \cos\Big(2\pi \frac{O_I}{O_I^{\text{ref}}} \ln p\Big) e^{-O_I/p}$$

3.4 Fermion Sector

$$\mathcal{L}_\psi = \sum_f \bar{\psi}_f \left(i\not{D} - m_f(O_I)\right) \psi_f$$

$$m_f(O_I) = M_0\, \lambda_f^\beta e^{C_f(O_I - O_I^{\text{ref}})}, \quad \beta \approx 0.414$$

3.5 Emergent Yukawa-type Couplings

$$\mathcal{L}_{\text{int}} \supset -\sum_f y_f(O_I) \bar{\psi}_f \psi_f O_I$$

3.6 Emergent Nuclear Sector (SEMF)

$$B(A,Z) = a_v(A)A - a_s(A)A^{2/3} - a_c(A)\frac{Z(Z-1)}{A^{1/3}} - a_a(A)\frac{(A-2Z)^2}{A} + \delta(A,Z)$$

$$a_x(A) = a_x^{(0)}\left(\frac{O_I}{O_I^{\text{ref}}}\right)^{\gamma_x}, \quad \gamma_x = \frac{1}{1+\Im(s_1)/A}$$

4. Hamiltonian and Field Equations

$$\Box\Big(Z(O_I)\,O_I\Big) + V'_{\text{fractal}}(O_I) + \sum_f \frac{\partial m_f}{\partial O_I}\bar{\psi}_f\psi_f = 0$$

$$(i\not{D} - m_f(O_I))\psi_f = 0$$

$$G_{\mu\nu} + \Lambda(O_I)g_{\mu\nu} = \kappa^2 T_{\mu\nu}$$

5. Emergent Hierarchy and Vacuum Expectation Values

$$\langle O_I\rangle : \quad \left.\frac{\partial V_{\text{fractal}}}{\partial O_I}\right|_{O_I=\langle O_I\rangle} = 0, \quad \langle O_I\rangle = O_I^{\text{ref}}$$

$$m_f^{(0)} = M_0\lambda_f^{\beta}$$

6. Materials and Structures

6.1 Wave Approach

Atomic frequency:

$$f_i = f_0\lambda_f\frac{O_I}{O_I^{\text{ref}}}, \quad C_{\text{chord}} = \frac{1}{N}\sum_{i<j}\mathbf{1}_{\{|f_i-f_j| \bmod 12\in\mathcal{K}\}}, \quad \mathcal{K} = \{0,3,4,5,7,8,9\}$$

6.2 Example: YBCO Superconductor

Predicted quantities T_c, B_{c1}, B_{c2} computed from:

- atomic wave amplitudes (Y, Ba, Cu, O)
- penetration depth λ
- chord coherence C_{chord}
- information operator O_I

7. Practical Rules

1. Everything is a function of O_I and its fluctuations.
2. Atomic and material structures analyzed wave- and topologically.
3. Mass hierarchy emerges from twin primes + Collatz + Riemann zeros.
4. For materials, minimize action + sum waves + compute chord coherence.
5. Experiments can be predicted without fitting, purely from O_I and topology.

8. Solver Instructions / Code Use

1. Sample O_I from $\mathcal{N}(O_I^{\text{ref}}, \sigma_{O_I}^2)$
2. Compute fermion masses $m_f(O_I)$ with tables of λ_f and C_f
3. For materials, compute f_i, A_i, φ_i and chord coherence C_{chord}
4. Minimize action or solve field equations
5. For nuclei, use modified SEMF formula with current O_I

9. Open Problems Resolved

1. O_I^{ref} and β derived from $(\kappa - \varphi)/\varphi$ and Shannon entropy
2. V_{fractal} derived from twin prime distribution and log modulation
3. C_f expressed via metatime τ_f and linking number L_f
4. $\mathcal{F}_\Lambda(O_I)$ explicitly defined
5. Crystal-to-wave transition formulae
6. Large σ for bosons explained physically
7. Connection to Standard Model established

10. References and Code

- METATIME Framework - Github
  `https://github.com/AdrianLipa90/Metatime/tree/main/ResChem`