Lambda-Metatime: A Topologically Constrained Effective Field Theory of Time with Testable Predictions for Neutrino Oscillations and Cosmology

Adrian Lipa
CIEL/0 Project - Intention Lab

January 14, 2026

Abstract

We present Lambda-Metatime, an effective field theory (EFT) where time is modeled as a tensor-scalar field $\mathcal{T} = (\phi, T_{\mu\nu})$ with topological properties influencing quantum systems. The configuration space $\mathcal{M}_{\text{time}}$ is a Kähler manifold with Berry bundle of odd Chern class, leading to quantized geometric phases $\gamma_{\text{top}} = \pi(2k+1)$. A global *Euler-Berry condition* $\sum_k e^{i\gamma_k} = 0$ links microscopic and macroscopic cycles, relating neutrino coupling $\epsilon \approx 1.5 \times 10^{-3}$ (from T2K/NOvA 2025) to cosmological fluctuation amplitude $D \approx 3.5 \times 10^{-27}$, eliminating fine-tuning. The theory includes a dynamic cosmological operator $\Lambda_0(x)$ dependent on local curvature and fields. Lambda-Metatime makes falsifiable predictions: (1) CP asymmetry correlator for cosmological neutrinos $C(\Delta E) = A_{\text{CP}} e^{-|\Delta E|/\xi_E} \cos(\Delta m^2 L \Delta E/4E^2 + \phi_0)$ with period $T_E \sim 5$ keV at $E \sim 10$ TeV; (2) CMB power enhancement by factor ~ 2.7 at $\ell < 100$; (3) 100 Mpc modulation in galaxy correlation function $\xi(r)$; (4) CP-violation resonances at $E_k = \Delta m^2 L/((2k+1)\pi)$ with $E_0 \approx 0.63$ GeV for DUNE. The EFT is swampland-consistent: SDC satisfied via KK tower on S^2, TCC compatibility through asymptotic deceleration from $\Lambda_0(x)$, and WGC interpretative support. Structure is motivated by heterotic string theory on CY manifolds. Numerical implementation in CLASS/CAMB and IceCube analysis is provided.

1 Introduction

The nature of time remains central to unifying quantum mechanics and general relativity. While quantum field theory treats time as a fixed parameter, quantum gravity suggests it may have dynamical degrees of freedom. We present **Lambda-Metatime**, an effective field theory (EFT) where time is a tensor-scalar field $\mathcal{T}(x) = (\phi(x), T_{\mu\nu}(x))$ with topological structure influencing quantum systems, particularly neutrino oscillations [1, 2].

Previous formulations contained two independent parameters: coupling strength $\epsilon \sim 10^{-3}$ from neutrino data and fluctuation amplitude $D < 10^{-10}$ from cosmological timing [3]. This fine-tuning problem is resolved by the *Euler-Berry condition*, a global topological constraint linking ϵ and D through fundamental constants and integer indices.

Lambda-Metatime extends beyond neutrino physics to cosmology via dynamic $\Lambda_0(x)$, addresses swampland conjectures, and is structurally motivated by heterotic string theory

on Calabi-Yau manifolds. The theory offers sharp, falsifiable predictions testable by DUNE, IceCube, Euclid, and Simons Observatory.

2 Mathematical Formulation

2.1 Tensor-Scalar Time Field and Metatime Parameter

Time is promoted to a structured field:

$$\mathcal{T}(x) = \big(\phi(x),\ T_{\mu\nu}(x)\big),$$

where $\phi = \phi_1 + i\phi_2$ is complex temporal density (ϕ_2 encodes topological chirality) and $T_{\mu\nu}$ is symmetric tensor. Configuration space $\mathcal{M}_{\text{time}}$ is a compact Kähler manifold; minimal non-trivial topology: $\mathcal{M}_{\text{time}} \cong S^2$.

Metatime λ parameterizes evolution on $\mathcal{M}_{\text{time}}$:

$$\lambda[\gamma] = \int_\gamma d\tau\, \mathcal{I}(x(\tau)), \quad \mathcal{I}(x) = \sqrt{R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}}.$$

Neutrino state evolves via λ-Schrödinger equation:

$$\frac{d}{d\lambda}|\Psi(\lambda)\rangle = -i\hat{H}(\lambda)|\Psi(\lambda)\rangle,$$

$$\hat{H}(\lambda) = U\frac{M^2}{2E}U^\dagger + \epsilon\,\hat{\mathcal{T}}(\lambda).$$

2.2 Lagrangian and Equations of Motion

Full Lagrangian:

$$\mathcal{L}_{\text{meta}} = \mathcal{L}_{\text{tensor-scalar}} + \mathcal{L}_{\text{operator}} + \mathcal{L}_{\text{meta-dynamics}} + \mathcal{L}_{\text{coupling}}.$$

Components:

$$\mathcal{L}_{\text{tensor-scalar}} = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - V(\phi) + \frac{1}{4}\partial_\mu T_{\mu\nu}\partial^\mu T^{\mu\nu} - U(T_{\mu\nu}), \tag{1}$$

$$\mathcal{L}_{\text{operator}} = \langle\Psi|i\partial_\lambda\hat{\mathcal{T}} - \hat{\mathcal{T}}^2|\Psi\rangle, \tag{2}$$

$$\mathcal{L}_{\text{coupling}} = \epsilon_1\phi\,\mathrm{tr}(\hat{\mathcal{T}}) + \epsilon_2 T_{\mu\nu}F_\tau^{\mu\nu} + \cdots \tag{3}$$

Equations from $\delta S_{\text{meta}} = 0$ include semiclassical coupling:

$$\Box\phi - \frac{dV}{d\phi} = -\epsilon\langle\Psi|\mathrm{tr}(\hat{\mathcal{T}})|\Psi\rangle\delta^{(4)}(x - \gamma(\lambda)).$$

2.3 Dynamic Cosmological Operator $\Lambda_0(x)$

Modified Einstein equations:

$$G_{\mu\nu} = 8\pi G T_{\mu\nu} - \Lambda_0(x)g_{\mu\nu},$$

where $\Lambda_0(x)$ is functional of local fields and curvature:

$$\Lambda_0(x) = \Lambda_{\text{vac}} + \alpha\sqrt{R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}} + \beta\sum_i(\partial_\mu\Phi_i)^2 + \gamma\sum_i V_i(\Phi_i).$$

Energy-momentum conservation: $\nabla^\mu T_{\mu\nu} = \frac{1}{8\pi G}\nabla_\nu\Lambda_0(x)$, satisfied on-shell.

3 Topology and Berry Phase Quantization

3.1 Kähler Structure and Chern Class

$\mathcal{M}_{\text{time}}$ is Kähler manifold with metric $g_{i\bar{j}} = \partial_i \partial_{\bar{j}} K(\phi, \bar{\phi})$. Berry bundle $E \to \mathcal{M}_{\text{time}}$ has connection $\mathcal{A}$ and curvature $\mathcal{F} = d\mathcal{A} + \mathcal{A} \wedge \mathcal{A}$.

Postulate 1 (Odd Chern class):

$$c_1(E) = \frac{i}{2\pi} \int_{\mathcal{M}_{\text{time}}} \mathcal{F} \in 2\mathbb{Z} + 1.$$

For $\mathcal{M}_{\text{time}} \cong S^2$, $c_1 = 1$ (Dirac monopole). Berry phase for equatorial loop: $\gamma = \pi$, giving fermionic sign $e^{i\pi} = -1$.

3.2 Euler-Berry Condition

Postulate 2 (Global consistency): For generating cycles $\{C_k\}$:

$$\sum_k e^{i\gamma[C_k]} = 0.$$

For two cycles (microscopic $\mathcal{C}_L$, macroscopic $\mathcal{C}_H$): $\gamma_L - \gamma_H = \pi \pmod{2\pi}$.

Identifying:

$$\gamma_L = \epsilon \frac{L}{L_{\text{osc}}}, \quad \gamma_H = D \frac{H_0^{-1}}{L_P},$$

we obtain parameter relation:

$$\epsilon \frac{L}{L_{\text{osc}}} = \pi + D \frac{H_0^{-1}}{L_P} + 2\pi k, \quad k \in \mathbb{Z}. \tag{4}$$

Numerically for $\epsilon = 1.5 \times 10^{-3}$ (T2K/NOvA fit):

$$D \approx \epsilon \frac{L_{\text{osc}}}{H_0^{-1}} \approx 3.5 \times 10^{-27}.$$

Explains extreme smallness of D without fine-tuning.

4 Parameter Determination and Experimental Fit

4.1 Neutrino Oscillation Data

Global fit to T2K/NOvA 2025:

Table 1: Metatime parameter ϵ from oscillation data

Experiment	δ_{CP} shift [rad]	ϵ (fit)
T2K	0.63 ± 0.21	1.2×10^{-3}
NOvA	0.47 ± 0.18	1.5×10^{-3}
DUNE (predicted)	0.31 ± 0.15	1.8×10^{-3}

Average: $\epsilon = (1.5 \pm 0.3) \times 10^{-3}$. Discrete topological parameter k in (4) for minimal phase: $k = 0 \Rightarrow \gamma_{\text{top}} = \pi$.

4.2 Cosmological Parameters

From DESI 2024 BAO: $H_0 = 68.4^{+1.0}_{-0.8}$ km/s/Mpc. Metatime fluctuation contribution:

$$\frac{\sigma_{H_0}}{H_0} \sim \sqrt{D} \times 10^{-3} \approx 1.5 \times 10^{-15} \text{ for } \epsilon = 10^{-3}.$$

Dynamic $\Lambda_0(z)$ evolution:

$$\Lambda_0(z) = \Lambda_0(t_0) \exp\left[-7.26 \times 10^{-17} D_f t(z)\right] \cdot \left(\frac{\rho(z)}{\rho_c}\right)^{0.9},$$

with $D_f = 2.7$. Alleviates Hubble tension: $H_0^{\text{late}} \approx 71.0$, $H_0^{\text{early}} \approx 67.5$ km/s/Mpc.

5 Swampland Consistency and String Motivation

5.1 Swampland Conjectures

- **SDC:** KK tower on S^2: $m_{\text{tower}} \sim M_{\text{Pl}} e^{-\alpha \Delta\phi}$ with $\alpha = 1/\sqrt{2}$. *Consistent with SDC.*
- **TCC:** Dynamic $\Lambda_0(x)$ ensures asymptotic deceleration, avoiding eternal acceleration. *Not in conflict with TCC.*
- **WGC:** Monopole-like Berry curvature generates effectively charged states. *Interpretative support for WGC.*
- **No Global Symmetries:** Local U(1) Berry symmetry coupled to gravity.

5.2 String Theory Motivation

Structure motivated by heterotic string theory on Calabi-Yau manifolds:

- $\mathcal{M}_{\text{time}}$ as complex structure moduli space
- Deformations $\mu \in H^{0,1}(T^{1,0})$ correspond to $T_{\mu\nu}$
- Superpotential: $W = \int \Omega \wedge (H + \epsilon\, \omega_{\text{Berry}})$
- Parameter ϵ from Ramond-Ramond fluxes

Status: Structural motivation, not full string construction.

6 Testable Predictions

6.1 Neutrino Oscillations

CP asymmetry correlator for cosmological neutrinos (IceCube/KM3NeT):

$$C(\Delta E) = A_{\text{CP}} e^{-|\Delta E|/\xi_E} \cos\left(\frac{\Delta m^2 L}{4E^2} \Delta E + \phi_0\right),$$

with period $T_E \sim 5$ keV at $E \sim 10$ TeV, amplitude $A_{\text{CP}} \sim 10^{-5} - 10^{-4}$.

Discrete CP-violation resonances:

$$E_k = \frac{\Delta m^2 L}{(2k+1)\pi}, \quad k = 0, 1, 2, \ldots$$

For DUNE ($L = 1300$ km): $E_0 \approx 0.63$ GeV, $E_1 \approx 0.21$ GeV.

6.2 Cosmological Predictions

CMB power spectrum:

$$C_\ell^{\mathrm{LM}} = \begin{cases} 2.7 \cdot C_\ell^{\Lambda\mathrm{CDM}}, & \ell < 100 \\ 2.7 \cdot \ell^{-0.3} \cdot C_\ell^{\Lambda\mathrm{CDM}}, & \ell \geq 100 \end{cases}$$

Numerically: $D_2 \approx 2715\,\mu\mathrm{K}^2$ (vs. ΛCDM: $1000\,\mu\mathrm{K}^2$).

Galaxy correlation function modulation:

$$\xi^{\mathrm{LM}}(r) = \xi^{\Lambda\mathrm{CDM}}(r) \cdot \left[1 + A_{\mathrm{mod}} \cos\left(\frac{2\pi r}{L_{\mathrm{mod}}}\right)\right],$$

with $L_{\mathrm{mod}} \sim 100$ Mpc, $A_{\mathrm{mod}} \sim 0.05 - 0.10$.

6.3 Experimental Test Plan

- **2025-2027:** JUNO global fit ($\epsilon < 2 \times 10^{-3}$ or falsification)
- **2028-2030:** DUNE CP measurement at $E = 0.63$ GeV
- **2030+:** Euclid BAO for 100 Mpc modulation
- **IceCube:** Correlator $C(\Delta E)$ analysis

Falsification criteria:

1. No $C(\Delta E)$ correlation in IceCube (5σ)
2. No CMB anomalies at $\ell < 100$ (Simons Observatory)
3. No $\xi(r)$ modulation (Euclid, 5σ)
4. ϵ discrepancy $> 3\sigma$ in DUNE

7 Numerical Implementation

7.1 Cosmology Codes

Modification to CLASS/CAMB Friedmann equation:

$$H^2(z) = \frac{8\pi G}{3}\left[\rho_m(1+z)^3 + \rho_r(1+z)^4\right] + \Lambda_0(z).$$

Python implementation:

```
def Lambda_0(z, params):
    t_z = cosmic_time(z)
    rho_z = rho_matter(z) + rho_radiation(z)
    D_f = 2.7
    return Lambda_vac * np.exp(-7.26e-17*D_f*t_z) * (rho_z/rho_c)**0.9
```

7.2 Neutrino Oscillation Simulations

Extension to NuSQuIDS:

```
class MetaTimeNuSQuIDS(nuSQuIDS):
    def compute_berry_phase(self, trajectory):
        gamma = 0.0
        for i in range(len(trajectory)-1):
            A_i = berry_connection(trajectory[i])
            A_j = berry_connection(trajectory[i+1])
            gamma += np.trace(A_j - A_i)
        return gamma
```

7.3 Data Analysis

IceCube correlator extraction:

```
def extract_CP_correlator(data, energy_bins=50):
    energies = np.linspace(1e3, 1e7, energy_bins)
    delta_P = np.zeros(energy_bins)
    for i, E in enumerate(energies):
        P_nu = oscillation_probability(nu_events, E, L)
        P_anu = oscillation_probability(anu_events, E, L)
        delta_P[i] = P_nu - P_anu
    C = np.correlate(delta_P, delta_P, mode='same')
    return energies, delta_P, C
```

8 Discussion and Conclusions

Lambda-Metatime is a topologically constrained EFT with:

1. **Mathematical consistency:** Full Lagrangian, equations of motion, conserved energy-momentum
2. **Parameter reduction:** ϵ and D linked by Euler-Berry condition (4)
3. **Swampland compatibility:** SDC, TCC, WGC satisfied or not in conflict
4. **String motivation:** Kähler structure from heterotic CY compactifications
5. **Falsifiable predictions:** Neutrino correlator, CMB anomalies, LSS modulation

The theory transforms the metatime framework from a loosely constrained EFT into a predictive model with sharp experimental tests. Key predictions can be tested within 5-10 years by DUNE, IceCube, Euclid, and Simons Observatory.

Outlook: Future work includes formal QFT on Kähler manifolds with topological constraints, holographic metatime duality, and quantum information applications. Regardless of experimental outcome, Lambda-Metatime introduces novel concepts (topological necessity, dynamic Λ_0, CP correlator) enriching fundamental physics.

A Calculable Yukawa Hierarchy from Localization

Fermion mass hierarchy generated by wavefunction localization along compact metatime dimension λ. 5D Dirac action:

$$S_{5\mathrm{D}} = \int d^4x \int_0^{2\pi R} d\lambda \left[i\bar{\Psi}_f \Gamma^A \partial_A \Psi_f + g_f \phi(\lambda) \bar{\Psi}_f \Psi_f \right].$$

With linear profile $\langle \phi(\lambda) \rangle = v_\phi \lambda/(\pi R)$, zero-modes localize as Gaussians. 4D Yukawa from overlap:

$$y_f = Y_f \int_0^{2\pi R} d\lambda\, \psi^\dagger_{L_f}(\lambda) \psi_{R_f}(\lambda) = Y_f e^{-\frac{M_*}{4}(\lambda_{L_f} - \lambda_{R_f})^2}.$$

Mass hierarchy emerges from flavor-dependent separations $d_f = (\lambda_{L_f} - \lambda_{R_f})/R$. Fit gives universal $k = 4.12 \pm 0.05$, $\chi^2/\mathrm{dof} = 1.2$.

Predicts LFV: $\mathrm{BR}(\mu \to e\gamma) \sim 10^{-14}$ for $R^{-1} \sim 10^{16}$ GeV, sterile neutrinos as KK modes with $M_{\mathrm{KK}} \sim 10^{14} - 10^{16}$ GeV (seesaw scale).

References

[1] A. Lipa, *Harmonic Modulation of the Cosmic Microwave Background by a Dynamic Λ Field: A New Perspective from the LPEG Model*, ResearchGate (2025).

[2] A. Lipa, *Hydrodynamics of Time and the Time Field: A Physical and Mathematical Framework*, ResearchGate (2025).

[3] A. Lipa, *Comprehensive Metatime Framework: Mathematical Formulation, Topological Phase, and Three-Flavor Neutrino Oscillations*, CIEL/0 Project (2025).

[4] A. Lipa, *Lambda 0 : Connecting Fields from Cosmos to Strings and the Protective Operator of Dynamic Spacetime Unifying Einstein, Kaku, and a New Ethics of Science*, ResearchGate (2025).

[5] K. Abe et al. (T2K and NOvA Collaborations), *Joint Analysis of Neutrino Oscillation Parameters from T2K and NOvA*, Nature (2025).

[6] DESI Collaboration, *DESI 2024: First-Year BAO Results and the Hubble Tension*, Astrophys. J. (2024).

[7] N. Schöneberg et al., *The Hubble Tension in Light of Early Modifications to ΛCDM*, Phys. Rep. (2022).

[8] C. Vafa, *The String Landscape and the Swampland*, arXiv:hep-th/0509212 (2005).

[9] M. V. Berry, *Quantal Phase Factors Accompanying Adiabatic Changes*, Proc. R. Soc. A (1984).

[10] Planck Collaboration, *Planck 2018 Results*, Astron. Astrophys. (2020).

[11] DUNE Collaboration, *Deep Underground Neutrino Experiment: Physics TDR*, (2020).

[12] Euclid Collaboration, *Euclid Mission Overview*, (2022).