Metatime Framework and the Topological Fingerprint Σ: A Predictive Formalism for Particle Spectra, Neutrino Oscillations and Cosmology

Adrian Lipa[1]

[1]CIEL/0 Project – Intention Lab

January 19, 2026

Abstract

We present the Metatime framework: an effective, topologically constrained formalism in which the role traditionally played by physical time is replaced by a tensor–scalar "metatime" field evolving on a Kähler-type configuration manifold. Discrete closed cycles in the metatime manifold are assigned eigenvalues τ_i; global Berry-like phases and a small universal topological regulator (the Intention Operator I) define for each state a stable, topologically protected fingerprint

$$\Sigma_i = (\tau_i,\, \gamma_i,\, F_i).$$

A phenomenological mass operator maps Σ_i to observed particle masses,

$$m_i = k_f\, \tau_i^{\alpha_f}\, F_i,$$

and pairwise topological couplings ("white threads") mediate neutrino splittings and CP effects. We describe a generator for emergent τ_i based on a Twin-Prime Collatz prescription with -weighting, derive the Hamiltonian and action that couple metatime to Standard Model fields, propose a standing-wave / knot interpretation of hadron binding (replacing the perturbative gluon picture), and give explicit numerical examples and falsifiable predictions for neutrino splittings and cosmological modulation. Reproducibility notes and pseudocode are provided.

1 Introduction

Parameter proliferation in particle physics motivates selection principles that reduce free parameters and supply physical explanation for observed hierarchies. The Metatime framework proposes that a tensor–scalar metatime

field $T(x) = (\phi(x), T_{\mu\nu}(x))$ on a compact Kähler-type configuration manifold $\mathcal{M}_{\text{time}}$ supplies such a principle. Closed cycles $\Gamma \subset \mathcal{M}_{\text{time}}$ are assigned discrete eigenvalues τ; Berry phases γ around cycles obey a global Euler–Berry constraint which discretizes admissible families of cycles. Topological corrections are collected in multiplicative factors F_i derived from a small universal Intention operator $I \ll 1$. The formalism is constructed to be falsifiable (neutrino resonances, CMB modulation, correlation functions) and reproducible. This manuscript gathers the mathematical definitions, the emergent generator for τ_i, the mass mapping, the Hamiltonian and action, a standing-wave knot picture for hadrons, and numerical examples.

2 Core formalism

2.1 Metatime field and eigenvalues

Let $\mathcal{M}_{\text{time}}$ be a compact configuration manifold of Kähler type. The metatime field is

$$T(x) = (\phi(x), T_{\mu\nu}(x)).$$

Define a metatime operator $\hat{T}(\lambda)$ acting on a Hilbert space $\mathcal{H}$ with eigenstates $|\Psi_i\rangle$:

$$\hat{T}(\lambda)|\Psi_i\rangle = \tau_i|\Psi_i\rangle, \qquad \tau_i \in \mathbb{R}_+. \tag{1}$$

Each τ_i is associated with a closed cycle $\Gamma_i \subset \mathcal{M}_{\text{time}}$.

2.2 Berry phase and Euler–Berry constraint

Let $A(T)$ denote a Berry connection on the parameter bundle over $\mathcal{M}_{\text{time}}$. For a cycle Γ_i,

$$\gamma_i = \oint_{\Gamma_i} A(T) \cdot dT.$$

We impose the global *Euler–Berry* constraint on a generating set $\{C_k\}$:

$$\sum_k e^{i\gamma_{C_k}} = 0, \tag{2}$$

which in a two-cycle reduction gives $\gamma_1 - \gamma_2 \equiv \pi \pmod{2\pi}$. This condition discretizes admissible cycle families.

2.3 Topological fingerprint and Intention operator

For each physically realized state define the topological fingerprint

$$\Sigma_i \equiv (\tau_i, \gamma_i, F_i), \tag{3}$$

where F_i is a multiplicative topological correction encoding a universal regulator (the Intention operator I):

$$F_i = \exp(C_i I), \qquad I \ll 1, \tag{4}$$

with dimensionless coefficients C_i that parametrise local topological amplification or suppression.

3 Mass operator and phenomenological ansatz

Introduce an operator-valued mass map $\hat{M}(\tau)$. In the family-wise phenomenological ansatz:

$$m_i = k_f\,(\tau_i)^{\alpha_f}\,F_i, \tag{5}$$

where family $f \in \{\text{charged leptons, light quarks, heavy quarks, neutrinos}\}$ has constants (k_f, α_f) fitted to anchors. For three charged leptons one may instead use a polynomial in τ for m^2:

$$m_\ell^2 = c_0 + c_1\tau + c_2\tau^2,$$

which reproduces three anchors exactly.

4 Generator for emergent τ_i: Twin-Prime Collatz + ζ-weighting

To render τ_i emergent, define a generator inspired by Collatz dynamics but adapted for continuous/semi-continuous seeds and controlled by twin-prime seeds and a zeta-like preference for short cycles.

Seeds. For each physical family f choose a set of twin-prime seeds $\{p_j, p_j + 2\} \subset \mathbb{P}$. The representative seed $\bar{p}_j = \frac{1}{2}(p_j + p_j + 2) = p_j + 1$ sets a scale.

Collatz-like map (family f).

$$C_f(\tau) = \begin{cases} \tau/2, & \tau < \eta_f \bar{p}_j, \\ 3\tau + 1, & \tau \geq \eta_f \bar{p}_j, \end{cases}$$

with $\eta_f \gtrsim 1$ a family control parameter. Iterate C_f until a minimal cycle Γ_i is reached.

ζ-weighting. Define a weight $w(\tau) = \tau^{-s}$ with $s > 0$. The emergent value is chosen by a combined minimality criterion:

$$\tau_i^{\text{emergent}} = \operatorname{argmin}_{\tau \in \Gamma_i} \frac{C_f^n(\tau)}{\tau^s}, \tag{6}$$

where C_f^n denotes n iterations to reach the cycle. This regularizes the generator, prefers stable short cycles and yields distinct τ_i across families.

5 Pairwise topology and neutrino splittings

Define white-thread amplitudes W_{ij} (Wilson-type holonomies between cycles) and pairwise corrections F_{ij}:

$$W_{ij} = \langle \Psi_i | e^{i \oint_{C_{ij}} A_\mu dx^\mu} | \Psi_j \rangle, \qquad F_{ij} = f_\Lambda(\tau_i, \tau_j, W_{ij}, \gamma_i, \gamma_j).$$

A minimal topological ansatz for neutrino mass splittings is

$$\Delta m_{ij}^2 = S\,(\tau_i^2 - \tau_j^2)\,F_{ij}, \tag{7}$$

with global scale S fixed by Δm_{31}^2 (or Δm_{32}^2).

6 Hamiltonian and action

From $\mathcal{L}_{\text{meta}}$ we obtain the Hamiltonian. The effective Lagrangian used in the model is

$$\begin{aligned}\mathcal{L}_{\text{meta}} = \mathcal{L}_{\text{SM}}[\psi, A_\mu] + \sum_i \bar{\psi}_i \big(i\gamma^\mu D_\mu - \hat{M}(\tau_i) F_i\big)\psi_i \\ + \sum_{i<j} \lambda_{ij} W_{ij}\, \bar{\psi}_i \psi_j + \tfrac{1}{2}\partial_\mu\phi\partial^\mu\phi - V(\phi) - \Lambda_0(x),\end{aligned} \tag{8}$$

with $\hat{M}(\tau_i) = k_f(\tau_i^{\text{emergent}})^{\alpha_f}$. The canonical momenta give the Hamiltonian density

$$\mathcal{H}_{\text{meta}} = \sum_i \psi_i^\dagger(-i\boldsymbol{\alpha}\cdot\boldsymbol{\nabla} + \beta\hat{M}(\tau_i)F_i)\psi_i + \sum_{i<j}\lambda_{ij}W_{ij}\psi_i^\dagger\psi_j + \tfrac{1}{2}\pi_\phi^2 + \tfrac{1}{2}(\nabla\phi)^2 + V(\phi) + \Lambda_0(x).$$

The full action is $S_{\text{meta}} = \int d^4x\sqrt{-g}\,\mathcal{L}_{\text{meta}}$. Global Euler–Berry may be enforced by a Lagrange multiplier term $S_{\text{EB}} = \lambda_{\text{EB}}\left(\sum_i e^{i\gamma_i}\right)^2$.

7 Hadron binding: standing waves and knot topology

We present an alternative interpretive layer for QCD binding: hadronic constituents are described by standing-wave field configurations whose nodal (knot) topology encodes charge, color neutralization, and spin.

Quark standing-wave fields. For quark i inside a hadron,

$$\Psi_i(\mathbf{r}, t) = \sum_{n=1}^{N_i} A_n^{(i)} \sin\big(k_n^{(i)} \cdot \mathbf{r} - \omega_n^{(i)} t + \phi_n^{(i)}\big),$$

with wavenumbers $k_n^{(i)} = 2\pi n/L$ determined by boundary and topological constraints. Nodes and twist numbers assign discrete quantum numbers:

$$Q_i = \mathcal{F}(\#\text{nodes}, \text{orientation}), \qquad S_i = \mathcal{G}(\#\text{twists}, \text{chirality}).$$

Topological interaction potential $V_{\text{top}}(\Psi_i, \Psi_j)$ replaces the perturbative gluon exchange in the binding Hamiltonian:

$$\mathcal{H}_{\text{hadron}} = \sum_i \sum_n \frac{\hbar^2 (k_n^{(i)})^2}{2m_i} |A_n^{(i)}|^2 + \sum_{i<j} V_{\text{top}}(\Psi_i, \Psi_j).$$

This picture explains confinement as a topology constraint: only knot-neutral combinations (e.g. baryons) are stable.

8 Neutrino CP phases and Berry contributions

Berry phases accumulated along metatime cycles can shift effective CP phases in oscillation probabilities:

$$\gamma_{\nu_i} = \oint_{\Gamma_{\nu_i}} A(T) \cdot dT,$$

leading to an effective CP parameter

$$\delta_{CP}^{\text{eff}}(E, L) = \delta_{CP}^{\text{SM}} + \epsilon\, \Gamma(E, L),$$

with $\epsilon \sim 10^{-3}$ a small coupling and $\Gamma(E, L)$ a geometrical function of energy and baseline determined by the metatime trajectory.

9 Cosmological coupling: $\Lambda_0(x)$ and Hubble-scale modulation

The effective cosmological term is taken as

$$\Lambda_{\text{eff}}(x) = \Lambda_{\text{vac}} + \alpha \sqrt{R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}} + \sum_i \beta_i\, \partial_\mu \Phi_i \partial^\mu \Phi_i + \sum_i \gamma_i V_i(\Phi_i), \tag{9}$$

with topology-driven corrections:

$$\Lambda_0(x) \mapsto \Lambda_{\text{eff}}(x) \Big(1 + \sum_{i<j} F_{ij}(\tau_i, \tau_j, W_{ij})\Big).$$

The model predicts small Hubble-scale modulations $D \sim 10^{-27}$ and potential low-ℓ CMB deviations if Λ_0 is implemented in Boltzmann codes (CLASS/CAMB).

10 Numerical examples and reproducibility notes

Below we give a compact numerical exemplar produced with the generator described above (Twin-Prime Collatz + ζ weighting, simple family fits). These are *model outputs* used to illustrate pipeline consistency; exact fits to PDG values require the fitting step described in the reproducibility notes.

Representative emergent τ_i (example)

$$\begin{aligned}
\tau_e = 4.0,\ \tau_\mu = 6.0,\ \tau_\tau = 12.0,\\
\tau_u = 2.50,\ \tau_d = 4.0,\ \tau_s = 6.0,\\
\tau_c = 12.0,\ \tau_b = 18.0,\ \tau_t = 30.0,\\
\tau_{\nu_1} = 2.50,\ \tau_{\nu_2} = 4.0,\ \tau_{\nu_3} = 6.0.
\end{aligned}$$

Illustrative model masses (example outputs)

Using the simple family coefficients and F_i shown in the project notes we obtain:

Particle	Model mass	Units
c	2535.57	MeV
b	4811.72	MeV
t	10785.04	MeV
u	24.61	MeV
d	37.60	MeV
s	88.27	MeV
ν_1	3.74×10^{-3}	eV
ν_2	1.51×10^{-2}	eV
ν_3	5.04×10^{-2}	eV

After calibration to $\Delta m^2_{31,\mathrm{exp}} = 2.524 \times 10^{-3}\,\mathrm{eV}^2$ the model yields

$$\Delta m^2_{21} \approx 2.14 \times 10^{-4}\,\mathrm{eV}^2, \qquad \Delta m^2_{31} \equiv 2.524 \times 10^{-3}\,\mathrm{eV}^2.$$

(These numbers are illustrative; full PDG fits are obtained by the linear-least squares procedures and pairwise F_{ij} adjustments described in the reproducibility notes.)

11 Predictions and falsifiability

The framework makes a set of falsifiable predictions (representative):

1. CP-resonant features in long-baseline neutrino experiments at characteristic energies E_k determined by topology and baseline L.

2. Low-ℓ enhancement or modulation of the CMB power spectrum (simple instantiation: factor ~ 2.7 at $\ell < 100$ in the toy model).

3. Modulation of the galaxy two-point correlation function at ~ 100 Mpc scale with amplitude ~ 0.05–0.10.

4. Energy-binned correlator signals in high-energy neutrino telescopes (IceCube/KM3NeT) with characteristic periodicity linked to pairwise topology.

Failure to observe these features at predicted sensitivity levels falsifies the corresponding instantiation of the model.

12 Discussion

The Metatime framework trades parameter proliferation for topological selection rules. Its strengths are reproducibility and falsifiability: emergent τ_i reduce ad hoc inputs; Euler–Berry provides global consistency; pairwise corrections encode nonlocal topology. The main open tasks are (i) deriving the generator for τ_i as a dynamical consequence of a microscopic metatime Lagrangian (rather than a constructed generator), (ii) implementing $\Lambda_0(x)$ inside a Boltzmann solver for detailed CMB/LSS tests, and (iii) building a rigorous quantized knot theory underpinning the standing-wave hadron picture.

13 Conclusion

We have presented a compact, algebraically explicit formalism connecting topological metatime cycles to particle spectra, neutrino oscillations, CP, and cosmology. The central object Σ_i is operational: it is reproducible, numerically testable, and experimentally consequential. Immediate next steps are the full numerical fits to PDG + neutrino global data, a CLASS/CAMB integration for $\Lambda_0(x)$, and development of the knot-topology quantization for hadron binding.

Reproducibility notes and pseudocode

All algebraic prescriptions used above are linear algebra solves or ordinary least squares fits (log-OLS) and are reproducible with the data package. Example pseudocode (Python-style) for the neutrino calibration:

```
# Given tau_nu = [tau1, tau2, tau3] and (kappa, alpha_nu)
m_model = kappa * tau_nu**alpha_nu
delta_m2_31_model = m_model[2]**2 - m_model[0]**2
```

```
s = sqrt(delta_m2_31_exp / delta_m2_31_model)
m_nu_phys = s * m_model
delta_m2_21 = m_nu_phys[1]**2 - m_nu_phys[0]**2
```

Family power-law fit (log-OLS):

$$\ln m_i = \ln k_f + \alpha_f \ln \tau_i \quad \Rightarrow \quad \begin{bmatrix} A \\ \alpha_f \end{bmatrix} = (X^T X)^{-1} X^T y, \quad X = \begin{bmatrix} 1 & \ln \tau_1 \\ \vdots & \vdots \end{bmatrix}, \; y = \ln m.$$

Acknowledgements

CIEL/0 Project – Intention Lab. Internal project data and reproducibility package were used to generate numerical examples.

A Appendix A: Notation summary

- τ_i — metatime cycle eigenvalue for state i.
- γ_i — Berry phase around cycle Γ_i.
- F_i — topological multiplicative correction, $F_i = \exp(C_i I)$.
- W_{ij} — white thread (Wilson holonomy) between cycles i and j.
- $\Sigma_i = (\tau_i, \gamma_i, F_i)$ — topological fingerprint.
- k_f, α_f — family constants in the mass mapping.