

Modeling of optimal phytosanitary policies in crops of economic importance in the state of Sonora.

Gabriel Adrián Salcedo Varela

Departamento de matemáticas, División de Ciencias Exactas y Naturales
Universidad de Sonora

November 29, 2019

Contents

mm

40

60

80

100

120

1 Motivation

2 Objective


3 Epidemical Model

- Controlled Model

4 Existence Theory

5 Simulations

Tomato Leaf Curl Virus



mm

40

60

80

100

120



mm

40

60

80

100

120

40

Objective

Model optimal phytosanitary policies for diseases in agricultural crops.

60

80

mm

40

60

80

100

120



40 Holt, J. Colvin, and V. Muniyappa.

Identifying control strategies for tomato leaf curl virus disease using an epidemiological model.

Journal of Applied Ecology, 36(5):625–633, oct 1999.

60

80

Plant Model without control

Tomato Leaf Curl Virus Disease Using an Epidemiological Model

mm

40

60


80

100

120

Hypothesis:

- infection by infected plants and vectors,
- output and input for plants and vectors.



Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.



R. E. Shun-xiang, W. A. Zhen-zhong, Q. I. Bao-li, X. I. Yuan.

The pest status of *bemisia tabaci* in china and non-chemical control strategies*.

Insect Science, 8(3):279–288, 2001.



H. A. Smith and M. C. Giurcanu.

New Insecticides for Management of Tomato Yellow Leaf Curl, a Virus Vectored by the Silverleaf Whitefly, *Bemisia tabaci*.

Journal of Insect Science, 14(1):4–7, jan 2014.

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

$$\frac{dS_p}{dt} = -\beta_p S_p I_v + r(L_p + I_p),$$

$$\frac{dL_p}{dt} = \beta_p S_p I_v - bL_p - rL_p,$$

$$\frac{dI_p}{dt} = bL_p - rI_p,$$


$$\frac{dS_v}{dt} = -\beta_v S_v I_p - \gamma S_v - (1 - \theta)\mu,$$

$$\frac{dI_v}{dt} = \beta_v S_v I_p - \gamma I_v - \theta\mu,$$

$$S_p(0) = S_{p0}, L_p(0) = L_{p0}, I_p(0) = I_{p0},$$

$$S_v(0) = S_{v0}, I_v(0) = I_{v0}.$$

Par.	Value	Descrip.
β_p	0.1	plant latent rate
r	0.01	plant remove rate
b	0.075	plant infectious rate
γ	0.06	vector die or depar rate
μ	0.3	immigration rate
θ	0.2	infected vectors arrival
β_v	0.003	vector infected rate



$$R_0 = \sqrt{\frac{\beta_v \mu b \beta_p}{r^2 (r + b) \gamma}}.$$

60



40

If $R_0 < 1$,

$$\lim_{t \rightarrow \infty} (S_p, L_p, I_p, S_v, I_v) = (N_p, 0, 0, \frac{\mu}{\gamma}, 0).$$

If $R_0 < 1$,

$$\lim_{t \rightarrow \infty} (S_p, L_p, I_p, S_v, I_v) = (S_p^*, L_p^*, I_p^*, S_v^*, I_v^*).$$



Plant Model with control

Tomato Leaf Curl Virus Disease Using an Epidemiological Model

mm

40

60

80

100

120

40

$$\frac{dS_p}{dt} = -\beta_p S_p I_v + (r + u_1) L_p + (r + u_2) I_p,$$

$$\frac{dL_p}{dt} = \beta_p S_p I_v - b L_p - (r + u_1) L_p,$$

60

$$\frac{dI_p}{dt} = b L_p - (r + u_2) I_p,$$

$$\frac{dS_v}{dt} = -\beta_v S_v I_p - (\gamma + u_3) S_v - (1 - \theta) \mu,$$

80

$$\frac{dI_v}{dt} = \beta_v S_v I_p - (\gamma + u_3) I_v - \theta \mu,$$

Minimize

$$J(u_1, u_2, u_3) = \int_0^T A_1 I_p(t) + A_2 L_p(t) + A_3 I_v(t) + c_1 u_1(t)^2 + c_2 u_2(t)^2 + c_3 u_3(t)^2 dt,$$

subject to

$$\begin{cases} \frac{dS_p}{dt} = -\beta_p S_p I_v + (r + u_1) L_p + (r + u_2) I_p, \\ \frac{dL_p}{dt} = \beta_p S_p I_v - b L_p - (r + u_1) L_p, \\ \frac{dI_p}{dt} = b L_p - (r + u_2) I_p, \\ \frac{dS_v}{dt} = -\beta_v S_v I_p - (\gamma + u_3) S_v - (1 - \theta) \mu, \\ \frac{dI_v}{dt} = \beta_v S_v I_p - (\gamma + u_3) I_v - \theta \mu, \\ S_p(0) = S_{p0}, L_p(0) = L_{p0}, I_p(0) = I_{p0}, S_v(0) = S_{v0}, I_v(0) = I_{v0}. \end{cases}$$

$T \in (0, \infty)$ be fixed. Consider the control system:

$$\begin{cases} \dot{x}(s) = f(s, u(s), x(s)) & s \in [t_0, T], \\ x(t_0) = x_0, \end{cases}$$

with terminal state constraint

$$x(T; t_0, x_0, u(\cdot)) \in M,$$

where $M \subseteq \mathbb{R}^n$ is fixed.

$M : \mathbb{R}_+ \rightarrow 2^{\mathbb{R}^n}$ is a moving target in \mathbb{R}^n if for any $t \in \mathbb{R}_+$, $M(t)$ is a measurable.

$$J(t_0, x_0; u(\cdot)) = \int_{t_0}^T g(s, u(s), x(s)) ds + h(T, x(T)) \equiv J^T(t_0, x_0, u(\cdot)).$$

Problem $(OC)^T$

Given $(t_0, x_0) \in \mathbb{R}_+ \times \mathbb{R}^n$ with $\tilde{\mathcal{U}}_{x_0}^M[t_0, T] \neq \emptyset$, find a $\bar{u}(\cdot) \in \tilde{\mathcal{U}}_{x_0}^M[t_0, T]$ such that

$$J^T(t_0, x_0; \bar{u}(\cdot)) = \inf_{u(\cdot) \in \tilde{\mathcal{U}}_{x_0}^M[t_0, T]} J^T(t_0, x_0; u(\cdot)).$$

(C1)

$f : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is measurable, satisfies a lipchitz condition in x , and $|f(t, u, 0)| \leq L$, for every $(t, u) \in \mathbb{R}_+ \times U$.

$\omega : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$, increasing, and $\omega(r, 0) = 0$ for every $r \geq 0$.

(C2)

$g : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}$ and $h : \mathbb{R}^n \rightarrow \mathbb{R}$ are measurable, and

$$|g(s, u, x_1) - g(s, u, x_2)| + |h(x_1) - h(x_2)| \leq \omega(|x_1| \vee |x_2|, |x_1 - x_2|)$$

for every $(s, u) \in \mathbb{R}_+ \times U, x_1, x_2 \in \mathbb{R}^n$.

For any $(t, x) \in [0, T] \times \mathbb{R}^n$,

mm

$$\mathbb{E}(t, x) = \{(z^0, z) \in \mathbb{R} \times \mathbb{R}_+ \mid z^0 \geq g(t, u, x), z = f(t, u, x), u \in U\}.$$

Cesari property:

$$\bigcap_{\delta > 0} \text{co} \mathbb{E}(t, B_\delta(x)) = \mathbb{E}(t, x).$$

(C3)

For almost all $t \in [0, T]$, Cesari property holds for any $x \in \mathbb{R}^n$.

Existence Theorem

Let (C1)-(C3) hold. Let $M \subseteq \mathbb{R}^n$ be a non-empty closed set. Let $(t_0, x_0) \in [0, T] \times \mathbb{R}^n$ be given and $\tilde{\mathcal{U}}_x^M[t_0, T] \neq \emptyset$. Then problem $(OC)^T$ admits at least one optimal pair.

40

60

80

Pontryagin's Maximum Principle

If $u^*(t)$ and $x^*(t)$ are optimal for the problem $(OC)^T$, then there exists a piecewise differentiable adjoint variable $\lambda(t)$ such that

$$H(t, x^*(t), u(t), \lambda(t)) \leq H(t, x^*(t), u^*(t), \lambda(t))$$

for all controls u at each time t , where the Hamiltonian H is

$$H = g(t, x(t), u(t)) + \lambda(t)f(t, x(t), u(t)),$$

and

$$\lambda'(t) = -\frac{\partial H(t, x^*(t), u^*(t), \lambda(t))}{\partial x},$$

$$\lambda(T) = 0.$$

mm

Step 1. Make an initial guess for \vec{u} over the interval.

Step 2. Using the initial condition $x_1 = x(t_0) = a$ and the values for \vec{u} , solve \vec{x} forward in time according to its differential equation in the optimality system.

Step 3. Using the transversality condition $\lambda_{N+1} = \lambda(t_1) = 0$ and the values for \vec{u} and \vec{x} , solve $\vec{\lambda}$ backward in time according to its differential equation in the optimality system.

Step 4. Update \vec{u} by entering the new \vec{x} and $\vec{\lambda}$ values into the characterization of the optimal control.

Step 5. Check convergence. If the values of the variables in this iteration and the last iteration are negligibly close, output the current values as solutions. If values are not close, return to Step 2.

mm

40

60

80

100


120

40

60

80

Case with one controls



Case with two controls

mm

40

60

80


100

120

40

60

80



Case with three controls

mm

40

60

80

100

120

40

60

80

