

Modeling of optimal phytosanitary policies in crops of economic importance in the state of Sonora.

Gabriel Adrián Salcedo Varela

Departamento de matemáticas, División de Ciencias Exactas y Naturales
Universidad de Sonora

November 29, 2019

Contents

mm

40

60

80

100

120

1 Motivation

2 Objective


3 Epidemical Model

- Model without control
- \hat{c} Controlled Model


4 Existence Theory

5 Simulations

Tomato Leaf Curl Virus



Tomato Leaf Curl Virus



mm

40

60

80

100

120



mm

40

60

80

100

120

40

Objective

Model optimal phytosanitary policies for diseases in agricultural crops.

60

80

mm

40

60

80

100

120



40 Holt, J. Colvin, and V. Muniyappa.

Identifying control strategies for tomato leaf curl virus disease using an epidemiological model.

Journal of Applied Ecology, 36(5):625–633, oct 1999.

60

80

Tomato Yellow Leaf Virus Disease Using an Epidemiological Model

Gabriel Adrián Salcedo Varela

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.

80

Others Controls

mm

40

60

80

100

120

Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

Biological control

- Parasitoids,
- predators
- fungi.

Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyr.



R. E. Shun-xiang, W. A. Zhen-zhong, Q. I. Bao-li, X. I. Yuan.

The pest status of *bemisia tabaci* in china and non-chemical control strategies*.

Insect Science, 8(3):279–288, 2001.



H. A. Smith and M. C. Giurcanu.

New Insecticides for Management of Tomato Yellow Leaf Curl, a Virus Vectored by the Silverleaf Whitefly, *Bemisia tabaci*.

Journal of Insect Science, 14(1):4–7, jan 2014.

$$\frac{dS_p}{dt} = -\beta_p S_p I_v + r(L_p + I_p),$$

$$\frac{dL_p}{dt} = \beta_p S_p I_v - bL_p - rL_p,$$

$$\frac{dI_p}{dt} = bL_p - rI_p,$$


$$\frac{dS_v}{dt} = -\beta_v S_v I_p - \gamma S_v - (1 - \theta)\mu,$$

$$\frac{dI_v}{dt} = \beta_v S_v I_p - \gamma I_v - \theta\mu,$$

$$S_p(0) = S_{p0}, L_p(0) = L_{p0}, I_p(0) = I_{p0},$$

$$S_v(0) = S_{v0}, I_v(0) = I_{v0}.$$


Par.	Value	Descrip.
β_p	0.1	plant latent rate
r	0.01	plant remove rate
b	0.075	plant infectious rate
γ	0.06	vector die or depar rate
μ	0.3	immigration rate
θ	0.2	infected vectors arrival
β_v	0.003	vector infected rate



$$R_0 = \sqrt{\frac{\beta_v \mu b \beta_p}{r^2 (r + b) \gamma}}.$$

If $R_0 < 1$,


$$\lim_{t \rightarrow \infty} (S_p, L_p, I_p, S_v, I_v) = (N_p, 0, 0, \frac{\mu}{\gamma}, 0).$$



$$R_0 = \sqrt{\frac{\beta_v \mu b \beta_p}{r^2(r+b)\gamma}}.$$

If $R_0 > 1$,

$$\lim_{t \rightarrow \infty} (S_p, L_p, I_p, S_v, I_v) = (S_p^*, L_p^*, I_p^*, S_v^*, I_v^*).$$



Plant Model with control

Tomato Yellow Leaf Curl Virus Disease Using an Epidemiological Model 100 120

40

$$\frac{dS_p}{dt} = -\beta_p S_p I_v + (r + u_1) L_p + (r + u_2) I_p,$$

$$\frac{dL_p}{dt} = \beta_p S_p I_v - b L_p - (r + u_1) L_p,$$

60

$$\frac{dI_p}{dt} = b L_p - (r + u_2) I_p,$$

$$\frac{dS_v}{dt} = -\beta_v S_v I_p - (\gamma + u_3) S_v - (1 - \theta) \mu,$$

80

$$\frac{dI_v}{dt} = \beta_v S_v I_p - (\gamma + u_3) I_v - \theta \mu,$$

Minimize

$$J(u_1, u_2, u_3) = \int_0^T A_1 I_p(t) + A_2 L_p(t) + A_3 I_v(t) + c_1 u_1(t)^2 + c_2 u_2(t)^2 + c_3 u_3(t)^2 dt,$$

subject to

$$\begin{cases} \frac{dS_p}{dt} = -\beta_p S_p I_v + (r + u_1) L_p + (r + u_2) I_p, \\ \frac{dL_p}{dt} = \beta_p S_p I_v - b L_p - (r + u_1) L_p, \\ \frac{dI_p}{dt} = b L_p - (r + u_2) I_p, \\ \frac{dS_v}{dt} = -\beta_v S_v I_p - (\gamma + u_3) S_v - (1 - \theta) \mu, \\ \frac{dI_v}{dt} = \beta_v S_v I_p - (\gamma + u_3) I_v - \theta \mu, \\ S_p(0) = S_{p0}, L_p(0) = L_{p0}, I_p(0) = I_{p0}, S_v(0) = S_{v0}, I_v(0) = I_{v0}. \end{cases}$$

Existence Theory

mm

40

60

80

100

120

$$\begin{cases} \dot{x}(s) = f(s, u(s), x(s)) & s \in [t_0, T], \\ x(t_0) = x_0, \end{cases}$$

with terminal state constraint

$$x(T; t_0, x_0, u(\cdot)) \in M, M \subseteq \mathbb{R}^n$$

$$J(t_0, x_0; u(\cdot)) = \int_{t_0}^T g(s, u(s), x(s)) ds + h(T, x(T)).$$

Problem $(OC)^T$

$(t_0, x_0) \in \mathbb{R}_+ \times \mathbb{R}^n$ with $\tilde{\mathcal{U}}_{x_0}[t_0, T] \neq \emptyset$, find a $\bar{u}(\cdot) \in \tilde{\mathcal{U}}_{x_0}[t_0, T]$ s.t.

$$J(t_0, x_0; \bar{u}(\cdot)) = \inf_{u(\cdot) \in \tilde{\mathcal{U}}_{x_0}[t_0, T]} J(t_0, x_0; u(\cdot)).$$

Hypothesis:

(C-1) $f : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is measurable, satisfies a lipchitz condition in x ,
 $|f(t, u, 0)| \leq L, \forall (t, u) \in \mathbb{R}_+ \times U$.

(C-2) $g : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}$,
 $h : \mathbb{R}^n \rightarrow \mathbb{R}$ are measurable,
and

$$\begin{aligned} & |g(s, u, x_1) - g(s, u, x_2)| + \\ & |h(x_1) - h(x_2)| \\ & \leq \omega(|x_1| \vee |x_2|, |x_1 - x_2|) \end{aligned}$$

$$\forall (s, u) \in \mathbb{R}_+ \times U, x_1, x_2 \in \mathbb{R}^n.$$

(C-3) For a.a. $t \in [0, T]$, Cesari property holds $\forall x \in \mathbb{R}^n$.

Hypothesis:

(C-1) $f : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is measurable, satisfies a lipchitz condition in x ,
 $|f(t, u, 0)| \leq L, \forall (t, u) \in \mathbb{R}_+ \times U$.

(C-2) $g : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}$,
 $h : \mathbb{R}^n \rightarrow \mathbb{R}$ are measurable,
and

$$\begin{aligned} & |g(s, u, x_1) - g(s, u, x_2)| + \\ & |h(x_1) - h(x_2)| \\ & \leq \omega(|x_1| \vee |x_2|, |x_1 - x_2|) \end{aligned}$$

$\forall (s, u) \in \mathbb{R}_+ \times U, x_1, x_2 \in \mathbb{R}^n$.

(C-3) For a.a. $t \in [0, T]$, Cesari property holds $\forall x \in \mathbb{R}^n$.

$\omega : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$, increasing, $\omega(r, 0) = 0 \forall r \geq 0$.

Hypothesis:

(C-1) $f : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is measurable, satisfies a lipchitz condition in x ,
 $|f(t, u, 0)| \leq L, \forall (t, u) \in \mathbb{R}_+ \times U$.

(C-2) $g : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}$,
 $h : \mathbb{R}^n \rightarrow \mathbb{R}$ are measurable, and

$$\begin{aligned} & |g(s, u, x_1) - g(s, u, x_2)| + \\ & |h(x_1) - h(x_2)| \\ & \leq \omega(|x_1| \vee |x_2|, |x_1 - x_2|) \end{aligned}$$

$$\forall (s, u) \in \mathbb{R}_+ \times U, x_1, x_2 \in \mathbb{R}^n.$$

(C-3) For a.a. $t \in [0, T]$, Cesari property holds $\forall x \in \mathbb{R}^n$.

80 — 100 — 120

$$\begin{aligned} \mathbb{E}(t, x) = \{ & (z^0, z) \in \mathbb{R} \times \mathbb{R}_+ | \\ & z^0 \geq g(t, u, x), \\ & z = f(t, u, x), u \in U \}. \end{aligned}$$

Cesari property

$$\bigcap_{\delta} \bar{\text{co}} \mathbb{E}(t, B_{\delta}(x)) = \mathbb{E}(t, x)$$

Existence Theorem

Let (C1)-(C3) hold. Then problem $(OC)^T$ admits at least one optimal pair.

40

60

80

mm

40

60

80

100

120

$$H = g(t, x(t), u(t)) + \lambda(t)f(t, x(t), u(t)).$$

40

$$H = A_1 I_v + A_2 L_p + A_3 I_v \sum_{i=1}^3 c_i u_i^2 + \lambda_1 (-\beta_p S_p I_v + (r + u_1) L_p + (r + u_2) I_p) \\ + \lambda_2 (\beta_p S_p I_v - b L_p - (r + u_1) L_p) + \lambda_3 (b L_p - (r + u_2) I_p) \\ + \lambda_4 (-\beta_v S_v I_p - (\gamma + u_3) S_v - (1 - \theta) \mu) + \lambda_5 (\beta_v S_v I_p - (\gamma + u_3) I_v - \theta \mu).$$

80

Pontryagin's Maximum Principle

If $u^*(t)$ and $x^*(t)$ are optimal for the problem $(OC)^T$, then there exists a piecewise differentiable adjoint variable $\lambda(t)$ s.t.

$$H(t, x^*(t), u(t), \lambda(t)) \leq H(t, x^*(t), u^*(t), \lambda(t))$$

$\forall u$ at t ,

$$\lambda'(t) = -\frac{\partial H(t, x^*(t), u^*(t), \lambda(t))}{\partial x},$$

$$\lambda(T) = 0.$$

$$\frac{d\lambda_1}{dt} = \beta_p(\lambda_1 - \lambda_2),$$

$$\frac{d\lambda_2}{dt} = -A_2 + (r + u_1)(\lambda_2 - \lambda_1) + b(\lambda_2 - \lambda_3),$$

$$\frac{d\lambda_3}{dt} = -A_1 + (r + u_2)(\lambda_3 - \lambda_1) + \beta_v S_v(\lambda_4 - \lambda_5),$$

$$\frac{d\lambda_4}{dt} = \beta_v I_p(\lambda_4 - \lambda_5) + (\gamma + u_3)\lambda_4,$$

$$\frac{d\lambda_5}{dt} = -A_3 + \beta_p S_p(\lambda_1 - \lambda_2) + (\gamma + u_3)\lambda_5,$$

80

$$u_1^* \min \left(\max \left(0, \frac{I_p(\lambda_2 - \lambda_1)}{2c_1} \right), u_{1m} \right)$$

$$u_2^* \min \left(\max \left(0, \frac{I_p(\lambda_3 - \lambda_1)}{2c_2} \right), u_{2m} \right)$$

$$u_3^*$$

The most popular

mm

40

60

80

100

120

Algorithm 2 Forward Backward Sweep

Input: $t_0, t_f, x_0, h, \text{tol}, \lambda_f$

Output: x^*, u^*, λ

procedure FORWARD_BACKWARD_SWEEP($g, \lambda_{\text{function}}, u, x_0, \lambda_f, h, n_{\text{max}}$)

while $\epsilon > \text{tol}$ **do**

 40 $u_{\text{old}} \leftarrow u$

$x_{\text{old}} \leftarrow x$

$x \leftarrow \text{RUNGE_KUTTA_FORWARD}(g, u, x_0, h)$

$\lambda_{\text{old}} \leftarrow \lambda$

$\lambda \leftarrow \text{RUNGE_KUTTA_BACKWARD}(\lambda_{\text{function}}, x, \lambda_f, h)$

$u_1 \leftarrow \text{OPTIMALITY_CONDITION}(u, x, \lambda)$

$u \leftarrow \alpha u_1 + (1 - \alpha) u_{\text{old}}, \quad \alpha \in [0, 1]$

▷ convex combination

 60 $\epsilon_u \leftarrow ||u - u_{\text{old}}||$

$\epsilon_u \leftarrow \frac{||u||}{||u||}$

$\epsilon_x \leftarrow \frac{||x - x_{\text{old}}||}{||x||}$

▷ relative error

$\epsilon_x \leftarrow \frac{||x||}{||x||}$

$\epsilon_\lambda \leftarrow \frac{||\lambda - \lambda_{\text{old}}||}{||\lambda||}$

 80 $\epsilon \leftarrow \max\{\epsilon_u, \epsilon_x, \epsilon_\lambda\}$

end while


return x^*, u^*, λ

▷ Optimal pair

end procedure


Case with one controls

mm



Case with two controls

mm



Case with three controls

