

# **mm**Modeling of optimal phytosanitary policies in crops of economic importance in the state of Sonora.

40

Gabriel Adrián Salcedo Varela

Departamento de matemáticas, División de Ciencias Exactas y Naturales  
Universidad de Sonora

November 30, 2019

# Contents

- 1 Motivation
- 2 Objective
- 3 Epidemical Model
  - Model without control
  - Controlled Model
- 4 Existence Theory
- 5 Optimal Control Characterization
- 6 Simulations

# Tomato Leaf Curl Virus



10

60

80

100

120



60

80

# Tomato Leaf Curl Virus



mm

10

60

80

100

120



60

80

mm

40

60

80

100

120



mm

40

60

80

100

120

40

Objetive

Model optimal phytosanitary policies for diseases in agricultural crops.

60


80

# Plant Model without control

Tomato Muri Virus Disease Using an Epidemiological Model

## Hypothesis:

- Remove from latent and infected plants,
- plants become latent plants by infected vectors,
- latent plants become infectious plants,
- vectors become infected vectors by infected plants,
- vectors die per day,
- immigration from alternative hosts.



# Others Controls

**mm**

**40**

**60**

**80**

**100**

**120**

## Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

## Biological control

- Parasitoids,
- predators
- fungi.

## Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyrr.

**80**

# Others Controls

**mm**

**40**

**60**

**80**

**100**

**120**

## Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

## Biological control

- Parasitoids,
- predators
- fungi.

## Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyrr.

**80**

# Others Controls

**mm**

**40**

**60**

**80**

**100**

**120**

## Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

## Biological control

- Parasitoids,
- predators
- fungi.

## Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyrr.

**80**

# Others Controls

mm

40

60

80

100

120

## Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

## Biological control

- Parasitoids,
- predators
- fungi.

## Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyrr.

80

# Others Controls

**mm**

**40**

**60**

**80**

**100**

**120**

## Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

## Biological control

- Parasitoids,
- predators
- fungi.

## Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyrr.

**80**

# Others Controls

**mm**

**40**

**60**

**80**

**100**

**120**

## Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

## Biological control

- Parasitoids,
- predators
- fungi.

## Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyrr.

**80**

# Others Controls

**mm**

**40**

**60**

**80**

**100**

**120**

## Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

## Biological control

- Parasitoids,
- predators
- fungi.

## Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyrr.

**80**

# Others Controls

**mm**

**40**

**60**

**80**

**100**

**120**

## Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

## Biological control

- Parasitoids,
- predators
- fungi.

## Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyrr.

**80**

# Others Controls

mm

40

60

80

100

120

## Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

## Biological control

- Parasitoids,
- predators
- fungi.

## Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyrr.

80

# Others Controls

**mm**

**40**

**60**

**80**

**100**

**120**

## Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

## Biological control

- Parasitoids,
- predators
- fungi.

## Insecticide

- Pymetrozine,
- flupyradifurone,
- cyazypyrr.

**80**

## Others Controls

mm

40

60

80

100

120

### Cultural Control

- physical barriers,
- planting dates,
- removal of infested plants,
- host plant resistance.

### Biological control

- Parasitoids,
- predators
- fungi.



R. E. Shun-xiang, W. A. Zhen-zhong, Q. I. Bao-li, X. I. Yuan.

The pest status of *bemisia tabaci* in China and non-chemical control strategies\*.

*Insect Science*, 8(3):279–288, 2001.



H. A. Smith and M. C. Giurcanu.

80  
New Insecticides for Management of Tomato Yellow Leaf Curl, a Virus Vectored by the Silverleaf Whitefly, *Bemisia tabaci*.

*Journal of Insect Science*, 14(1):4–7, jan 2014.

mm

40

60

80

100

120



J. Holt, J. Colvin, and V. Muniyappa.

Identifying control strategies for tomato leaf curl virus disease using an epidemiological model.

*Journal of Applied Ecology*, 36(5):625–633, oct 1999.

60

80

$$\frac{dS_p}{dt} = -\beta_p S_p I_v + \textcolor{blue}{r}(L_p + I_p),$$

$$\frac{dL_p}{dt} = \beta_p S_p I_v - b L_p - \textcolor{blue}{r} L_p,$$

$$\frac{dI_p}{dt} = b L_p - \textcolor{blue}{r} I_p,$$

$$\frac{dS_v}{dt} = -\beta_v S_v I_p - \textcolor{brown}{g} S_v + (1 - \theta)\mu,$$

$$\frac{dI_v}{dt} = \beta_v S_v I_p - \textcolor{brown}{g} I_v + \theta\mu,$$

$$S_p(0) = S_{p_0}, L_p(0) = L_{p_0}, I_p(0) = I_{p_0},$$

$$S_v(0) = S_{v_0}, I_v(0) = I_{v_0}.$$

80

80

100

120

Par.	Unit	40	60	80	100	120
$\beta_p$	vector <sup>-1</sup> day <sup>-1</sup>					
$r$	day <sup>-1</sup>					
$b$	day <sup>-1</sup>					
$\gamma$	day <sup>-1</sup>	40				
$\mu$	plant <sup>-1</sup> day <sup>-1</sup>					
$\theta$	proportion					
$\beta_v$	plant <sup>-1</sup> day <sup>-1</sup>					

$$\frac{dS_p}{dt} = -\beta_p S_p I_v + \textcolor{blue}{r}(L_p + I_p),$$

$$\frac{dL_p}{dt} = \beta_p S_p I_v - b L_p - \textcolor{blue}{r} L_p,$$


$$\frac{dI_p}{dt} = b L_p - \textcolor{blue}{r} I_p,$$

$$\frac{dS_v}{dt} = -\beta_v S_v I_p - \textcolor{brown}{y} S_v + (1 - \theta) \mu,$$

$$\frac{dI_v}{dt} = \beta_v S_v I_p - \textcolor{brown}{y} I_v + \theta \mu,$$

$$S_p(0) = S_{p_0}, L_p(0) = L_{p_0}, I_p(0) = I_{p_0},$$

$$S_v(0) = S_{v_0}, I_v(0) = I_{v_0}.$$



80

$$R_0 = \sqrt{\frac{\beta_v \mu b \beta_p}{r^2(r+b)\gamma}}.$$

40

60

80

100

120


60

80

$$R_0 = \sqrt{\frac{\beta_v \mu b \beta_p}{r^2(r+b)\gamma}}.$$

If  $R_0 < 1$ ,


$$\lim_{t \rightarrow \infty} (S_p, L_p, I_p, S_v, I_v) = (N_p, 0, 0, \frac{\mu}{\gamma}, 0).$$



$$R_0 = \sqrt{\frac{\beta_v \mu b \beta_p}{r^2(r+b)\gamma}}.$$

If  $R_0 > 1$ ,

$$\lim_{t \rightarrow \infty} (S_p, L_p, I_p, S_v, I_v) = (S_p^*, L_p^*, I_p^*, S_v^*, I_v^*).$$



# Plant Model with control

Tomato Mosaic Virus Disease Using an Epidemiological Model

$$\frac{dS_p}{dt} = -\beta_p S_p I_v + (r + u_1) L_p + (r + u_2) I_p,$$

$$\frac{dL_p}{dt} = \beta_p S_p I_v - b L_p - (r + u_1) L_p,$$

$$\frac{dI_p}{dt} = b L_p - (r + u_2) I_p,$$

$$\frac{dS_v}{dt} = -\beta_v S_v I_p - (\gamma + u_3) S_v + (1 - \theta)\mu,$$

$$\frac{dI_v}{dt} = \beta_v S_v I_p - (\gamma + u_3) I_v + \theta\mu,$$

## Controls:

- $u_1$ : replanting latent plant,
- $u_2$ : replanting infected plants,
- $u_3$ : fumigation.

$$\min_{\bar{u}(\cdot) \in \tilde{\mathcal{U}}_{x_0}[t_0, T]} J(u_1, u_2, u_3) = \int_0^T (A_1 I_p(t) + A_2 L_p(t) + A_3 I_v(t) \\ + c_1 u_1(t)^2 + c_2 u_2(t)^2 + c_3 u_3(t)^2) dt,$$

s.t.

$$\frac{dS_p}{dt} = -\beta_p S_p I_v + (r + u_1) L_p + (r + u_2) I_p,$$

$$\frac{dL_p}{dt} = \beta_p S_p I_v - b L_p - (r + u_1) L_p,$$

$$\frac{dI_p}{dt} = b L_p - (r + u_2) I_p,$$

$$\frac{dS_v}{dt} = -\beta_v S_v I_p - (\gamma + u_3) S_v + (1 - \theta) \mu,$$

$$\frac{dI_v}{dt} = \beta_v S_v I_p - (\gamma + u_3) I_v + \theta \mu,$$

$$S_p(0) = S_{p_0}, L_p(0) = L_{p_0},$$

$$I_p(0) = I_{p_0}, S_v(0) = S_{v_0}, I_v(0) = I_{v_0}.$$

# Existence Theory

mm

40

60

80

100

120

$$\begin{cases} \dot{x}(s) = f(s, u(s), x(s)) & s \in [t_0, T], \\ x(t_0) = x_0, \end{cases}$$

with terminal state constraint

$$x(T; t_0, x_0, u(\cdot)) \in M, M \subseteq \mathbb{R}^n.$$

$$J(t_0, x_0; u(\cdot)) = \int_{t_0}^T g(s, u(s), x(s)) ds + h(T, x(T)).$$

mm

40

60

80

100

120

## Problem ( $OC$ )

$(t_0, x_0) \in \mathbb{R}_+ \times \mathbb{R}^n$  with  $\tilde{\mathcal{U}}_{x_0}[t_0, T] \neq \emptyset$ , find a  $\bar{u}(\cdot) \in \tilde{\mathcal{U}}_{x_0}[t_0, T]$  s.t.

$$J(t_0, x_0; \bar{u}(\cdot)) = \inf_{u(\cdot) \in \tilde{\mathcal{U}}_{x_0}[t_0, T]} J(t_0, x_0; u(\cdot)).$$

60

80

## Hypothesis:

- (C-1)  $f : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is measurable, satisfies a lipchitz condition in  $x$ ,  $|f(t, u, 0)| \leq L, \forall (t, u) \in \mathbb{R}_+ \times U$ .
- (C-2)  $g : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $h : \mathbb{R}^n \rightarrow \mathbb{R}$  are measurable, and
- $$\begin{aligned} & |g(s, u, x_1) - g(s, u, x_2)| + \\ & |h(x_1) - h(x_2)| \\ & \leq \omega(|x_1| \vee |x_2|, |x_1 - x_2|) \end{aligned}$$
- $$\forall (s, u) \in \mathbb{R}_+ \times U, x_1, x_2 \in \mathbb{R}^n.$$
- (C-3) For a.a.  $t \in [0, T]$ , Cesari property holds  $\forall x \in \mathbb{R}^n$ .

80      100      120

## Hypothesis:

- (C-1)  $f : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is measurable, satisfies a lipchitz condition in  $x$ ,  $|f(t, u, 0)| \leq L, \forall (t, u) \in \mathbb{R}_+ \times U$ .
- (C-2)  $g : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $h : \mathbb{R}^n \rightarrow \mathbb{R}$  are measurable, and
- $$\begin{aligned} & |g(s, u, x_1) - g(s, u, x_2)| + \\ & |h(x_1) - h(x_2)| \\ & \leq \omega(|x_1| \vee |x_2|, |x_1 - x_2|) \end{aligned}$$
- $$\forall (s, u) \in \mathbb{R}_+ \times U, x_1, x_2 \in \mathbb{R}^n.$$
- (C-3) For a.a.  $t \in [0, T]$ , Cesari property holds  $\forall x \in \mathbb{R}^n$ .

$\omega : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ , increasing,  $\omega(r, 0) = 0 \forall r \geq 0$ .

## Hypothesis:

- (C-1)  $f : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is measurable, satisfies a lipchitz condition in  $x$ ,  $|f(t, u, 0)| \leq L, \forall (t, u) \in \mathbb{R}_+ \times U$ .
- (C-2)  $g : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $h : \mathbb{R}^n \rightarrow \mathbb{R}$  are measurable, and
- $$\begin{aligned} & |g(s, u, x_1) - g(s, u, x_2)| + \\ & |h(x_1) - h(x_2)| \\ & \leq \omega(|x_1| \vee |x_2|, |x_1 - x_2|) \end{aligned}$$
- $$\forall (s, u) \in \mathbb{R}_+ \times U, x_1, x_2 \in \mathbb{R}^n.$$
- (C-3) For a.a.  $t \in [0, T]$ , Cesari property holds  $\forall x \in \mathbb{R}^n$ .

80      100      120

$$\begin{aligned} \mathbb{E}(t, x) = & \{(z^0, z) \in \mathbb{R} \times \mathbb{R}^n | \\ & z^0 \geq g(t, u, x), \\ & z = f(t, u, x), u \in U\}. \end{aligned}$$

## Cesari property

$$\bigcap_{\delta} \bar{co} \mathbb{E}(t, B_\delta(x)) = \mathbb{E}(t, x)$$

## Hypothesis:

- (C-1)  $f : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  is measurable, satisfies a lipchitz condition in  $x$ ,  $|f(t, u, 0)| \leq L, \forall (t, u) \in \mathbb{R}_+ \times U$ .
- (C-2)  $g : \mathbb{R}_+ \times U \times \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $h : \mathbb{R}^n \rightarrow \mathbb{R}$  are measurable, and
- $$\begin{aligned} & |g(s, u, x_1) - g(s, u, x_2)| + \\ & |h(x_1) - h(x_2)| \\ & \leq \omega(|x_1| \vee |x_2|, |x_1 - x_2|) \end{aligned}$$
- $$\forall (s, u) \in \mathbb{R}_+ \times U, x_1, x_2 \in \mathbb{R}^n.$$
- (C-3) For a.a.  $t \in [0, T]$ , Cesari property holds  $\forall x \in \mathbb{R}^n$ .

## Existence Theorem

Let (C1)-(C3) hold. Then problem (*OC*) admits at least one optimal pair.

mm

$$H = g(t, x(t), u(t)) + \lambda(t)f(t, x(t), u(t)),$$

mm

$$H = g(t, x(t), u(t)) + \lambda(t)f(t, x(t), u(t)),$$

$$\begin{cases} \dot{x}(s) = f(s, u(s), x(s)) & s \in [t_0, T], \\ x(t_0) = x_0, \end{cases}$$

$$J(t_0, x_0; u(\cdot)) = \int_{t_0}^T g(s, u(s), x(s)) ds + h(x(T)).$$

80

## Pontryagin's Maximum Principle

If  $u^*(t)$  and  $x^*(t)$  are optimal for the problem ( $OC$ ), then there exists a piecewise differentiable adjoint variable  $\lambda(t)$  s.t.

$$H(t, x^*(t), u(t), \lambda(t)) \leq H(t, x^*(t), u^*(t), \lambda(t))$$

$\forall u$  at  $t$ ,

$$\lambda'(t) = -\frac{\partial H(t, x^*(t), u^*(t), \lambda(t))}{\partial x},$$

$$\lambda(T) = 0.$$

mm

$$\begin{aligned} H = & A_1 I_v + A_2 L_p + A_3 I_p + \sum_{i=1}^3 c_i u_i^2 \\ & + \lambda_1(-\beta_p S_p I_v + (r + u_1) L_p + (r + u_2) I_p) \\ & + \lambda_2(\beta_p S_p I_v - b L_p - (r + u_1) L_p) \\ & + \lambda_3(b L_p - (r + u_2) I_p) \\ & + \lambda_4(-\beta_v S_v I_p - (\gamma + u_3) S_v + (1 - \theta) \mu) \\ & + \lambda_5(\beta_v S_v I_p - (\gamma + u_3) I_v + \theta \mu). \end{aligned}$$

**mm**

40

120

$$\frac{d\lambda_1}{dt} = \beta_p(\lambda_1 - \lambda_2),$$

$$\frac{d\lambda_2}{dt} = -A_2 + (r + u_1)(\lambda_2 - \lambda_1) + b(\lambda_2 - \lambda_3),$$

$$\frac{d\lambda_3}{dt} = A_1 + (r + u_2)(\lambda_3 - \lambda_1) + \beta_v S_v(\lambda_4 - \lambda_5),$$

$$\frac{d\lambda_4}{dt} = \beta_v I_p(\lambda_4 - \lambda_5) + (\gamma + u_3)\lambda_4,$$

$$\frac{d\lambda_5}{dt} = -A_3 + \beta_p S_p(\lambda_1 - \lambda_2) + (\gamma + u_3)\lambda_5,$$

6

8

$$u_1^* = \min \left( \max \left( 0, \frac{L_p(\lambda_2 - \lambda_1)}{2c_1} \right), u_{1_{max}} \right)$$

$$u_2^* = \min \left( \max \left( 0, \frac{I_p(\lambda_3 - \lambda_1)}{2c_2} \right), u_{2_{max}} \right)$$

$$u_3^* = \min \left( \max \left( 0, \frac{S_v \lambda_4 + I_v \lambda_5}{2c_3} \right), u_{3_{max}} \right)$$

# The most popular

**mm**

**40**

**60**

**80**

**100**

**120**

## Algorithm 2 Forward Backward Sweep

**Input:**  $t_0, t_f, x_0, h, \text{tol}, \lambda_f$

**Output:**  $x^*, u^*, \lambda$

**procedure** FORWARD - BACKWARD - SWEEP( $g, \lambda_{\text{function}}, u, x_0, \lambda_f, h, n_{\max}$ )

**while**  $\epsilon > \text{tol}$  **do**

$u_{\text{ld}} \leftarrow u$

$x_{\text{old}} \leftarrow x$

$x \leftarrow \text{RUNGE\_KUTTA\_FORWARD}(g, u, x_0, h)$

$\lambda_{\text{old}} \leftarrow \lambda$

$\lambda \leftarrow \text{RUNGE\_KUTTA\_BACKWARD}(\lambda_{\text{function}}, x, \lambda_f, h)$

$u_1 \leftarrow \text{OPTIMALITY\_CONDITION}(u, x, \lambda)$

$u \leftarrow \alpha u_1 + (1 - \alpha) u_{\text{old}}, \quad \alpha \in [0, 1]$

        ▷ convex combination

$\epsilon_u \leftarrow \frac{\|u - u_{\text{old}}\|}{\|u\|}$

$\epsilon_x \leftarrow \frac{\|x - x_{\text{old}}\|}{\|x\|}$

        ▷ relative error

$\epsilon_\lambda \leftarrow \frac{\|\lambda - \lambda_{\text{old}}\|}{\|\lambda\|}$

$\epsilon \leftarrow \max\{\epsilon_u, \epsilon_x, \epsilon_\lambda\}$


**end while**

**return**  $x^*, u^*, \lambda$


    ▷ Optimal pair

**end procedure**

## Case with one controls




## Case with two controls




## Case with three controls

mm 40 60 80 100 120




## Case with three controls

mm 40 60 80 100 120



# Cost Function Comparation



mm

40

T. M. Hilker, J. S. Allen, A. Bokil, J. Briggs, Y. Feng, K. A.

Garrett, L. J. Gross, F. M. Hamelin, M. J. Jeger, C. A. Manore,  
A. G. Power, M. G. Redinbaugh, M. A. Rúa, and N. J. Cunniffe.

Modeling Virus Coinfection to Inform Management of Maize Lethal  
Necrosis in Kenya.

40

*Phytopathology*, 107(10):PHYTO-03-17-008, aug 2017.

P. Grandits, R. M. Kovacevic, and V. M. Veliov.

Optimal control and the value of information for a stochastic  
epidemiological SIS-model.

60

*Journal of Mathematical Analysis and Applications*,

476(2):665–695, aug 2019.

J. Yong and X. Y. Zhou.

*Stochastic controls: Hamiltonian systems and HJB equations*,

80  
Volume 43.

Springer Science & Business Media, 1999.