

R Code for Examples in the book

"Statistics: The Art and Science of Learning from Data" by Agresti, Franklin and Klingenberg, 5th edition

Chapter 14

Example 6: Friends and Happiness – Tukey Test

Reading in data

```
gss <-
read.csv(file='https://raw.githubusercontent.com/artofstat/data/master/Chapte
r14/gss_1998_happy_numfriends.csv')
head(data)
##
     NumFriends X1.VERY.HAPPY X2.PRETTY.HAPPY X3.NOT.TOO.HAPPY
## 1
               1
                              8
                                              33
                                                                 10
## 2
               2
                             35
                                              82
                                                                 17
## 3
               3
                             38
                                              79
                                                                 15
## 4
               4
                             36
                                              75
                                                                  7
## 5
               5
                             28
                                              57
                                                                 10
## 6
                             30
                                              35
                                                                  5
```

To convert the data into long format, you can use the pivot_longer() function from the tidyverse library

```
library(tidyverse)
gss <-
  gss %>%
  rename_at(2:4, ~ c('very_happy', 'pretty_happy', 'not_too_happy')) %>%
  pivot_longer(2:4, names_to = 'Happiness', values_to = 'Count') %>%
  uncount(Count) %>%
  relocate(Happiness)
attach(gss)
tapply(NumFriends, Happiness, summary)
## $not_too_happy
##
      Min. 1st Qu.
                    Median
                              Mean 3rd Qu.
                                              Max.
##
             2.000
                     4.000
                                     6.000 50.000
     1.000
                             6.271
##
## $pretty_happy
##
      Min. 1st Qu.
                    Median
                              Mean 3rd Qu.
                                              Max.
                     4.000
##
     1.000
             2.500
                             5.667
                                     6.000 60.000
##
## $very_happy
```

```
## Min. 1st Qu. Median Mean 3rd Qu. Max.
## 1.000 3.000 5.000 7.577 10.000 75.000
```

To perform a Tukey test

```
myAnova <- aov(NumFriends ~ Happiness)</pre>
tukeyTest <- TukeyHSD(myAnova)</pre>
tukeyTest
##
     Tukey multiple comparisons of means
##
       95% family-wise confidence level
##
## Fit: aov(formula = NumFriends ~ Happiness)
##
## $Happiness
                                     diff
##
                                                 lwr
                                                          upr
                                                                   p adj
## pretty_happy-not_too_happy -0.6039216 -2.5876540 1.379811 0.7547887
## very_happy-not_too_happy
                               1.3061908 -0.7860212 3.398403 0.3079598
## very_happy-pretty_happy
                               1.9101124 0.6170907 3.203134 0.0015952
```