ADRIANO MONTEIRO

NUTRITION COACH

Sleep, Recovery & Mental Resilience

The foundation your training, your food and your mood are built on

COMPLETE GUIDE 5 OF 5

adrianonutrition.com

Contents

Start here

Why sleep is the foundation

How sleep works (the short version)

Set up the night

Caffeine, alcohol, food and supplements

The evening routine

Daytime: light, naps and deep rest

When sleep does not improve

ADHD and sleep

The stress equation and recovery

Building resilience

A recovery day, stitched together

Your 14-day plan

Pick your case: starter plans

Worksheets

> **Disclaimer.** This educational resource presents sleep, recovery and stress-management strategies for personal development. It does not constitute medical, psychological or therapeutic advice. The author is a nutrition coach, not a licensed healthcare provider. If you have persistent insomnia, symptoms of sleep apnea, a mental health condition, ADHD, a substance dependence or any other clinical concern, consult a qualified professional before making changes. Cold exposure and breathing practices are not suitable for everyone. Individual results vary.

BEFORE YOU READ

Start here

You can have a perfect meal plan and a perfect program and still get poor results if you sleep five hours, run on stress and never recover. This guide covers the three things that sit under everything else: **sleep, recovery and resilience.**

This area is full of exaggerated claims: neurotransmitter boosts of several hundred percent, brain changes in weeks, miracle protocols. Most of those numbers come from tiny studies or none at all. In this guide I tell you how strong the evidence is for each tool and I leave out the numbers I could not verify. The basics are boring and they work: a regular schedule, light at the right times, a cool dark room, less caffeine and alcohol, daily movement, and people.

How to use this guide

- **Start with Chapters 1 to 5.** Sleep is the biggest lever and the most fixable.
- **Pick ONE change per week.** The 14-day plan in Chapter 12 shows the order.
- **Struggling for weeks despite good habits?** Read Chapter 7. Insomnia and sleep apnea are medical problems with real treatments.
- **If you have ADHD,** Chapter 8 is for you. Sleep problems are very common and often missed.

CHAPTER 1

Why sleep is the foundation

How much do adults need?

A consensus statement from the American Academy of Sleep Medicine and the Sleep Research Society recommends that adults sleep **7 or more hours per night** on a regular basis to promote optimal health. Sleeping less than 7 hours regularly is associated with adverse health outcomes, including weight gain, diabetes, high blood pressure, heart disease and impaired performance.[1] Individual needs vary a little. A useful test: if you need an alarm to wake up, nap easily in the day or rely on caffeine to function, you may be short.

Sleep and appetite

In a laboratory study of 12 healthy young men, two nights of 4-hour sleep reduced leptin (a hormone that signals fullness) by about 18% and increased ghrelin (a hunger hormone) by about 28%, and the men reported about a 24% increase in hunger and stronger appetite for calorie-dense foods.[2] The sample was small and the exposure extreme, but the direction matches what many people feel after a bad night: more hunger, more cravings.

Sleep and fat loss

In a small randomized crossover study, 10 overweight adults followed the same calorie-restricted diet for two weeks, once with 8.5 hours in bed and once with 5.5 hours. On both occasions they lost a similar amount of **weight,** but on the short-sleep arm they lost about **55% less fat** and about 60% more lean mass.[3] It was a small study, so do not treat the exact percentages as precise. The practical point stands: if you diet while short of sleep, you may lose a worse kind of weight.

Sleep and performance

In a study of Stanford basketball players, extending sleep to about 10 hours per night for five to seven weeks was associated with faster sprint times, better shooting accuracy and improved mood.[4] There was no control group, so some of this may be due to other factors. A review of sleep in athletes concluded more broadly that sleep loss impairs performance and recovery and that sleep extension is a promising strategy.[5]

What I removed

Many guides say that "you cannot out-train or out-diet bad sleep" and then quote percentages that nobody can source. The statement is a good rule of thumb, and I use it as coaching advice, but I do not attach fake precision to it.

CHAPTER 2

How sleep works (the short version)

Sleep is not one state. Over a night you move through repeating cycles of **light sleep, deep sleep (slow-wave) and REM sleep,** each lasting roughly 90 minutes on average, though this varies by person and by night.

Stage	What happens	Why it matters
Light sleep	Transition. Heart rate and temperature drop.	Roughly half the night. Easy to wake from.
Deep sleep	Slow brain waves. Hard to wake.	Physical restoration. More in the first half of the night.
REM sleep	Vivid dreams. Brain active, muscles relaxed.	Emotional processing and memory. More in the second half of the night.

Two systems decide when you sleep. **Sleep pressure** builds the longer you are awake (adenosine is part of this). Your **body clock** (circadian rhythm) sends alerting and sleepy signals on a roughly 24-hour schedule, anchored mostly by light. Good sleep happens when both line up: you are tired enough and it is the right time on the clock.

Why cutting sleep short hurts

Because REM sleep is concentrated in the last hours of the night, going to bed late and waking early with an alarm trims REM the most. It is one reason a short night can leave you irritable and flat, even if you "got through" the day.

CHAPTER 3

Set up the night

1. A consistent schedule

Go to bed and wake up at about the same time every day, within an hour or so, including weekends. This is a standard recommendation in sleep medicine, because a regular schedule stabilizes your body clock. If your schedule is irregular because of shifts or kids, the principle still holds: anchor **one** thing, usually the wake-up time.

2. Light: bright by day, dim by night

Light is the strongest signal for your body clock. A practical rule that matches the science of circadian rhythms:

- **Get outdoor light soon after waking,** ideally 10 to 20 minutes. Even a cloudy sky is far brighter than indoor lighting.

- **Dim the lights in the evening.** Lamps instead of bright overhead lights for the last 1 to 2 hours.
- **Dark room at night.** Blackout curtains or a sleep mask. Cover small LEDs.

In a controlled study, 12 adults who read on a light-emitting e-reader for several evenings before bed took longer to fall asleep, felt less sleepy at night, had **suppressed melatonin** and a later body clock, and were less alert the next morning compared with reading a printed book.[6] A device on a low brightness setting is a smaller problem than a bright one, and what you do on it matters too (a stressful email is worse than a book), so the sensible rule is: **screens off or very dim for the last hour,** and the phone charges outside the bedroom.

3. Cool, quiet, comfortable

Core body temperature falls as you go to sleep, and a room that is too warm makes that harder. A review of thermal environment and sleep found that heat exposure tends to increase wakefulness and reduce deep sleep and REM.[7] Many sleep guides suggest a bedroom around 16 to 19 degrees C (60 to 67 F). Treat that range as a starting point and adjust to what feels right for you.

Room setting	Aim	Cheap fix
Temperature	Cool, a little cooler than you think	Fan, lighter bedding, window open, lower thermostat
Light	As dark as possible	Blackout curtains, sleep mask, tape over LEDs
Noise	Quiet or steady	Earplugs, white noise or fan
Bed	Used for sleep and sex only	No work, no scrolling in bed

CHAPTER 4

Caffeine, alcohol, food and supplements

Caffeine: later than you think

Caffeine has a half-life of roughly 5 to 6 hours in many adults, so half of an afternoon dose is still active at bedtime. In a randomized crossover trial, 400 mg of caffeine taken 0, 3 or 6 hours before bedtime significantly reduced total sleep time, **even when taken 6 hours before bed**

(objectively about 40 minutes less sleep), and people did not always notice it.[8] A sensible rule: **stop caffeine at least 8 hours before bed,** and earlier if you are sensitive. For many people that means no coffee after about 1 to 2 PM.

You may have read that you should delay your first coffee for 90 to 120 minutes after waking so that adenosine clears. This is a popular idea, but I could not find human trials showing that it improves energy or sleep. If waiting suits you, fine, but it is not required. Do not feel guilty about a morning coffee.

Alcohol: a sedative, not a sleep aid

Alcohol may help you fall asleep faster, but a systematic review found that it reduces REM sleep in the first half of the night and disrupts sleep in the second half, with more waking and lighter sleep.[9] Higher doses have larger effects. If you drink, finish at least 3 to 4 hours before bed, drink water alongside, and keep it modest. If you are working on cutting alcohol out, expect your sleep to feel worse for a few weeks before it improves, which is common, and talk to your doctor if you were drinking heavily.

Food and timing

- **Avoid a very large meal right before bed.** It can cause discomfort and reflux.
- **Do not go to bed hungry** either. A small protein-rich snack is fine (Greek yogurt, a glass of milk).
- **Spread fluids across the day** so you are not waking for the toilet.

Melatonin: modest and specific

A meta-analysis of 19 studies in people with sleep problems found that melatonin reduced the time to fall asleep by about 7 minutes and increased total sleep by about 8 minutes, a small effect.[10] It can be helpful for shifting your body clock (jet lag, shift work, or a delayed schedule), generally in low doses taken at the right time. Supplement products vary in actual dose. It is not a substitute for fixing the habits above, and you should check with your doctor if you take other medication, are pregnant or have a medical condition.

A note on other sleep supplements

Magnesium, glycine, theanine, ashwagandha and others are popular. Evidence for most is limited or mixed. Habits first. If you want to try one, add it alone, for two weeks, and keep notes.

CHAPTER 5

The evening routine

Wind down on purpose

Your brain does not switch off like a lamp. Give it a runway of 30 to 60 minutes. The activity matters less than the pattern: the same cues, in the same order, signal "sleep is coming."

1. **60 minutes before bed:** screens off or very dim. Dim the lights. Do the tomorrow list on paper.
2. **45 minutes before:** a warm shower or bath, if you like. Brush teeth.
3. **30 minutes before:** something calm. A book, stretching, slow breathing, quiet music.
4. **At bedtime:** lights out, room cool and dark. If you cannot sleep within about 20 minutes, see the tip below.

A warm shower helps (a bit)

A meta-analysis found that a warm bath or shower (about 40 to 42.5 degrees C) taken **1 to 2 hours before bed** improved sleep quality and shortened the time to fall asleep by about 10 minutes.[11] The proposed reason is that the body cools after the heat, which helps trigger sleep. A very hot shower right before bed may do the opposite for some people.

Exercise helps sleep

A meta-analysis found that both single sessions and regular exercise improved sleep, with small to moderate effects, including less time to fall asleep and more deep sleep.[12] For most people, exercise at any time is better than none. If you find evening workouts hard to wind down from, move the hard sessions earlier and keep the evening gentle.

Racing mind? Write it down

If your mind races at bedtime, do a two-minute "brain dump": list what you are worried about and one next step for each. Then put the paper away. This is a coaching habit with limited direct evidence, but it costs two minutes.

If you cannot sleep

If you are still awake after about 20 minutes, get up and do something quiet in dim light (read a book, stretch) and go back when sleepy. Do not look at the clock. Staying in bed, frustrated, teaches your brain that bed means struggle.

CHAPTER 6

Daytime: light, naps and deep rest

Morning light

Outdoor light in the first hour or two after waking anchors your body clock. I recommend 10 minutes or more, no sunglasses unless needed for safety, never looking at the sun. On a dark winter morning, a bright lamp is a reasonable substitute.

Naps: short and early

A review of napping research found that a short nap (about 10 to 20 minutes) improved alertness and performance with little grogginess, while longer naps risk sleep inertia (feeling groggy).[13] Nap early in the afternoon, ideally before 3 PM, and skip naps if you have insomnia, since they can lower sleep pressure at night.

NSDR and yoga nidra: useful, but not a miracle

NSDR (non-sleep deep rest) is a modern name for guided relaxation practices such as yoga nidra. You lie down and follow a recording that takes you through breathing and body awareness for 10 to 30 minutes. Many people find it relaxing and it is a free tool for the afternoon dip.

You may have read that NSDR "increases dopamine by 65%" or "replaces hours of sleep." The 65% figure comes from a very small brain imaging study of **eight** experienced yoga nidra practitioners.[14] It is interesting, but it does not show that it works for you, and there is no good evidence that it replaces sleep. Use it for rest and to calm down, not as a replacement for a night's sleep.

Protocol

10 to 20 minutes, lying down, a free guided recording, once a day (for example, after lunch or after training). If you fall asleep, you probably needed it. Do not set expectations beyond "I feel calmer."

CHAPTER 7

When sleep does not improve

Insomnia is treatable, and it is not just "bad habits"

If you have trouble falling or staying asleep at least three nights a week for three months or more, and it affects your day, you may have insomnia disorder. The first-line treatment is **cognitive behavioral therapy for insomnia (CBT-I),** not sleeping pills. The American College of Physicians recommends CBT-I as the initial treatment for chronic insomnia in adults.[15] A meta-analysis of 20 randomized trials found that CBT-I improved insomnia severity, sleep onset latency and total sleep time in adults, with durable effects.[16]

CBT-I component	What it is
Sleep restriction	Temporarily limit time in bed to your actual sleep time, then widen it. Do this with guidance.
Stimulus control	Bed only for sleep. Get up if awake more than about 20 minutes.
Cognitive work	Challenge unhelpful sleep beliefs ("I will be useless tomorrow").
Relaxation and routine	Wind-down habits, breathing, a fixed schedule.

CBT-I is delivered by therapists, by programs and by apps, and is widely available online. Ask your doctor.

Signs to see a doctor

- Loud snoring, gasping, or pauses in breathing at night, or morning headaches. These can mean **sleep apnea,** which is common, serious and treatable.
- Falling asleep at the wheel or in dangerous situations.
- Restless legs or kicking at night.

- Insomnia that lasts weeks despite good habits, or that comes with low mood or severe anxiety.
- You rely on alcohol or over-the-counter pills to sleep every night.

A caution about sleeping pills

Prescription and over-the-counter sleeping pills may help in the short term but carry risks and are not first-line treatment for long-term insomnia. Do not start or stop prescribed sleep medication without your doctor.

CHAPTER 8

ADHD and sleep

Sleep problems are very common in people with ADHD. A review of the research found that adults and children with ADHD more often have difficulty falling asleep, a delayed body clock (evening-type), restless sleep and daytime sleepiness, and that other sleep disorders such as sleep apnea and restless legs syndrome are also more common.[17] Sleep loss, in turn, can look like ADHD and make real ADHD symptoms worse, so it is worth treating on its own.

Common ADHD sleep problem	What to try
Cannot switch off at night	Fixed wind-down with an alarm to start it. Brain dump on paper. Calm audio rather than video.
Bedtime procrastination ("revenge" scrolling)	Phone charges outside the room. Delete the app that drags you in. Treat bedtime as a start time for the routine, not lights out.
Late body clock	Fixed wake time every day. Morning light. Ask your doctor about low-dose, well-timed melatonin.
Medication timing	Stimulants can affect sleep. Talk to your prescriber about timing and dose, never change it alone.
Hard to wake up	Alarm across the room. Light on a timer. A planned reward (coffee, shower) in the first 10 minutes.

From my experience

As an adult with ADHD, I learned that I do better when the evening has fewer decisions. The same three or four steps, in the same order, with phone alarms to start and finish them. Your version will be different, but fewer decisions help most brains.

CHAPTER 9

The stress equation and recovery

Load versus capacity

A simple model I use with clients: **stress load** (work, training, family, money, emotions) against **recovery capacity** (sleep, food, rest days, social support, coping tools). When load stays above capacity for too long, you tend to see poor sleep, irritability, low motivation, a higher resting heart rate, more injuries and more illness. This is a practical model from coaching, not a laboratory formula, but it is useful because it gives you two levers: reduce the load, or raise the capacity.

Signs you are over capacity	What to do this week
Sleep worse than usual for several nights	Cut one commitment. Protect bedtime.
Resting heart rate noticeably above your normal for days	Reduce training volume 30 to 50% for a week.
Irritable, low mood, no drive	Add a rest day and a walk. Talk to someone.
Performance dropping despite effort	Deload. Eat enough. Sleep more. See a doctor if it lasts.
Frequent colds or small injuries	Back off, eat and sleep, and check with a doctor if it persists.

Raise capacity every day

Frequency	Habit
Daily	7+ hours of sleep window. A 20 to 30 minute walk outdoors. One or two real conversations. 5 to 10 minutes of calm (breathing, NSDR, meditation, prayer, music).

Weekly	At least one full rest day from hard training. One easy recovery session. One social plan. Time in nature.
Monthly	A lighter week every 4 to 8 weeks. A review of your goals and workload.

Active recovery

Active recovery is low-intensity movement on easy days: walking, easy cycling, swimming, mobility work. A useful guide is the **talk test:** you can hold a conversation. Some coaches use a heart rate under about 120 beats per minute as a rule of thumb, but this depends on the person. Easy movement can ease stiffness and feel good, and it should leave you feeling better, not more tired. It is not a license to turn every rest day into a workout.

Deloads and rest days

Hard training breaks you down, and adaptation happens during recovery. Plan an easier week every 4 to 8 weeks, reduce volume by about 30 to 50%, and keep the movements. If you are in CrossFit or similar high-intensity training, build rest days into the week rather than waiting until you are forced to take them.

CHAPTER 10

Building resilience

Resilience is the ability to handle stress and recover from setbacks. It is not a fixed trait. It comes from a set of skills and supports that you can build: managing your emotions (see Guide 4), seeing things flexibly, keeping your body healthy, and having people to lean on.

Reframe the stress response

In an experiment, participants who were told that the physical arousal they felt before a stressful task (a fast heartbeat, say) can help performance showed better stress responses and performed better than those who were told to ignore it.[18] Results of this approach have been mixed in later work, so treat it as a promising tool: "My body is getting ready" can be a better script than "I must calm down at once."

Journaling

Writing about emotions is covered in Guide 4. The research shows a **small** benefit on average, so use it as a cheap tool, not a cure.[19] A simple structure: morning, 3 minutes (one priority, one worry and a next step). Evening, 3 minutes (what went well, one thing to try tomorrow, one word for how you feel).

Social connection

A meta-analysis of 148 studies (over 300,000 people) found that people with stronger social relationships had about a **50% greater likelihood of survival** over the follow-up period than those with weaker ones, an effect comparable to well-known risk factors such as smoking.[20] Most of these studies are observational, so cause and effect are not proven, and the often-quoted "as bad as 15 cigarettes a day" is a loose way of expressing the comparison. The practical advice is simple: keep a few real relationships alive. One or two real conversations a day, one social plan a week.

Noticing: a mindfulness without the cushion

Psychologist Ellen Langer defines mindfulness as actively noticing new things, which she argues reduces autopilot and rigidity.[21] You can practice it without sitting: during one meal a day, notice three details of taste or texture; on a regular walk, notice one new sound; in a talk with someone you know well, notice something new. Evidence for this version is more limited than for formal mindfulness programs (see Guide 4), but it costs nothing and makes ordinary moments more vivid.

Taking in the good

Our brains register threats quickly and good moments weakly. A popular practice, "taking in the good," asks you to stay with a positive experience for 10 to 20 seconds so it registers. The idea is plausible and it is a pleasant habit, but I am not aware of strong trials. Try it as an evening ritual at the end of your journaling and judge by how it feels.

Cold exposure: what we know

Cold showers and ice baths are everywhere, and so are big numbers. One often-quoted study found that immersion in cold water (14 degrees C) raised dopamine by 250%, but it had **six** participants and measured blood levels, not brain activity.[22] A large Dutch randomized trial of 3,018 adults found that ending the daily shower with 30 to 90 seconds of cold water reduced

sickness absence from work by 29%, but did not reduce the number of days people reported being ill.[23] That suggests cold may change how people feel or cope, not necessarily their immune function.

If you like cold showers and they make you feel alert, there is no reason to stop. Do not use them as a substitute for sleep, avoid prolonged cold-water immersion alone, and check with a doctor if you have heart or blood pressure problems. Do not do breath-holding in water.

Be cautious with hacks

If a recovery tool promises a huge, precise effect ("+250%," "+65%," "replaces 2 hours of sleep"), ask: how many people were studied, and what was actually measured? If the answer is a handful of participants and a blood marker, treat it as a hypothesis.

CHAPTER 11

A recovery day, stitched together

Here is what a good day can look like. Nothing here takes more than 30 to 40 extra minutes. The lift is consistency, not time. Adjust the clock to your life.

Time	What
6:30	Wake at your usual time. Outside for 10 minutes. Water.
7:00 to 8:00	Breakfast with protein (Guide 2). Coffee if you like.
13:00	Lunch, a short walk. Optional 10 to 20 minute NSDR or nap if you need it.
14:00	Last caffeine of the day.
17:30	Training or easy movement. Hard sessions earlier, if you can.
20:00	Dinner finished. Lights dim. Phone on the charger outside the bedroom.
21:00	Warm shower. Three-line journal. Brain dump for tomorrow.
21:30 to 22:00	Quiet reading, breathing. Bed. Room cool and dark.

CHAPTER 12

Your 14-day plan

Days	Change	Why this order
1 to 3	Fix your **wake-up time** and get 10 minutes of outdoor light	Anchors your body clock. Costs nothing.
4 to 6	**Caffeine cutoff** 8 hours before bed. Phone charges outside the bedroom	Biggest sleep gains for most people.
7 to 9	**Wind-down routine:** dim lights, screens off for the last hour, brain dump	Makes sleep come easier.
10 to 12	**Room:** cool, dark, quiet. Alcohol off on weeknights	Improves sleep quality.
13 to 14	Review: sleep log, mood, energy. Decide what stays	You keep what works.

Keep a simple log for the two weeks: bedtime, wake time, how you slept (1 to 5), energy at 3 PM (1 to 5). Do not obsess over numbers, since worrying about sleep can make it worse. The pattern over two weeks is what counts.

CHAPTER 13

Pick your case: starter plans

If this is you…	Your starter plan
I sleep 5 to 6 hours	Move bedtime 15 minutes earlier every 3 days until you reach 7 or more hours. Do not try to jump an hour at once.
I fall asleep fine but wake at 3 AM	Check alcohol, caffeine and room temperature. Avoid looking at the clock. If it persists, see your doctor (apnea, stress, mood).
I cannot switch off	Brain dump, breathing, calm audio. Consider CBT-I (Chapter 7).
I work shifts	Anchor sleep to one fixed block. Blackout room. Light at wake-up. Caffeine early in the shift. Talk to your doctor about melatonin timing.

I am a new parent or caregiver	Protect any 4 to 5 hour block. Nap when you can. Lower training load. Accept this is a season.
I train hard and feel flat	Deload week. Eat enough. Sleep 8 hours for 7 days. See a doctor if it lasts.
I am stopping alcohol	Expect rough sleep for weeks. Keep the schedule. Light and exercise help. Medical advice if drinking was heavy.

APPENDIX

Worksheets

1. Two-week sleep log

Date	Bedtime	Wake	Sleep quality 1 to 5	Energy 3 PM 1 to 5	Caffeine last at

2. My evening routine

Time	Step

3. Load and capacity check (weekly)

Question	My answer
What is draining me most this week?	

What is my best recovery habit?
What can I remove or reduce?
Who will I talk to this week?
Rest day planned? When?

About the author

Adriano Monteiro is a nutrition coach with Precision Nutrition certifications. He has lost 38 kg of fat, has competed in CrossFit for 8 years, has been sober for 4 years and has been in treatment for ADHD for 2 years. He writes and coaches for people who want lasting change without having to suffer in silence, and works with clients in English from Brazil.

Want a sleep and recovery plan built around YOUR week?

Book a free 20-minute call. We will look at your schedule, your training and your stress, and choose the first change to make. No pressure, no sales pitch, just a plan.

Book your free 20-minute call

https://calendly.com/adriano-nutrition/free-20-min-discovery-call

References

1. Watson, N. F., Badr, M. S., Belenky, G., et al. (2015). Recommended amount of sleep for a healthy adult: A joint consensus statement of the American Academy of Sleep Medicine and Sleep Research Society. *Sleep*, 38(6), 843-844.
2. Spiegel, K., Tasali, E., Penev, P., & Van Cauter, E. (2004). Brief communication: Sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Annals of Internal Medicine*, 141(11), 846-850.
3. Nedeltcheva, A. V., Kilkus, J. M., Imperial, J., et al. (2010). Insufficient sleep undermines dietary efforts to reduce adiposity. *Annals of Internal Medicine*, 153(7), 435-441.
4. Mah, C. D., Mah, K. E., Kezirian, E. J., & Dement, W. C. (2011). The effects of sleep extension on the athletic performance of collegiate basketball players. *Sleep*, 34(7), 943-950.
5. Fullagar, H. H. K., Skorski, S., Duffield, R., et al. (2015). Sleep and athletic performance: The effects of sleep loss on exercise performance, and physiological and cognitive responses to exercise. *Sports Medicine*, 45(2), 161-186.
6. Chang, A. M., Aeschbach, D., Duffy, J. F., & Czeisler, C. A. (2015). Evening use of light-emitting eReaders negatively affects sleep, circadian timing, and next-morning alertness. *Proceedings of the National Academy of Sciences*, 112(4), 1232-1237.
7. Okamoto-Mizuno, K., & Mizuno, K. (2012). Effects of thermal environment on sleep and circadian rhythm. *Journal of Physiological Anthropology*, 31, 14.
8. Drake, C., Roehrs, T., Shambroom, J., & Roth, T. (2013). Caffeine effects on sleep taken 0, 3, or 6 hours before going to bed. *Journal of Clinical Sleep Medicine*, 9(11), 1195-1200.
9. Ebrahim, I. O., Shapiro, C. M., Williams, A. J., & Fenwick, P. B. (2013). Alcohol and sleep I: Effects on normal sleep. *Alcoholism: Clinical and Experimental Research*, 37(4), 539-549.
10. Ferracioli-Oda, E., Qawasmi, A., & Bloch, M. H. (2013). Meta-analysis: Melatonin for the treatment of primary sleep disorders. *PLoS ONE*, 8(5), e63773.
11. Haghayegh, S., Khoshnevis, S., Smolensky, M. H., Diller, K. R., & Castriotta, R. J. (2019). Before-bedtime passive body heating by warm shower or bath to improve sleep: A systematic review and meta-analysis. *Sleep Medicine Reviews*, 46, 124-135.
12. Kredlow, M. A., Capozzoli, M. C., Hearon, B. A., Calkins, A. W., & Otto, M. W. (2015). The effects of physical activity on sleep: A meta-analytic review. *Journal of Behavioral Medicine*, 38(3), 427-449.
13. Milner, C. E., & Cote, K. A. (2009). Benefits of napping in healthy adults: Impact of nap length, time of day, age, and experience with napping. *Journal of Sleep Research*, 18(2), 272-281.
14. Kjaer, T. W., Bertelsen, C., Piccini, P., et al. (2002). Increased dopamine tone during meditation-induced change of consciousness. *Cognitive Brain Research*, 13(2), 255-259.
15. Qaseem, A., Kansagara, D., Forciea, M. A., Cooke, M., & Denberg, T. D. (2016). Management of chronic insomnia disorder in adults: A clinical practice guideline from the American College of Physicians. *Annals of Internal Medicine*, 165(2), 125-133.
16. Trauer, J. M., Qian, M. Y., Doyle, J. S., Rajaratnam, S. M. W., & Cunnington, D. (2015). Cognitive behavioral therapy for chronic insomnia: A systematic review and meta-analysis. *Annals of Internal Medicine*, 163(3), 191-204.
17. Hvolby, A. (2015). Associations of sleep disturbance with ADHD: Implications for treatment. *ADHD Attention Deficit and Hyperactivity Disorders*, 7(1), 1-18.
18. Jamieson, J. P., Nock, M. K., & Mendes, W. B. (2012). Mind over matter: Reappraising arousal improves cardiovascular and cognitive responses to stress. *Journal of Experimental Psychology: General*, 141(3), 417-422.

19. Frattaroli, J. (2006). Experimental disclosure and its moderators: A meta-analysis. *Psychological Bulletin*, 132(6), 823-865.

20. Holt-Lunstad, J., Smith, T. B., & Layton, J. B. (2010). Social relationships and mortality risk: A meta-analytic review. *PLoS Medicine*, 7(7), e1000316.

21. Langer, E. J., & Moldoveanu, M. (2000). The construct of mindfulness. *Journal of Social Issues*, 56(1), 1-9.

22. Sramek, P., Simeckova, M., Jansky, L., Savlikova, J., & Vybiral, S. (2000). Human physiological responses to immersion into water of different temperatures. *European Journal of Applied Physiology*, 81(5), 436-442.

23. Buijze, G. A., Sierevelt, I. N., van der Heijden, B. C. J. M., Dijkgraaf, M. G., & Frings-Dresen, M. H. W. (2016). The effect of cold showering on health and work: A randomized controlled trial. *PLoS ONE*, 11(9), e0161749.